

On Some Properties of Real Hilbert Space. Part II

Hiroshi Yamazaki
Shinshu University
Nagano

Yasumasa Suzuki
Take, Yokosuka-shi
Japan

Takao Inoué
The Iida Technical High School
Nagano

Yasunari Shidama
Shinshu University
Nagano

Summary. This paper is a continuation of our paper [22]. We give an analogue of the necessary and sufficient condition for summable set (i.e. the main theorem of [22]) with respect to summable set by a functional L in real Hilbert space. After presenting certain useful lemmas, we prove our main theorem that the summability for an orthonormal infinite set in real Hilbert space is equivalent to its summability with respect to the square of norm, say $H(x) = (x, x)$. Then we show that the square of norm H commutes with infinite sum operation if the orthonormal set under our consideration is summable. Our main theorem is due to [8].

MML Identifier: BHSP_7.

WWW: http://mizar.org/JFM/Vol15/bhsp_7.html

The articles [16], [19], [6], [1], [17], [9], [4], [5], [20], [18], [12], [13], [14], [3], [7], [10], [15], [11], [2], [21], and [22] provide the notation and terminology for this paper.

1. NECESSARY AND SUFFICIENT CONDITION FOR SUMMABLE SET

In this paper X is a real unitary space and x, y are points of X .

The following propositions are true:

- (1) Let Y be a subset of X and L be a functional in X . Then Y is summable set by L if and only if for every real number e such that $0 < e$ there exists a finite subset Y_0 of X such that Y_0 is non empty and $Y_0 \subseteq Y$ and for every finite subset Y_1 of X such that Y_1 is non empty and $Y_1 \subseteq Y$ and Y_0 misses Y_1 holds $|\text{setofunc}(Y_1, \text{the carrier of } X, \mathbb{R}, L, +\mathbb{R})| < e$.
- (2) Let given X . Suppose the addition of X is commutative and associative and the addition of X has a unity. Let S be a finite orthogonal family of X . Suppose S is non empty. Let I be a function from the carrier of X into the carrier of X . Suppose $S \subseteq \text{dom } I$ and for every y such that $y \in S$ holds $I(y) = y$. Let H be a function from the carrier of X into \mathbb{R} . Suppose $S \subseteq \text{dom } H$ and for every y such that $y \in S$ holds $H(y) = (y|y)$. Then $(\text{setofunc}(S, \text{the carrier of } X, \text{the carrier of } X, I, \text{the addition of } X) | \text{setofunc}(S, \text{the carrier of } X, \text{the carrier of } X, I, \text{the addition of } X)) = \text{setofunc}(S, \text{the carrier of } X, \mathbb{R}, H, +\mathbb{R})$.
- (3) Let given X . Suppose the addition of X is commutative and associative and the addition of X has a unity. Let S be a finite orthogonal family of X . Suppose S is non empty. Let H be a functional in X . Suppose $S \subseteq \text{dom } H$ and for every x such that $x \in S$ holds $H(x) = (x|x)$. Then $(\text{Setsum}(S) | \text{Setsum}(S)) = \text{setofunc}(S, \text{the carrier of } X, \mathbb{R}, H, +\mathbb{R})$.

- (4) Let Y be an orthogonal family of X and Z be a subset of X . If Z is a subset of Y , then Z is an orthogonal family of X .
- (5) Let Y be an orthonormal family of X and Z be a subset of X . If Z is a subset of Y , then Z is an orthonormal family of X .

2. EQUIVALENCE OF SUMMABILITY

One can prove the following propositions:

- (6) Let given X . Suppose the addition of X is commutative and associative and the addition of X has a unity and X is a Hilbert space. Let S be an orthonormal family of X and H be a functional in X . Suppose $S \subseteq \text{dom}H$ and for every x such that $x \in S$ holds $H(x) = (x|x)$. Then S is summable_set if and only if S is summable set by H .
- (7) Let S be a subset of X . Suppose S is non empty and summable_set. Let e be a real number. Suppose $0 < e$. Then there exists a finite subset Y_0 of X such that Y_0 is non empty and $Y_0 \subseteq S$ and for every finite subset Y_1 of X such that $Y_0 \subseteq Y_1$ and $Y_1 \subseteq S$ holds $|(\text{sum}S | \text{sum}S) - (\text{Setsum}(Y_1) | \text{Setsum}(Y_1))| < e$.
- (8) Let given X . Suppose the addition of X is commutative and associative and the addition of X has a unity and X is a Hilbert space. Let S be an orthonormal family of X . Suppose S is non empty. Let H be a functional in X . Suppose $S \subseteq \text{dom}H$ and for every x such that $x \in S$ holds $H(x) = (x|x)$. If S is summable_set, then $(\text{sum}S | \text{sum}S) = \text{SumByfunc}(S, H)$.

REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinal1.html>.
- [2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.
- [3] Czesław Byliński. Binary operations. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/binop_1.html.
- [4] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [5] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [6] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [7] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.
- [8] P. R. Halmos. *Introduction to Hilbert Space*. American Mathematical Society, 1987.
- [9] Krzysztof Hryniewiecki. Basic properties of real numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/real_1.html.
- [10] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/vectsp_1.html.
- [11] Bogdan Nowak and Andrzej Trybulec. Hahn-Banach theorem. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/hahnban.html>.
- [12] Jan Popiolek. Some properties of functions modul and signum. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/absvalue.html>.
- [13] Jan Popiolek. Introduction to Banach and Hilbert spaces — part I. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/bhsp_1.html.
- [14] Jan Popiolek. Introduction to Banach and Hilbert spaces — part III. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/bhsp_3.html.
- [15] Andrzej Trybulec. Semilattice operations on finite subsets. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/setwiseo.html>.
- [16] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.

- [17] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [18] Wojciech A. Trybulec. Vectors in real linear space. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/rlvect_1.html.
- [19] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/subset_1.html.
- [20] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/relat_1.html.
- [21] Hiroshi Yamazaki, Yasunari Shidama, and Yatsuka Nakamura. Bessel's inequality. *Journal of Formalized Mathematics*, 15, 2003. http://mizar.org/JFM/Voll15/bhsp_5.html.
- [22] Hiroshi Yamazaki, Yasumasa Suzuki, Takao Inoué, and Yasunari Shidama. On some properties of real Hilbert space, I. *Journal of Formalized Mathematics*, 15, 2003. http://mizar.org/JFM/Voll15/bhsp_6.html.

Received April 17, 2003

Published January 2, 2004
