## **Cardinal Arithmetics**

Grzegorz Bancerek Warsaw University Białystok

**Summary.** In the article addition, multiplication and power operation of cardinals are introduced. Presented are some properties of equipotence of Cartesian products, basic cardinal arithmetics laws (transformativity, associativity, distributivity), and some facts about finite sets.

MML Identifier: CARD\_2.
WWW: http://mizar.org/JFM/Vol2/card\_2.html

The articles [13], [12], [9], [14], [7], [8], [3], [4], [5], [11], [2], [6], [1], and [10] provide the notation and terminology for this paper.

For simplicity, we adopt the following convention: A, B are ordinal numbers, K, M, N are cardinal numbers, x,  $x_1$ ,  $x_2$ , y,  $y_1$ ,  $y_2$ , X, Y, Z,  $X_1$ ,  $X_2$ ,  $Y_1$ ,  $Y_2$  are sets, and f is a function.

Next we state several propositions:

- (2)<sup>1</sup>  $\overline{\overline{X}} \leq \overline{\overline{Y}}$  iff there exists f such that  $X \subseteq f^{\circ}Y$ .
- (3)  $\overline{\overline{f^{\circ}X}} \leq \overline{\overline{X}}$ .
- (4) If  $\overline{\overline{X}} < \overline{\overline{Y}}$ , then  $Y \setminus X \neq \emptyset$ .
- (5) If  $x \in X$  and  $X \approx Y$ , then  $Y \neq \emptyset$  and there exists x such that  $x \in Y$ .
- (6)  $2^X \approx 2^{\overline{X}}$  and  $\overline{\overline{2^X}} = \overline{\overline{2^{\overline{X}}}}$ .
- (7) If  $Z \in Y^X$ , then  $Z \approx X$  and  $\overline{\overline{Z}} = \overline{\overline{X}}$ .

Let us consider M, N. The functor M + N yielding a cardinal number is defined as follows:

(Def. 1) 
$$M + N = \overline{M + N}$$
.

Let us observe that the functor M + N is commutative. The functor  $M \cdot N$  yields a cardinal number and is defined as follows:

(Def. 2)  $M \cdot N = \overline{[:M,N:]}$ .

Let us observe that the functor  $M \cdot N$  is commutative. The functor  $M^N$  yields a cardinal number and is defined as follows:

(Def. 3)  $M^N = \overline{\overline{M^N}}$ .

The following propositions are true:

<sup>&</sup>lt;sup>1</sup> The proposition (1) has been removed.

- $(11)^2$  [:X, Y:]  $\approx$  [:Y, X:] and  $\overline{\overline{[:X,Y:]}} = \overline{\overline{[:Y,X:]}}$ .
- (12)  $[:[:X,Y:],Z:] \approx [:X,[:Y,Z:]:]$  and  $\overline{\overline{[:[:X,Y:],Z:]}} = \overline{\overline{[:X,[:Y,Z:]:]}}$ .
- (13)  $X \approx [:X, \{x\}:]$  and  $\overline{\overline{X}} = \overline{[:X, \{x\}:]}$ .
- (14)  $[:X,Y:] \approx [:\overline{\overline{X}},Y:]$  and  $[:X,Y:] \approx [:X,\overline{\overline{Y}}:]$  and  $[:X,Y:] \approx [:\overline{\overline{X}},\overline{\overline{Y}}:]$  and  $\overline{\overline{[:X,Y:]}} = \overline{\overline{[:\overline{\overline{X}},Y:]}}$ and  $\overline{\overline{[:X,Y:]}} = \overline{\overline{[:X,\overline{\overline{Y}}:]}}$  and  $\overline{\overline{[:X,Y:]}} = \overline{\overline{[:\overline{\overline{X}},\overline{\overline{Y}}:]}}$ .
- (15) If  $X_1 \approx Y_1$  and  $X_2 \approx Y_2$ , then  $[X_1, X_2:] \approx [Y_1, Y_2:]$  and  $\overline{[X_1, X_2:]} = \overline{[Y_1, Y_2:]}$ .
- (16) If  $x_1 \neq x_2$ , then  $A + B \approx [:A, \{x_1\}:] \cup [:B, \{x_2\}:]$  and  $\overline{\overline{A + B}} = \overline{[:A, \{x_1\}:] \cup [:B, \{x_2\}:]}$ .
- (17) If  $x_1 \neq x_2$ , then  $K + M \approx [:K, \{x_1\}:] \cup [:M, \{x_2\}:]$  and  $K + M = \overline{[:K, \{x_1\}:] \cup [:M, \{x_2\}:]}$ .
- (18)  $A \cdot B \approx [:A, B:]$  and  $\overline{\overline{A \cdot B}} = \overline{[:A, B:]}$ .
- (19)  $0 = \overline{\overline{0}}$  and  $1 = \overline{\overline{1}}$  and  $2 = \overline{\overline{\operatorname{succ} 1}}$ .
- (20) 1 = 1.
- $(22)^3$  2 = {0,1} and 2 = succ 1.
- (23) If  $X_1 \approx Y_1$  and  $X_2 \approx Y_2$  and  $x_1 \neq x_2$  and  $y_1 \neq y_2$ , then  $[:X_1, \{x_1\}:] \cup [:X_2, \{x_2\}:] \approx [:Y_1, \{y_1\}:] \cup [:Y_2, \{y_2\}:]$  and  $\overline{[:X_1, \{x_1\}:] \cup [:X_2, \{x_2\}:]} = \overline{[:Y_1, \{y_1\}:] \cup [:Y_2, \{y_2\}:]}$ .
- (24)  $\overline{\overline{A+B}} = \overline{\overline{A}} + \overline{\overline{B}}.$
- (25)  $\overline{\overline{A \cdot B}} = \overline{\overline{A}} \cdot \overline{\overline{B}}.$
- $(26) [X, \{0\}] \cup [Y, \{1\}] \approx [Y, \{0\}] \cup [X, \{1\}] \text{ and } \overline{[X, \{0\}] \cup [Y, \{1\}]} = \overline{[Y, \{0\}] \cup [X, \{1\}]}.$
- (27)  $[:X_1, X_2:] \cup [:Y_1, Y_2:] \approx [:X_2, X_1:] \cup [:Y_2, Y_1:] \text{ and } \overline{[:X_1, X_2:] \cup [:Y_1, Y_2:]} = \overline{[:X_2, X_1:] \cup [:Y_2, Y_1:]}.$
- (28) If  $x \neq y$ , then  $\overline{\overline{X}} + \overline{\overline{Y}} = \overline{[:X, \{x\}:] \cup [:Y, \{y\}:]}$ .
- $(29) \quad M+0=M.$
- $(31)^4$  (K+M)+N=K+(M+N).
- $(32) \quad K \cdot 0 = 0.$
- $(33) \quad K \cdot 1 = K.$
- $(35)^5 \quad (K \cdot M) \cdot N = K \cdot (M \cdot N).$
- $(36) \quad 2 \cdot K = K + K.$
- $(37) \quad K \cdot (M+N) = K \cdot M + K \cdot N.$
- (38)  $K^0 = 1.$
- (39) If  $K \neq 0$ , then  $0^K = 0$ .
- (40)  $K^1 = K$  and  $1^K = 1$ .
- $(41) \quad K^{M+N} = K^M \cdot K^N.$

<sup>&</sup>lt;sup>2</sup> The propositions (8)–(10) have been removed.

<sup>&</sup>lt;sup>3</sup> The proposition (21) has been removed.

<sup>&</sup>lt;sup>4</sup> The proposition (30) has been removed.

<sup>&</sup>lt;sup>5</sup> The proposition (34) has been removed.

- (42)  $(K \cdot M)^N = K^N \cdot M^N$ .
- $(43) \quad K^{M \cdot N} = (K^M)^N.$
- (44)  $2^{\overline{X}} = \overline{\overline{2^X}}.$
- $(45) \quad K^2 = K \cdot K.$
- (46)  $(K+M)^2 = K \cdot K + 2 \cdot K \cdot M + M \cdot M.$
- (47)  $\overline{\overline{X \cup Y}} \leq \overline{\overline{X}} + \overline{\overline{Y}}.$
- (48) If X misses Y, then  $\overline{\overline{X \cup Y}} = \overline{\overline{X}} + \overline{\overline{Y}}$ .

In the sequel *m*, *n* are natural numbers. The following propositions are true:

- $(49) \quad n+m=n+m.$
- (50)  $n \cdot m = n \cdot m$ .
- (51)  $\overline{\overline{n+m}} = \overline{\overline{n}} + \overline{\overline{m}}.$
- (52)  $\overline{\overline{n \cdot m}} = \overline{\overline{n}} \cdot \overline{\overline{m}}.$
- (53) For all finite sets X, Y such that X misses Y holds  $card(X \cup Y) = cardX + cardY$ .
- (54) For every finite set *X* such that  $x \notin X$  holds  $\operatorname{card}(X \cup \{x\}) = \operatorname{card} X + 1$ .
- (57)<sup>6</sup> For all finite sets X, Y holds  $\overline{\overline{X}} \leq \overline{\overline{Y}}$  iff card  $X \leq$  card Y.
- (58) For all finite sets *X*, *Y* holds  $\overline{\overline{X}} < \overline{\overline{Y}}$  iff card *X* < card *Y*.
- (59) For every set X such that  $\overline{\overline{X}} = 0$  holds  $X = \emptyset$ .
- (60) For every set *X* holds  $\overline{\overline{X}} = 1$  iff there exists *x* such that  $X = \{x\}$ .
- (61) For every finite set X holds  $X \approx \operatorname{card} X$  and  $X \approx \overline{\operatorname{card} X}$  and  $X \approx \operatorname{Seg} \operatorname{card} X$ .
- (62) For all finite sets X, Y holds  $\operatorname{card}(X \cup Y) \leq \operatorname{card} X + \operatorname{card} Y$ .
- (63) For all finite sets X, Y such that  $Y \subseteq X$  holds  $\operatorname{card}(X \setminus Y) = \operatorname{card} X \operatorname{card} Y$ .
- (64) For all finite sets X, Y holds  $\operatorname{card}(X \cup Y) = (\operatorname{card} X + \operatorname{card} Y) \operatorname{card}(X \cap Y)$ .
- (65) For all finite sets X, Y holds  $\operatorname{card}[:X, Y:] = \operatorname{card} X \cdot \operatorname{card} Y$ .
- (67)<sup>7</sup> For all finite sets X, Y such that  $X \subset Y$  holds card X < card Y and  $\overline{\overline{X}} < \overline{\overline{Y}}$ .
- (68) If  $\overline{\overline{X}} \leq \overline{\overline{Y}}$  or  $\overline{\overline{X}} < \overline{\overline{Y}}$  and if Y is finite, then X is finite.

In the sequel  $x_1$ ,  $x_2$ ,  $x_3$ ,  $x_4$ ,  $x_5$ ,  $x_6$ ,  $x_7$ ,  $x_8$  denote sets. Next we state a number of propositions:

- (69) card{ $x_1, x_2$ }  $\leq 2$ .
- (70) card{ $x_1, x_2, x_3$ }  $\leq 3$ .
- (71) card{ $x_1, x_2, x_3, x_4$ }  $\leq 4$ .
- (72) card{ $x_1, x_2, x_3, x_4, x_5$ }  $\leq 5$ .

<sup>&</sup>lt;sup>6</sup> The propositions (55) and (56) have been removed.

<sup>&</sup>lt;sup>7</sup> The proposition (66) has been removed.

- (73) card{ $x_1, x_2, x_3, x_4, x_5, x_6$ }  $\leq 6$ .
- (74) card{ $x_1, x_2, x_3, x_4, x_5, x_6, x_7$ }  $\leq 7$ .
- (75) card{ $x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8$ }  $\leq 8$ .
- (76) If  $x_1 \neq x_2$ , then card $\{x_1, x_2\} = 2$ .
- (77) If  $x_1 \neq x_2$  and  $x_1 \neq x_3$  and  $x_2 \neq x_3$ , then card $\{x_1, x_2, x_3\} = 3$ .
- (78) If  $x_1 \neq x_2$  and  $x_1 \neq x_3$  and  $x_1 \neq x_4$  and  $x_2 \neq x_3$  and  $x_2 \neq x_4$  and  $x_3 \neq x_4$ , then card $\{x_1, x_2, x_3, x_4\} = 4$ .

## REFERENCES

- [1] Grzegorz Bancerek. Cardinal numbers. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/card\_1.html.
- [2] Grzegorz Bancerek. The fundamental properties of natural numbers. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/nat\_1.html.
- [3] Grzegorz Bancerek. The ordinal numbers. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/ordinal1. html.
- [4] Grzegorz Bancerek. Sequences of ordinal numbers. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/ ordinal2.html.
- [5] Grzegorz Bancerek. Zermelo theorem and axiom of choice. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/ Voll/wellord2.html.
- [6] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/finseq\_1.html.
- [7] Czesław Byliński. Functions and their basic properties. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/ funct\_1.html.
- [8] Czesław Byliński. Functions from a set to a set. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/funct\_ 2.html.
- [9] Czesław Byliński. Some basic properties of sets. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/ zfmisc\_1.html.
- [10] Agata Darmochwał. Finite sets. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/finset\_1.html.
- [11] Krzysztof Hryniewiecki. Basic properties of real numbers. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/ Voll/real\_1.html.
- [12] Andrzej Trybulec. Enumerated sets. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/enumset1.html.
- [13] Andrzej Trybulec. Tarski Grothendieck set theory. Journal of Formalized Mathematics, Axiomatics, 1989. http://mizar.org/JFM/ Axiomatics/tarski.html.
- [14] Edmund Woronowicz. Relations and their basic properties. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/ Voll/relat\_1.html.

Received March 6, 1990

Published January 2, 2004