

Coherent Space

Jarosław Kotowicz
Warsaw University
Białystok

Konrad Raczkowski
Warsaw University
Białystok

Summary. Coherent Space, web of coherent space and two categories: category of coherent spaces and category of tolerances on same fixed set.

MML Identifier: COH_SP.

WWW: http://mizar.org/JFM/Vol4/coh_sp.html

The articles [9], [6], [11], [12], [13], [10], [2], [5], [1], [3], [8], [7], and [4] provide the notation and terminology for this paper.

1. COHERENT SPACE AND WEB OF COHERENT SPACE

In this paper x, y, a, b, X, A denote sets.

Let I_1 be a set. We say that I_1 is binary complete if and only if:

(Def. 2)¹ For every A such that $A \subseteq I_1$ and for all a, b such that $a \in A$ and $b \in A$ holds $a \cup b \in I_1$ holds $\bigcup A \in I_1$.

Let us note that there exists a set which is subset-closed, binary complete, and non empty.

A coherent space is a subset-closed binary complete non empty set.

In the sequel C, D denote coherent spaces.

One can prove the following propositions:

- (1) $\emptyset \in C$.
- (2) 2^X is a coherent space.
- (3) $\{\emptyset\}$ is a coherent space.
- (4) If $x \in \bigcup C$, then $\{x\} \in C$.

Let C be a coherent space. The functor $\text{Web}(C)$ yields a tolerance of $\bigcup C$ and is defined by:

(Def. 3) For all x, y holds $\langle x, y \rangle \in \text{Web}(C)$ iff there exists X such that $X \in C$ and $x \in X$ and $y \in X$.

In the sequel T denotes a tolerance of $\bigcup C$.

The following propositions are true:

- (5) $T = \text{Web}(C)$ iff for all x, y holds $\langle x, y \rangle \in T$ iff $\{x, y\} \in C$.
- (6) $a \in C$ iff for all x, y such that $x \in a$ and $y \in a$ holds $\{x, y\} \in C$.

¹ The definition (Def. 1) has been removed.

- (7) $a \in C$ iff for all x, y such that $x \in a$ and $y \in a$ holds $\langle x, y \rangle \in \text{Web}(C)$.
- (8) If for all x, y such that $x \in a$ and $y \in a$ holds $\{x, y\} \in C$, then $a \subseteq \bigcup C$.
- (9) If $\text{Web}(C) = \text{Web}(D)$, then $C = D$.
- (10) If $\bigcup C \in C$, then $C = 2^{\bigcup C}$.
- (11) If $C = 2^{\bigcup C}$, then $\text{Web}(C) = \bigvee_{\bigcup C}$.

Let X be a set and let E be a tolerance of X . The functor $\text{CohSp}(E)$ yields a coherent space and is defined by:

(Def. 4) For every a holds $a \in \text{CohSp}(E)$ iff for all x, y such that $x \in a$ and $y \in a$ holds $\langle x, y \rangle \in E$.

In the sequel E is a tolerance of X .

We now state four propositions:

- (12) $\text{Web}(\text{CohSp}(E)) = E$.
- (13) $\text{CohSp}(\text{Web}(C)) = C$.
- (14) $a \in \text{CohSp}(E)$ iff a is a set of mutually elements w.r.t. E .
- (15) $\text{CohSp}(E) = \text{TolSets}E$.

2. CATEGORY OF COHERENT SPACES

Let us consider X . The functor $\text{CSp}(X)$ yields a set and is defined by:

(Def. 5) $\text{CSp}(X) = \{x; x \text{ ranges over subsets of } 2^X : x \text{ is a coherent space}\}$.

Let us consider X . One can verify that $\text{CSp}(X)$ is non empty.

Let X be a set. Note that every element of $\text{CSp}(X)$ is subset-closed, binary complete, and non empty.

In the sequel C, C_1, C_2 are elements of $\text{CSp}(X)$.

One can prove the following proposition

- (16) If $\{x, y\} \in C$, then $x \in \bigcup C$ and $y \in \bigcup C$.

Let us consider X . The functor $\text{Funcs}_C X$ yields a set and is defined by:

(Def. 7)² $\text{Funcs}_C X = \bigcup \{(\bigcup y)^{\bigcup x} : x \text{ ranges over elements of } \text{CSp}(X), y \text{ ranges over elements of } \text{CSp}(X)\}$.

Let us consider X . One can verify that $\text{Funcs}_C X$ is non empty and functional.

In the sequel g denotes an element of $\text{Funcs}_C X$.

One can prove the following proposition

- (17) $x \in \text{Funcs}_C X$ iff there exist C_1, C_2 such that if $\bigcup C_2 = \emptyset$, then $\bigcup C_1 = \emptyset$ and x is a function from $\bigcup C_1$ into $\bigcup C_2$.

Let us consider X . The functor $\text{Maps}_C X$ yields a set and is defined by the condition (Def. 8).

(Def. 8) $\text{Maps}_C X = \{\langle \langle C, C_3 \rangle, f \rangle; C \text{ ranges over elements of } \text{CSp}(X), C_3 \text{ ranges over elements of } \text{CSp}(X), f \text{ ranges over elements of } \text{Funcs}_C X : (\bigcup C_3 = \emptyset \Rightarrow \bigcup C = \emptyset) \wedge f \text{ is a function from } \bigcup C \text{ into } \bigcup C_3 \wedge \bigwedge_{x,y} (\{x, y\} \in C \Rightarrow \{f(x), f(y)\} \in C_3)\}$.

Let us consider X . One can check that $\text{Maps}_C X$ is non empty.

In the sequel l, l_1, l_2, l_3 denote elements of $\text{Maps}_C X$.

One can prove the following two propositions:

² The definition (Def. 6) has been removed.

- (18) There exist g, C_1, C_2 such that
- (i) $l = \langle \langle C_1, C_2 \rangle, g \rangle$,
 - (ii) if $\bigcup C_2 = \emptyset$, then $\bigcup C_1 = \emptyset$,
 - (iii) g is a function from $\bigcup C_1$ into $\bigcup C_2$, and
 - (iv) for all x, y such that $\{x, y\} \in C_1$ holds $\{g(x), g(y)\} \in C_2$.
- (19) Let f be a function from $\bigcup C_1$ into $\bigcup C_2$. Suppose if $\bigcup C_2 = \emptyset$, then $\bigcup C_1 = \emptyset$ and for all x, y such that $\{x, y\} \in C_1$ holds $\{f(x), f(y)\} \in C_2$. Then $\langle \langle C_1, C_2 \rangle, f \rangle \in \text{Maps}_{\mathcal{C}}X$.

Let X be a set and let l be an element of $\text{Maps}_{\mathcal{C}}X$. Note that l_2 is function-like and relation-like. Let us consider X, l . The functor $\text{dom } l$ yields an element of $\text{CSp}(X)$ and is defined as follows:

(Def. 10)³ $\text{dom } l = (l_1)_1$.

The functor $\text{cod } l$ yielding an element of $\text{CSp}(X)$ is defined by:

(Def. 11) $\text{cod } l = (l_1)_2$.

One can prove the following proposition

(20) $l = \langle \langle \text{dom } l, \text{cod } l \rangle, l_2 \rangle$.

Let us consider X, C . The functor $\text{id}(C)$ yielding an element of $\text{Maps}_{\mathcal{C}}X$ is defined as follows:

(Def. 12) $\text{id}(C) = \langle \langle C, C \rangle, \text{id}_{\bigcup C} \rangle$.

The following proposition is true

(21) $\bigcup \text{cod } l \neq \emptyset$ or $\bigcup \text{dom } l = \emptyset$ but l_2 is a function from $\bigcup \text{dom } l$ into $\bigcup \text{cod } l$ but for all x, y such that $\{x, y\} \in \text{dom } l$ holds $\{l_2(x), l_2(y)\} \in \text{cod } l$.

Let us consider X, l_1, l_2 . Let us assume that $\text{cod } l_1 = \text{dom } l_2$. The functor $l_2 \cdot l_1$ yields an element of $\text{Maps}_{\mathcal{C}}X$ and is defined by:

(Def. 13) $l_2 \cdot l_1 = \langle \langle \text{dom } l_1, \text{cod } l_2 \rangle, (l_2)_2 \cdot (l_1)_2 \rangle$.

One can prove the following propositions:

(22) If $\text{dom } l_2 = \text{cod } l_1$, then $(l_2 \cdot l_1)_2 = (l_2)_2 \cdot (l_1)_2$ and $\text{dom}(l_2 \cdot l_1) = \text{dom } l_1$ and $\text{cod}(l_2 \cdot l_1) = \text{cod } l_2$.

(23) If $\text{dom } l_2 = \text{cod } l_1$ and $\text{dom } l_3 = \text{cod } l_2$, then $l_3 \cdot (l_2 \cdot l_1) = (l_3 \cdot l_2) \cdot l_1$.

(24) $(\text{id}(C))_2 = \text{id}_{\bigcup C}$ and $\text{dom } \text{id}(C) = C$ and $\text{cod } \text{id}(C) = C$.

(25) $l \cdot \text{id}(\text{dom } l) = l$ and $\text{id}(\text{cod } l) \cdot l = l$.

Let us consider X . The functor $\text{Dom}_{\text{CSp}}X$ yields a function from $\text{Maps}_{\mathcal{C}}X$ into $\text{CSp}(X)$ and is defined by:

(Def. 14) For every l holds $(\text{Dom}_{\text{CSp}}X)(l) = \text{dom } l$.

The functor $\text{Cod}_{\text{CSp}}X$ yields a function from $\text{Maps}_{\mathcal{C}}X$ into $\text{CSp}(X)$ and is defined by:

(Def. 15) For every l holds $(\text{Cod}_{\text{CSp}}X)(l) = \text{cod } l$.

The functor $\cdot_{\text{CSp}}X$ yields a partial function from $[\text{Maps}_{\mathcal{C}}X, \text{Maps}_{\mathcal{C}}X]$ to $\text{Maps}_{\mathcal{C}}X$ and is defined as follows:

(Def. 16) For all l_2, l_1 holds $\langle l_2, l_1 \rangle \in \text{dom } \cdot_{\text{CSp}}X$ iff $\text{dom } l_2 = \text{cod } l_1$ and for all l_2, l_1 such that $\text{dom } l_2 = \text{cod } l_1$ holds $(\cdot_{\text{CSp}}X)(\langle l_2, l_1 \rangle) = l_2 \cdot l_1$.

³ The definition (Def. 9) has been removed.

The functor $\text{Id}_{\text{CSp}} X$ yields a function from $\text{CSp}(X)$ into $\text{Maps}_{\text{C}} X$ and is defined by:

(Def. 17) For every C holds $(\text{Id}_{\text{CSp}} X)(C) = \text{id}(C)$.

We now state the proposition

(26) $\langle \text{CSp}(X), \text{Maps}_{\text{C}} X, \text{Dom}_{\text{CSp}} X, \text{Cod}_{\text{CSp}} X, \cdot_{\text{CSp}} X, \text{Id}_{\text{CSp}} X \rangle$ is a category.

Let us consider X . The X -coherent space category yielding a category is defined by:

(Def. 18) The X -coherent space category = $\langle \text{CSp}(X), \text{Maps}_{\text{C}} X, \text{Dom}_{\text{CSp}} X, \text{Cod}_{\text{CSp}} X, \cdot_{\text{CSp}} X, \text{Id}_{\text{CSp}} X \rangle$.

3. CATEGORY OF TOLERANCES

Let X be a set. The tolerances on X constitute a set defined as follows:

(Def. 19) $x \in$ the tolerances on X iff x is a tolerance of X .

Let X be a set. Note that the tolerances on X is non empty.

Let X be a set. The tolerances on subsets of X constitute a set defined as follows:

(Def. 20) The tolerances on subsets of $X = \bigcup \{\text{the tolerances on } Y : Y \text{ ranges over subsets of } X\}$.

Let X be a set. One can check that the tolerances on subsets of X is non empty.

One can prove the following propositions:

(27) $x \in$ the tolerances on subsets of X iff there exists A such that $A \subseteq X$ and x is a tolerance of A .

(28) $\nabla_a \in$ the tolerances on a .

(30)⁴ $\emptyset \in$ the tolerances on subsets of X .

(31) If $a \subseteq X$, then $\nabla_a \in$ the tolerances on subsets of X .

(32) If $a \subseteq X$, then $\text{id}_a \in$ the tolerances on subsets of X .

(33) $\nabla_X \in$ the tolerances on subsets of X .

(34) $\text{id}_X \in$ the tolerances on subsets of X .

Let us consider X . The functor $\text{TOL}(X)$ yields a set and is defined by:

(Def. 21) $\text{TOL}(X) = \{\langle t, Y \rangle; t \text{ ranges over elements of the tolerances on subsets of } X, Y \text{ ranges over elements of } 2^X; t \text{ is a tolerance of } Y\}$.

Let us consider X . One can verify that $\text{TOL}(X)$ is non empty.

In the sequel T, T_1, T_2 denote elements of $\text{TOL}(X)$.

One can prove the following propositions:

(35) $\langle \emptyset, \emptyset \rangle \in \text{TOL}(X)$.

(36) If $a \subseteq X$, then $\langle \text{id}_a, a \rangle \in \text{TOL}(X)$.

(37) If $a \subseteq X$, then $\langle \nabla_a, a \rangle \in \text{TOL}(X)$.

(38) $\langle \text{id}_X, X \rangle \in \text{TOL}(X)$.

(39) $\langle \nabla_X, X \rangle \in \text{TOL}(X)$.

Let us consider X, T . Then T_2 is an element of 2^X . Then T_1 is a tolerance of T_2 .

Let us consider X . The functor $\text{Funcs}_{\text{T}} X$ yields a set and is defined by:

⁴ The proposition (29) has been removed.

(Def. 22) $\text{Funcs}_T X = \bigcup \{ \langle (T_3)_2 \rangle^{T_2} : T \text{ ranges over elements of } \text{TOL}(X), T_3 \text{ ranges over elements of } \text{TOL}(X) \}$.

Let us consider X . Observe that $\text{Funcs}_T X$ is non empty and functional.

In the sequel f is an element of $\text{Funcs}_T X$.

One can prove the following proposition

(40) $x \in \text{Funcs}_T X$ iff there exist T_1, T_2 such that if $(T_2)_2 = \emptyset$, then $(T_1)_2 = \emptyset$ and x is a function from $(T_1)_2$ into $(T_2)_2$.

Let us consider X . The functor $\text{Maps}_T X$ yielding a set is defined by the condition (Def. 23).

(Def. 23) $\text{Maps}_T X = \{ \langle \langle T, T_3 \rangle, f \rangle ; T \text{ ranges over elements of } \text{TOL}(X), T_3 \text{ ranges over elements of } \text{TOL}(X), f \text{ ranges over elements of } \text{Funcs}_T X : ((T_3)_2 = \emptyset \Rightarrow T_2 = \emptyset) \wedge f \text{ is a function from } T_2 \text{ into } (T_3)_2 \wedge \bigwedge_{x,y} (\langle x, y \rangle \in T_1 \Rightarrow \langle f(x), f(y) \rangle \in (T_3)_1) \}$.

Let us consider X . Note that $\text{Maps}_T X$ is non empty.

In the sequel m, m_1, m_2, m_3 are elements of $\text{Maps}_T X$.

The following propositions are true:

(41) There exist f, T_1, T_2 such that

(i) $m = \langle \langle T_1, T_2 \rangle, f \rangle$,

(ii) if $(T_2)_2 = \emptyset$, then $(T_1)_2 = \emptyset$,

(iii) f is a function from $(T_1)_2$ into $(T_2)_2$, and

(iv) for all x, y such that $\langle x, y \rangle \in (T_1)_1$ holds $\langle f(x), f(y) \rangle \in (T_2)_1$.

(42) Let f be a function from $(T_1)_2$ into $(T_2)_2$. Suppose if $(T_2)_2 = \emptyset$, then $(T_1)_2 = \emptyset$ and for all x, y such that $\langle x, y \rangle \in (T_1)_1$ holds $\langle f(x), f(y) \rangle \in (T_2)_1$. Then $\langle \langle T_1, T_2 \rangle, f \rangle \in \text{Maps}_T X$.

Let X be a set and let m be an element of $\text{Maps}_T X$. One can check that m_2 is function-like and relation-like.

Let us consider X, m . The functor $\text{dom } m$ yields an element of $\text{TOL}(X)$ and is defined by:

(Def. 25)⁵ $\text{dom } m = (m_1)_1$.

The functor $\text{cod } m$ yielding an element of $\text{TOL}(X)$ is defined as follows:

(Def. 26) $\text{cod } m = (m_1)_2$.

Next we state the proposition

(43) $m = \langle \langle \text{dom } m, \text{cod } m \rangle, m_2 \rangle$.

Let us consider X, T . The functor $\text{id}(T)$ yields an element of $\text{Maps}_T X$ and is defined as follows:

(Def. 27) $\text{id}(T) = \langle \langle T, T \rangle, \text{id}_{T_2} \rangle$.

Next we state the proposition

(44) $(\text{cod } m)_2 \neq \emptyset$ or $(\text{dom } m)_2 = \emptyset$ but m_2 is a function from $(\text{dom } m)_2$ into $(\text{cod } m)_2$ but for all x, y such that $\langle x, y \rangle \in (\text{dom } m)_1$ holds $\langle m_2(x), m_2(y) \rangle \in (\text{cod } m)_1$.

Let us consider X, m_1, m_2 . Let us assume that $\text{cod } m_1 = \text{dom } m_2$. The functor $m_2 \cdot m_1$ yielding an element of $\text{Maps}_T X$ is defined by:

(Def. 28) $m_2 \cdot m_1 = \langle \langle \text{dom } m_1, \text{cod } m_2 \rangle, (m_2)_2 \cdot (m_1)_2 \rangle$.

The following four propositions are true:

⁵ The definition (Def. 24) has been removed.

(45) If $\text{dom } m_2 = \text{cod } m_1$, then $(m_2 \cdot m_1)_2 = (m_2)_2 \cdot (m_1)_2$ and $\text{dom}(m_2 \cdot m_1) = \text{dom } m_1$ and $\text{cod}(m_2 \cdot m_1) = \text{cod } m_2$.

(46) If $\text{dom } m_2 = \text{cod } m_1$ and $\text{dom } m_3 = \text{cod } m_2$, then $m_3 \cdot (m_2 \cdot m_1) = (m_3 \cdot m_2) \cdot m_1$.

(47) $(\text{id}(T))_2 = \text{id}_{T_2}$ and $\text{dom id}(T) = T$ and $\text{cod id}(T) = T$.

(48) $m \cdot \text{id}(\text{dom } m) = m$ and $\text{id}(\text{cod } m) \cdot m = m$.

Let us consider X . The functor Dom_X yields a function from $\text{Maps}_T X$ into $\text{TOL}(X)$ and is defined as follows:

(Def. 29) For every m holds $\text{Dom}_X(m) = \text{dom } m$.

The functor Cod_X yields a function from $\text{Maps}_T X$ into $\text{TOL}(X)$ and is defined as follows:

(Def. 30) For every m holds $\text{Cod}_X(m) = \text{cod } m$.

The functor \cdot_X yields a partial function from $[\text{Maps}_T X, \text{Maps}_T X]$ to $\text{Maps}_T X$ and is defined by:

(Def. 31) For all m_2, m_1 holds $\langle m_2, m_1 \rangle \in \text{dom}(\cdot_X)$ iff $\text{dom } m_2 = \text{cod } m_1$ and for all m_2, m_1 such that $\text{dom } m_2 = \text{cod } m_1$ holds $\cdot_X(\langle m_2, m_1 \rangle) = m_2 \cdot m_1$.

The functor Id_X yielding a function from $\text{TOL}(X)$ into $\text{Maps}_T X$ is defined by:

(Def. 32) For every T holds $\text{Id}_X(T) = \text{id}(T)$.

Next we state the proposition

(49) $\langle \text{TOL}(X), \text{Maps}_T X, \text{Dom}_X, \text{Cod}_X, \cdot_X, \text{Id}_X \rangle$ is a category.

Let us consider X . The X -tolerance category is a category and is defined by:

(Def. 33) The X -tolerance category $= \langle \text{TOL}(X), \text{Maps}_T X, \text{Dom}_X, \text{Cod}_X, \cdot_X, \text{Id}_X \rangle$.

REFERENCES

- [1] Grzegorz Bancerek. Tarski's classes and ranks. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol12/classes1.html>.
- [2] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol11/funct_1.html.
- [3] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol11/funct_2.html.
- [4] Czesław Byliński. Introduction to categories and functors. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol11/cat_1.html.
- [5] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol11/partfun1.html>.
- [6] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol11/zfmisc_1.html.
- [7] Krzysztof Hryniewiecki. Relations of tolerance. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol12/toler_1.html.
- [8] Konrad Raczkowski and Paweł Sadowski. Equivalence relations and classes of abstraction. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol11/eqrel_1.html.
- [9] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [10] Andrzej Trybulec. Tuples, projections and Cartesian products. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol11/mcart_1.html.
- [11] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol11/subset_1.html.
- [12] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol11/relat_1.html.

- [13] Edmund Woronowicz. Relations defined on sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/relset_1.html.

Received December 29, 1992

Published January 2, 2004
