

# Enumerated Sets

Andrzej Trybulec  
Warsaw University  
Białystok

**Summary.** We prove basic facts about enumerated sets: definitional theorems and their immediate consequences, some theorems related to the decomposition of an enumerated set into union of two sets, facts about removing elements that occur more than once, and facts about permutations of enumerated sets (with the length  $\leq 4$ ). The article includes also schemes enabling instantiation of up to nine universal quantifiers.

MML Identifier: ENUMSET1.

WWW: <http://mizar.org/JFM/Vol1/enumset1.html>

The article [1] provides the notation and terminology for this paper.

In this paper  $x, x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, X$  are sets.

In this article we present several logical schemes. The scheme *UI1* deals with a set  $\mathcal{A}$  and a unary predicate  $\mathcal{P}$ , and states that:

$\mathcal{P}[\mathcal{A}]$

provided the parameters have the following property:

- For every  $x_1$  holds  $\mathcal{P}[x_1]$ .

The scheme *UI2* deals with sets  $\mathcal{A}, \mathcal{B}$  and a binary predicate  $\mathcal{P}$ , and states that:

$\mathcal{P}[\mathcal{A}, \mathcal{B}]$

provided the following condition is satisfied:

- For all  $x_1, x_2$  holds  $\mathcal{P}[x_1, x_2]$ .

The scheme *UI3* deals with sets  $\mathcal{A}, \mathcal{B}, \mathcal{C}$  and a ternary predicate  $\mathcal{P}$ , and states that:

$\mathcal{P}[\mathcal{A}, \mathcal{B}, \mathcal{C}]$

provided the parameters meet the following condition:

- For all  $x_1, x_2, x_3$  holds  $\mathcal{P}[x_1, x_2, x_3]$ .

The scheme *UI4* deals with sets  $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}$  and a 4-ary predicate  $\mathcal{P}$ , and states that:

$\mathcal{P}[\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}]$

provided the following requirement is met:

- For all  $x_1, x_2, x_3, x_4$  holds  $\mathcal{P}[x_1, x_2, x_3, x_4]$ .

The scheme *UI5* deals with sets  $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}, \mathcal{E}$  and a 5-ary predicate  $\mathcal{P}$ , and states that:

$\mathcal{P}[\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}, \mathcal{E}]$

provided the following requirement is met:

- For all  $x_1, x_2, x_3, x_4, x_5$  holds  $\mathcal{P}[x_1, x_2, x_3, x_4, x_5]$ .

The scheme *UI6* deals with sets  $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}, \mathcal{E}, \mathcal{F}$  and a 6-ary predicate  $\mathcal{P}$ , and states that:

$\mathcal{P}[\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}, \mathcal{E}, \mathcal{F}]$

provided the parameters meet the following condition:

- For all  $x_1, x_2, x_3, x_4, x_5, x_6$  holds  $\mathcal{P}[x_1, x_2, x_3, x_4, x_5, x_6]$ .

The scheme *UI7* deals with sets  $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}, \mathcal{E}, \mathcal{F}, \mathcal{G}$  and a 7-ary predicate  $\mathcal{P}$ , and states that:

$\mathcal{P}[\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}, \mathcal{E}, \mathcal{F}, \mathcal{G}]$

provided the parameters satisfy the following condition:

- For all  $x_1, x_2, x_3, x_4, x_5, x_6, x_7$  holds  $\mathcal{P}[x_1, x_2, x_3, x_4, x_5, x_6, x_7]$ .

The scheme *UI8* deals with sets  $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}, \mathcal{E}, \mathcal{F}, \mathcal{G}, \mathcal{H}$  and a 8-ary predicate  $\mathcal{P}$ , and states that:

$$\mathcal{P}[\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}, \mathcal{E}, \mathcal{F}, \mathcal{G}, \mathcal{H}]$$

provided the parameters satisfy the following condition:

- For all  $x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8$  holds  $\mathcal{P}[x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8]$ .

The scheme *UI9* deals with sets  $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}, \mathcal{E}, \mathcal{F}, \mathcal{G}, \mathcal{H}, \mathcal{I}$  and a 9-ary predicate  $\mathcal{P}$ , and states that:

$$\mathcal{P}[\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D}, \mathcal{E}, \mathcal{F}, \mathcal{G}, \mathcal{H}, \mathcal{I}]$$

provided the following condition is satisfied:

- For all sets  $x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9$  holds  $\mathcal{P}[x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8, x_9]$ .

Let us consider  $x_1, x_2, x_3$ . The functor  $\{x_1, x_2, x_3\}$  yields a set and is defined as follows:

(Def. 1)  $x \in \{x_1, x_2, x_3\}$  iff  $x = x_1$  or  $x = x_2$  or  $x = x_3$ .

Next we state three propositions:

(13)<sup>1</sup> If  $x \in \{x_1, x_2, x_3\}$ , then  $x = x_1$  or  $x = x_2$  or  $x = x_3$ .

(14) If  $x = x_1$  or  $x = x_2$  or  $x = x_3$ , then  $x \in \{x_1, x_2, x_3\}$ .

(15) For all  $x_1, x_2, x_3, X$  such that for every  $x$  holds  $x \in X$  iff  $x = x_1$  or  $x = x_2$  or  $x = x_3$  holds  $X = \{x_1, x_2, x_3\}$ .

Let us consider  $x_1, x_2, x_3, x_4$ . The functor  $\{x_1, x_2, x_3, x_4\}$  yielding a set is defined by:

(Def. 2)  $x \in \{x_1, x_2, x_3, x_4\}$  iff  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$ .

We now state three propositions:

(18)<sup>2</sup> If  $x \in \{x_1, x_2, x_3, x_4\}$ , then  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$ .

(19) If  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$ , then  $x \in \{x_1, x_2, x_3, x_4\}$ .

(20) For all  $x_1, x_2, x_3, x_4, X$  such that for every  $x$  holds  $x \in X$  iff  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  holds  $X = \{x_1, x_2, x_3, x_4\}$ .

Let us consider  $x_1, x_2, x_3, x_4, x_5$ . The functor  $\{x_1, x_2, x_3, x_4, x_5\}$  yields a set and is defined by:

(Def. 3)  $x \in \{x_1, x_2, x_3, x_4, x_5\}$  iff  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  or  $x = x_5$ .

One can prove the following three propositions:

(23)<sup>3</sup> If  $x \in \{x_1, x_2, x_3, x_4, x_5\}$ , then  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  or  $x = x_5$ .

(24) If  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  or  $x = x_5$ , then  $x \in \{x_1, x_2, x_3, x_4, x_5\}$ .

(25) For every set  $X$  such that for every  $x$  holds  $x \in X$  iff  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  or  $x = x_5$  holds  $X = \{x_1, x_2, x_3, x_4, x_5\}$ .

Let us consider  $x_1, x_2, x_3, x_4, x_5, x_6$ . The functor  $\{x_1, x_2, x_3, x_4, x_5, x_6\}$  yields a set and is defined as follows:

(Def. 4)  $x \in \{x_1, x_2, x_3, x_4, x_5, x_6\}$  iff  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  or  $x = x_5$  or  $x = x_6$ .

One can prove the following three propositions:

(28)<sup>4</sup> If  $x \in \{x_1, x_2, x_3, x_4, x_5, x_6\}$ , then  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  or  $x = x_5$  or  $x = x_6$ .

<sup>1</sup> The propositions (1)–(12) have been removed.

<sup>2</sup> The propositions (16) and (17) have been removed.

<sup>3</sup> The propositions (21) and (22) have been removed.

<sup>4</sup> The propositions (26) and (27) have been removed.

(29) If  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  or  $x = x_5$  or  $x = x_6$ , then  $x \in \{x_1, x_2, x_3, x_4, x_5, x_6\}$ .

(30) Let  $X$  be a set. Suppose that for every  $x$  holds  $x \in X$  iff  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  or  $x = x_5$  or  $x = x_6$ . Then  $X = \{x_1, x_2, x_3, x_4, x_5, x_6\}$ .

Let us consider  $x_1, x_2, x_3, x_4, x_5, x_6, x_7$ . The functor  $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7\}$  yields a set and is defined as follows:

(Def. 5)  $x \in \{x_1, x_2, x_3, x_4, x_5, x_6, x_7\}$  iff  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  or  $x = x_5$  or  $x = x_6$  or  $x = x_7$ .

The following propositions are true:

(33)<sup>5</sup> If  $x \in \{x_1, x_2, x_3, x_4, x_5, x_6, x_7\}$ , then  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  or  $x = x_5$  or  $x = x_6$  or  $x = x_7$ .

(34) If  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  or  $x = x_5$  or  $x = x_6$  or  $x = x_7$ , then  $x \in \{x_1, x_2, x_3, x_4, x_5, x_6, x_7\}$ .

(35) Let  $X$  be a set. Suppose that for every  $x$  holds  $x \in X$  iff  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  or  $x = x_5$  or  $x = x_6$  or  $x = x_7$ . Then  $X = \{x_1, x_2, x_3, x_4, x_5, x_6, x_7\}$ .

Let us consider  $x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8$ . The functor  $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\}$  yields a set and is defined as follows:

(Def. 6)  $x \in \{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\}$  iff  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  or  $x = x_5$  or  $x = x_6$  or  $x = x_7$  or  $x = x_8$ .

The following propositions are true:

(38)<sup>6</sup> If  $x \in \{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\}$ , then  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  or  $x = x_5$  or  $x = x_6$  or  $x = x_7$  or  $x = x_8$ .

(39) If  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  or  $x = x_5$  or  $x = x_6$  or  $x = x_7$  or  $x = x_8$ , then  $x \in \{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\}$ .

(40) Let  $X$  be a set. Suppose that for every  $x$  holds  $x \in X$  iff  $x = x_1$  or  $x = x_2$  or  $x = x_3$  or  $x = x_4$  or  $x = x_5$  or  $x = x_6$  or  $x = x_7$  or  $x = x_8$ . Then  $X = \{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\}$ .

$$(41) \quad \{x_1, x_2\} = \{x_1\} \cup \{x_2\}.$$

$$(42) \quad \{x_1, x_2, x_3\} = \{x_1\} \cup \{x_2, x_3\}.$$

$$(43) \quad \{x_1, x_2, x_3\} = \{x_1, x_2\} \cup \{x_3\}.$$

$$(44) \quad \{x_1, x_2, x_3, x_4\} = \{x_1\} \cup \{x_2, x_3, x_4\}.$$

$$(45) \quad \{x_1, x_2, x_3, x_4\} = \{x_1, x_2\} \cup \{x_3, x_4\}.$$

$$(46) \quad \{x_1, x_2, x_3, x_4\} = \{x_1, x_2, x_3\} \cup \{x_4\}.$$

$$(47) \quad \{x_1, x_2, x_3, x_4, x_5\} = \{x_1\} \cup \{x_2, x_3, x_4, x_5\}.$$

$$(48) \quad \{x_1, x_2, x_3, x_4, x_5\} = \{x_1, x_2\} \cup \{x_3, x_4, x_5\}.$$

$$(49) \quad \{x_1, x_2, x_3, x_4, x_5\} = \{x_1, x_2, x_3\} \cup \{x_4, x_5\}.$$

$$(50) \quad \{x_1, x_2, x_3, x_4, x_5\} = \{x_1, x_2, x_3, x_4\} \cup \{x_5\}.$$

$$(51) \quad \{x_1, x_2, x_3, x_4, x_5, x_6\} = \{x_1\} \cup \{x_2, x_3, x_4, x_5, x_6\}.$$

$$(52) \quad \{x_1, x_2, x_3, x_4, x_5, x_6\} = \{x_1, x_2\} \cup \{x_3, x_4, x_5, x_6\}.$$

<sup>5</sup> The propositions (31) and (32) have been removed.

<sup>6</sup> The propositions (36) and (37) have been removed.

- (53)  $\{x_1, x_2, x_3, x_4, x_5, x_6\} = \{x_1, x_2, x_3\} \cup \{x_4, x_5, x_6\}$ .
- (54)  $\{x_1, x_2, x_3, x_4, x_5, x_6\} = \{x_1, x_2, x_3, x_4\} \cup \{x_5, x_6\}$ .
- (55)  $\{x_1, x_2, x_3, x_4, x_5, x_6\} = \{x_1, x_2, x_3, x_4, x_5\} \cup \{x_6\}$ .
- (56)  $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7\} = \{x_1\} \cup \{x_2, x_3, x_4, x_5, x_6, x_7\}$ .
- (57)  $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7\} = \{x_1, x_2\} \cup \{x_3, x_4, x_5, x_6, x_7\}$ .
- (58)  $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7\} = \{x_1, x_2, x_3\} \cup \{x_4, x_5, x_6, x_7\}$ .
- (59)  $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7\} = \{x_1, x_2, x_3, x_4\} \cup \{x_5, x_6, x_7\}$ .
- (60)  $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7\} = \{x_1, x_2, x_3, x_4, x_5\} \cup \{x_6, x_7\}$ .
- (61)  $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7\} = \{x_1, x_2, x_3, x_4, x_5, x_6\} \cup \{x_7\}$ .
- (62)  $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\} = \{x_1\} \cup \{x_2, x_3, x_4, x_5, x_6, x_7, x_8\}$ .
- (63)  $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\} = \{x_1, x_2\} \cup \{x_3, x_4, x_5, x_6, x_7, x_8\}$ .
- (64)  $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\} = \{x_1, x_2, x_3\} \cup \{x_4, x_5, x_6, x_7, x_8\}$ .
- (65)  $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\} = \{x_1, x_2, x_3, x_4\} \cup \{x_5, x_6, x_7, x_8\}$ .
- (66)  $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\} = \{x_1, x_2, x_3, x_4, x_5\} \cup \{x_6, x_7, x_8\}$ .
- (67)  $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\} = \{x_1, x_2, x_3, x_4, x_5, x_6\} \cup \{x_7, x_8\}$ .
- (68)  $\{x_1, x_2, x_3, x_4, x_5, x_6, x_7, x_8\} = \{x_1, x_2, x_3, x_4, x_5, x_6, x_7\} \cup \{x_8\}$ .
- (69)  $\{x_1, x_1\} = \{x_1\}$ .
- (70)  $\{x_1, x_1, x_2\} = \{x_1, x_2\}$ .
- (71)  $\{x_1, x_1, x_2, x_3\} = \{x_1, x_2, x_3\}$ .
- (72)  $\{x_1, x_1, x_2, x_3, x_4\} = \{x_1, x_2, x_3, x_4\}$ .
- (73)  $\{x_1, x_1, x_2, x_3, x_4, x_5\} = \{x_1, x_2, x_3, x_4, x_5\}$ .
- (74)  $\{x_1, x_1, x_2, x_3, x_4, x_5, x_6\} = \{x_1, x_2, x_3, x_4, x_5, x_6\}$ .
- (75)  $\{x_1, x_1, x_2, x_3, x_4, x_5, x_6, x_7\} = \{x_1, x_2, x_3, x_4, x_5, x_6, x_7\}$ .
- (76)  $\{x_1, x_1, x_1\} = \{x_1\}$ .
- (77)  $\{x_1, x_1, x_1, x_2\} = \{x_1, x_2\}$ .
- (78)  $\{x_1, x_1, x_1, x_2, x_3\} = \{x_1, x_2, x_3\}$ .
- (79)  $\{x_1, x_1, x_1, x_2, x_3, x_4\} = \{x_1, x_2, x_3, x_4\}$ .
- (80)  $\{x_1, x_1, x_1, x_2, x_3, x_4, x_5\} = \{x_1, x_2, x_3, x_4, x_5\}$ .
- (81)  $\{x_1, x_1, x_1, x_2, x_3, x_4, x_5, x_6\} = \{x_1, x_2, x_3, x_4, x_5, x_6\}$ .
- (82)  $\{x_1, x_1, x_1, x_1\} = \{x_1\}$ .
- (83)  $\{x_1, x_1, x_1, x_1, x_2\} = \{x_1, x_2\}$ .
- (84)  $\{x_1, x_1, x_1, x_1, x_2, x_3\} = \{x_1, x_2, x_3\}$ .
- (85)  $\{x_1, x_1, x_1, x_1, x_2, x_3, x_4\} = \{x_1, x_2, x_3, x_4\}$ .
- (86)  $\{x_1, x_1, x_1, x_1, x_2, x_3, x_4, x_5\} = \{x_1, x_2, x_3, x_4, x_5\}$ .

- (87)  $\{x_1, x_1, x_1, x_1, x_1\} = \{x_1\}$ .  
 (88)  $\{x_1, x_1, x_1, x_1, x_1, x_2\} = \{x_1, x_2\}$ .  
 (89)  $\{x_1, x_1, x_1, x_1, x_1, x_2, x_3\} = \{x_1, x_2, x_3\}$ .  
 (90)  $\{x_1, x_1, x_1, x_1, x_1, x_2, x_3, x_4\} = \{x_1, x_2, x_3, x_4\}$ .  
 (91)  $\{x_1, x_1, x_1, x_1, x_1, x_1\} = \{x_1\}$ .  
 (92)  $\{x_1, x_1, x_1, x_1, x_1, x_1, x_2\} = \{x_1, x_2\}$ .  
 (93)  $\{x_1, x_1, x_1, x_1, x_1, x_1, x_2, x_3\} = \{x_1, x_2, x_3\}$ .  
 (94)  $\{x_1, x_1, x_1, x_1, x_1, x_1, x_1\} = \{x_1\}$ .  
 (95)  $\{x_1, x_1, x_1, x_1, x_1, x_1, x_1, x_2\} = \{x_1, x_2\}$ .  
 (96)  $\{x_1, x_1, x_1, x_1, x_1, x_1, x_1, x_1\} = \{x_1\}$ .  
 (98)<sup>7</sup>  $\{x_1, x_2, x_3\} = \{x_1, x_3, x_2\}$ .  
 (99)  $\{x_1, x_2, x_3\} = \{x_2, x_1, x_3\}$ .  
 (100)  $\{x_1, x_2, x_3\} = \{x_2, x_3, x_1\}$ .  
 (102)<sup>8</sup>  $\{x_1, x_2, x_3\} = \{x_3, x_2, x_1\}$ .  
 (103)  $\{x_1, x_2, x_3, x_4\} = \{x_1, x_2, x_4, x_3\}$ .  
 (104)  $\{x_1, x_2, x_3, x_4\} = \{x_1, x_3, x_2, x_4\}$ .  
 (105)  $\{x_1, x_2, x_3, x_4\} = \{x_1, x_3, x_4, x_2\}$ .  
 (107)<sup>9</sup>  $\{x_1, x_2, x_3, x_4\} = \{x_1, x_4, x_3, x_2\}$ .  
 (108)  $\{x_1, x_2, x_3, x_4\} = \{x_2, x_1, x_3, x_4\}$ .  
 (109)  $\{x_1, x_2, x_3, x_4\} = \{x_2, x_1, x_4, x_3\}$ .  
 (110)  $\{x_1, x_2, x_3, x_4\} = \{x_2, x_3, x_1, x_4\}$ .  
 (111)  $\{x_1, x_2, x_3, x_4\} = \{x_2, x_3, x_4, x_1\}$ .  
 (112)  $\{x_1, x_2, x_3, x_4\} = \{x_2, x_4, x_1, x_3\}$ .  
 (113)  $\{x_1, x_2, x_3, x_4\} = \{x_2, x_4, x_3, x_1\}$ .  
 (116)<sup>10</sup>  $\{x_1, x_2, x_3, x_4\} = \{x_3, x_2, x_1, x_4\}$ .  
 (117)  $\{x_1, x_2, x_3, x_4\} = \{x_3, x_2, x_4, x_1\}$ .  
 (118)  $\{x_1, x_2, x_3, x_4\} = \{x_3, x_4, x_1, x_2\}$ .  
 (119)  $\{x_1, x_2, x_3, x_4\} = \{x_3, x_4, x_2, x_1\}$ .  
 (123)<sup>11</sup>  $\{x_1, x_2, x_3, x_4\} = \{x_4, x_2, x_3, x_1\}$ .  
 (125)<sup>12</sup>  $\{x_1, x_2, x_3, x_4\} = \{x_4, x_3, x_2, x_1\}$ .

<sup>7</sup> The proposition (97) has been removed.  
<sup>8</sup> The proposition (101) has been removed.  
<sup>9</sup> The proposition (106) has been removed.  
<sup>10</sup> The propositions (114) and (115) have been removed.  
<sup>11</sup> The propositions (120)–(122) have been removed.  
<sup>12</sup> The proposition (124) has been removed.

## REFERENCES

- [1] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics, Axiomatics*, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.

*Received January 8, 1989*

*Published January 2, 2004*

---