

# Basic Properties of Extended Real Numbers

Noboru Endou  
Shinshu University  
Nagano

Katsumi Wasaki  
Shinshu University  
Nagano

Yasunari Shidama  
Shinshu University  
Nagano

**Summary.** We introduce product, quotient and absolute value, and we prove some basic properties of extended real numbers.

MML Identifier: EXTREAL1.

WWW: <http://mizar.org/JFM/Vol12/extreal1.html>

The articles [1], [5], [2], [3], and [4] provide the notation and terminology for this paper.

## 1. PRELIMINARIES

In this paper  $x, y, z$  denote extended real numbers and  $a$  denotes a real number.

The following propositions are true:

- (1) If  $x \neq +\infty$  and  $x \neq -\infty$ , then  $x$  is a real number.
- (2)  $-\infty < +\infty$ .
- (3) If  $x < y$ , then  $x \neq +\infty$  and  $y \neq -\infty$ .
- (4)  $x = +\infty$  iff  $-x = -\infty$  and  $x = -\infty$  iff  $-x = +\infty$ .
- (5)  $x - -y = x + y$ .
- (7)<sup>1</sup> If  $x \neq -\infty$  and  $y \neq +\infty$  and  $x \leq y$ , then  $x \neq +\infty$  and  $y \neq -\infty$ .
- (8) Suppose  $x = +\infty$  and  $y = -\infty$  and  $x = -\infty$  and  $y = +\infty$  and  $y \neq +\infty$  or  $z \neq -\infty$  but  $y \neq -\infty$  or  $z \neq +\infty$  and  $x \neq +\infty$  or  $z \neq -\infty$  but  $x \neq -\infty$  or  $z \neq +\infty$ . Then  $(x + y) + z = x + (y + z)$ .
- (9)  $x + -x = 0_{\mathbb{R}}$ .
- (11)<sup>2</sup> Suppose  $x = +\infty$  and  $y = -\infty$  and  $x = -\infty$  and  $y = +\infty$  and  $y = +\infty$  and  $z = +\infty$  and  $y = -\infty$  and  $z = -\infty$  and  $x = +\infty$  and  $z = +\infty$  and  $x = -\infty$  and  $z = -\infty$ . Then  $(x + y) - z = x + (y - z)$ .

## 2. OPERATIONS OF MULTIPLICATION, QUOTIENT AND ABSOLUTE VALUE ON EXTENDED REAL NUMBERS

Let  $x, y$  be extended real numbers. The functor  $x \cdot y$  yields an extended real number and is defined by the conditions (Def. 1).

<sup>1</sup> The proposition (6) has been removed.

<sup>2</sup> The proposition (10) has been removed.

- (Def. 1)(i) There exist real numbers  $a, b$  such that  $x = a$  and  $y = b$  and  $x \cdot y = a \cdot b$ , or
- (ii)  $0_{\mathbb{R}} < x$  and  $y = +\infty$  or  $0_{\mathbb{R}} < y$  and  $x = +\infty$  or  $x < 0_{\mathbb{R}}$  and  $y = -\infty$  or  $y < 0_{\mathbb{R}}$  and  $x = -\infty$  but  $x \cdot y = +\infty$ , or
  - (iii)  $x < 0_{\mathbb{R}}$  and  $y = +\infty$  or  $y < 0_{\mathbb{R}}$  and  $x = +\infty$  or  $0_{\mathbb{R}} < x$  and  $y = -\infty$  or  $0_{\mathbb{R}} < y$  and  $x = -\infty$  but  $x \cdot y = -\infty$ , or
  - (iv)  $x = 0_{\mathbb{R}}$  or  $y = 0_{\mathbb{R}}$  but  $x \cdot y = 0_{\mathbb{R}}$ .

Next we state two propositions:

- (13)<sup>3</sup> For all extended real numbers  $x, y$  and for all real numbers  $a, b$  such that  $x = a$  and  $y = b$  holds  $x \cdot y = a \cdot b$ .
- (17)<sup>4</sup> For all extended real numbers  $x, y$  holds  $x \cdot y = y \cdot x$ .

Let  $x, y$  be extended real numbers. Let us observe that the functor  $x \cdot y$  is commutative.

One can prove the following propositions:

- (18) If  $x = a$ , then  $0 < a$  iff  $0_{\mathbb{R}} < x$ .
- (19) If  $x = a$ , then  $a < 0$  iff  $x < 0_{\mathbb{R}}$ .
- (20) If  $0_{\mathbb{R}} < x$  and  $0_{\mathbb{R}} < y$  or  $x < 0_{\mathbb{R}}$  and  $y < 0_{\mathbb{R}}$ , then  $0_{\mathbb{R}} < x \cdot y$ .
- (21) If  $0_{\mathbb{R}} < x$  and  $y < 0_{\mathbb{R}}$  or  $x < 0_{\mathbb{R}}$  and  $0_{\mathbb{R}} < y$ , then  $x \cdot y < 0_{\mathbb{R}}$ .
- (22)  $x \cdot y = 0_{\mathbb{R}}$  iff  $x = 0_{\mathbb{R}}$  or  $y = 0_{\mathbb{R}}$ .
- (23)  $(x \cdot y) \cdot z = x \cdot (y \cdot z)$ .
- (24)  $-0_{\mathbb{R}} = 0_{\mathbb{R}}$ .
- (25)  $0_{\mathbb{R}} < x$  iff  $-x < 0_{\mathbb{R}}$  and  $x < 0_{\mathbb{R}}$  iff  $0_{\mathbb{R}} < -x$ .
- (26)  $-x \cdot y = x \cdot -y$  and  $-x \cdot y = (-x) \cdot y$ .
- (27) If  $x \neq +\infty$  and  $x \neq -\infty$  and  $x \cdot y = +\infty$ , then  $y = +\infty$  or  $y = -\infty$ .
- (28) If  $x \neq +\infty$  and  $x \neq -\infty$  and  $x \cdot y = -\infty$ , then  $y = +\infty$  or  $y = -\infty$ .
- (29) If  $x \neq +\infty$  and  $x \neq -\infty$ , then  $x \cdot (y + z) = x \cdot y + x \cdot z$ .
- (30) If  $y \neq +\infty$  or  $z \neq +\infty$  but  $y \neq -\infty$  or  $z \neq -\infty$  and  $x \neq +\infty$  and  $x \neq -\infty$ , then  $x \cdot (y - z) = x \cdot y - x \cdot z$ .

Let  $x, y$  be extended real numbers. Let us assume that  $x = -\infty$  or  $x = +\infty$  but  $y = -\infty$  or  $y = +\infty$  but  $y \neq 0_{\mathbb{R}}$ . The functor  $\frac{x}{y}$  yielding an extended real number is defined by the conditions (Def. 2).

- (Def. 2)(i) There exist real numbers  $a, b$  such that  $x = a$  and  $y = b$  and  $\frac{x}{y} = \frac{a}{b}$ , or
- (ii)  $x = +\infty$  and  $0_{\mathbb{R}} < y$  or  $x = -\infty$  and  $y < 0_{\mathbb{R}}$  but  $\frac{x}{y} = +\infty$ , or
  - (iii)  $x = -\infty$  and  $0_{\mathbb{R}} < y$  or  $x = +\infty$  and  $y < 0_{\mathbb{R}}$  but  $\frac{x}{y} = -\infty$ , or
  - (iv)  $y = -\infty$  or  $y = +\infty$  but  $\frac{x}{y} = 0_{\mathbb{R}}$ .

We now state three propositions:

- (32)<sup>5</sup> Let  $x, y$  be extended real numbers. Suppose  $y \neq 0_{\mathbb{R}}$ . Let  $a, b$  be real numbers. If  $x = a$  and  $y = b$ , then  $\frac{x}{y} = \frac{a}{b}$ .

<sup>3</sup> The proposition (12) has been removed.

<sup>4</sup> The propositions (14)–(16) have been removed.

<sup>5</sup> The proposition (31) has been removed.

(33) For all extended real numbers  $x, y$  such that  $x \neq -\infty$  but  $x \neq +\infty$  but  $y = -\infty$  or  $y = +\infty$  holds  $\frac{x}{y} = 0_{\mathbb{R}}$ .

(34) For every extended real number  $x$  such that  $x \neq -\infty$  and  $x \neq +\infty$  and  $x \neq 0_{\mathbb{R}}$  holds  $\frac{x}{x} = 1$ .

Let  $x$  be an extended real number. The functor  $|x|$  yields an extended real number and is defined as follows:

$$\text{(Def. 3)} \quad |x| = \begin{cases} x, & \text{if } 0_{\mathbb{R}} \leq x, \\ -x, & \text{otherwise.} \end{cases}$$

One can prove the following propositions:

(36)<sup>6</sup> For every extended real number  $x$  such that  $0_{\mathbb{R}} < x$  holds  $|x| = x$ .

(37) For every extended real number  $x$  such that  $x < 0_{\mathbb{R}}$  holds  $|x| = -x$ .

(38) For all real numbers  $a, b$  holds  $\overline{\mathbb{R}}(a \cdot b) = \overline{\mathbb{R}}(a) \cdot \overline{\mathbb{R}}(b)$ .

(39) For all real numbers  $a, b$  such that  $b \neq 0$  holds  $\overline{\mathbb{R}}\left(\frac{a}{b}\right) = \frac{\overline{\mathbb{R}}(a)}{\overline{\mathbb{R}}(b)}$ .

(40) For all extended real numbers  $x, y$  such that  $x \leq y$  and  $x < +\infty$  and  $-\infty < y$  holds  $0_{\mathbb{R}} \leq y - x$ .

(41) For all extended real numbers  $x, y$  such that  $x < y$  and  $x < +\infty$  and  $-\infty < y$  holds  $0_{\mathbb{R}} < y - x$ .

(42) If  $x \leq y$  and  $0_{\mathbb{R}} \leq z$ , then  $x \cdot z \leq y \cdot z$ .

(43) If  $x \leq y$  and  $z \leq 0_{\mathbb{R}}$ , then  $y \cdot z \leq x \cdot z$ .

(44) If  $x < y$  and  $0_{\mathbb{R}} < z$  and  $z \neq +\infty$ , then  $x \cdot z < y \cdot z$ .

(45) If  $x < y$  and  $z < 0_{\mathbb{R}}$  and  $z \neq -\infty$ , then  $y \cdot z < x \cdot z$ .

(46) Suppose  $x$  is a real number and  $y$  is a real number. Then  $x < y$  if and only if there exist real numbers  $p, q$  such that  $p = x$  and  $q = y$  and  $p < q$ .

(47) If  $x \neq -\infty$  and  $y \neq +\infty$  and  $x \leq y$  and  $0_{\mathbb{R}} < z$ , then  $\frac{x}{z} \leq \frac{y}{z}$ .

(48) If  $x \leq y$  and  $0_{\mathbb{R}} < z$  and  $z \neq +\infty$ , then  $\frac{x}{z} \leq \frac{y}{z}$ .

(49) If  $x \neq -\infty$  and  $y \neq +\infty$  and  $x \leq y$  and  $z < 0_{\mathbb{R}}$ , then  $\frac{y}{z} \leq \frac{x}{z}$ .

(50) If  $x \leq y$  and  $z < 0_{\mathbb{R}}$  and  $z \neq -\infty$ , then  $\frac{y}{z} \leq \frac{x}{z}$ .

(51) If  $x < y$  and  $0_{\mathbb{R}} < z$  and  $z \neq +\infty$ , then  $\frac{x}{z} < \frac{y}{z}$ .

(52) If  $x < y$  and  $z < 0_{\mathbb{R}}$  and  $z \neq -\infty$ , then  $\frac{y}{z} < \frac{x}{z}$ .

## REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinal1.html>.
- [2] Józef Białas. Infimum and supremum of the set of real numbers. Measure theory. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/supinf\\_1.html](http://mizar.org/JFM/Vol2/supinf_1.html).
- [3] Józef Białas. Series of positive real numbers. Measure theory. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/supinf\\_2.html](http://mizar.org/JFM/Vol2/supinf_2.html).
- [4] Józef Białas. Some properties of the intervals. *Journal of Formalized Mathematics*, 6, 1994. <http://mizar.org/JFM/Vol6/measure6.html>.

<sup>6</sup> The proposition (35) has been removed.

- [5] Krzysztof Hryniewiecki. Basic properties of real numbers. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/real\\_1.html](http://mizar.org/JFM/Vol1/real_1.html).

*Received September 7, 2000*

*Published January 2, 2004*

---