

# Real Function One-Side Differentiability

Ewa Burakowska  
Warsaw University  
Białystok

Beata Madras  
Warsaw University  
Białystok

**Summary.** We define real function one-side differentiability and one-side continuity. Main properties of one-side differentiability function are proved. Connections between one-side differential and differential real function at the point are demonstrated.

MML Identifier: FDIFF\_3.

WWW: [http://mizar.org/JFM/Vol3/fdiff\\_3.html](http://mizar.org/JFM/Vol3/fdiff_3.html)

The articles [11], [1], [12], [2], [14], [5], [3], [4], [13], [7], [8], [10], [9], and [6] provide the notation and terminology for this paper.

For simplicity, we adopt the following rules:  $h, h_1, h_2$  denote convergent to 0 sequences of real numbers,  $c$  denotes a constant sequence of real numbers,  $f, f_1, f_2$  denote partial functions from  $\mathbb{R}$  to  $\mathbb{R}$ ,  $x_0, r, r_1, g, g_1, g_2$  denote real numbers,  $n$  denotes a natural number, and  $a$  denotes a sequence of real numbers.

One can prove the following propositions:

- (1) If there exists  $r$  such that  $r > 0$  and  $[x_0 - r, x_0] \subseteq \text{dom } f$ , then there exist  $h, c$  such that  $\text{rng } c = \{x_0\}$  and  $\text{rng}(h + c) \subseteq \text{dom } f$  and for every  $n$  holds  $h(n) < 0$ .
- (2) If there exists  $r$  such that  $r > 0$  and  $[x_0, x_0 + r] \subseteq \text{dom } f$ , then there exist  $h, c$  such that  $\text{rng } c = \{x_0\}$  and  $\text{rng}(h + c) \subseteq \text{dom } f$  and for every  $n$  holds  $h(n) > 0$ .
- (3) Suppose for all  $h, c$  such that  $\text{rng } c = \{x_0\}$  and  $\text{rng}(h + c) \subseteq \text{dom } f$  and for every  $n$  holds  $h(n) < 0$  holds  $h^{-1}(f \cdot (h + c) - f \cdot c)$  is convergent and  $\{x_0\} \subseteq \text{dom } f$ . Let given  $h_1, h_2, c$ . Suppose  $\text{rng } c = \{x_0\}$  and  $\text{rng}(h_1 + c) \subseteq \text{dom } f$  and for every  $n$  holds  $h_1(n) < 0$  and  $\text{rng}(h_2 + c) \subseteq \text{dom } f$  and for every  $n$  holds  $h_2(n) < 0$ . Then  $\lim(h_1^{-1}(f \cdot (h_1 + c) - f \cdot c)) = \lim(h_2^{-1}(f \cdot (h_2 + c) - f \cdot c))$ .
- (4) Suppose for all  $h, c$  such that  $\text{rng } c = \{x_0\}$  and  $\text{rng}(h + c) \subseteq \text{dom } f$  and for every  $n$  holds  $h(n) > 0$  holds  $h^{-1}(f \cdot (h + c) - f \cdot c)$  is convergent and  $\{x_0\} \subseteq \text{dom } f$ . Let given  $h_1, h_2, c$ . Suppose  $\text{rng } c = \{x_0\}$  and  $\text{rng}(h_1 + c) \subseteq \text{dom } f$  and  $\text{rng}(h_2 + c) \subseteq \text{dom } f$  and for every  $n$  holds  $h_1(n) > 0$  and for every  $n$  holds  $h_2(n) > 0$ . Then  $\lim(h_1^{-1}(f \cdot (h_1 + c) - f \cdot c)) = \lim(h_2^{-1}(f \cdot (h_2 + c) - f \cdot c))$ .

Let us consider  $f, x_0$ . We say that  $f$  is left continuous in  $x_0$  if and only if:

(Def. 1)  $x_0 \in \text{dom } f$  and for every  $a$  such that  $\text{rng } a \subseteq ]-\infty, x_0[ \cap \text{dom } f$  and  $a$  is convergent and  $\lim a = x_0$  holds  $f \cdot a$  is convergent and  $f(x_0) = \lim(f \cdot a)$ .

We say that  $f$  is right continuous in  $x_0$  if and only if:

(Def. 2)  $x_0 \in \text{dom } f$  and for every  $a$  such that  $\text{rng } a \subseteq ]x_0, +\infty[ \cap \text{dom } f$  and  $a$  is convergent and  $\lim a = x_0$  holds  $f \cdot a$  is convergent and  $f(x_0) = \lim(f \cdot a)$ .

We say that  $f$  is right differentiable in  $x_0$  if and only if the conditions (Def. 3) are satisfied.

- (Def. 3)(i) There exists  $r$  such that  $r > 0$  and  $[x_0, x_0 + r] \subseteq \text{dom } f$ , and  
 (ii) for all  $h, c$  such that  $\text{rng } c = \{x_0\}$  and  $\text{rng}(h+c) \subseteq \text{dom } f$  and for every  $n$  holds  $h(n) > 0$  holds  $h^{-1}(f \cdot (h+c) - f \cdot c)$  is convergent.

We say that  $f$  is left differentiable in  $x_0$  if and only if the conditions (Def. 4) are satisfied.

- (Def. 4)(i) There exists  $r$  such that  $r > 0$  and  $[x_0 - r, x_0] \subseteq \text{dom } f$ , and  
 (ii) for all  $h, c$  such that  $\text{rng } c = \{x_0\}$  and  $\text{rng}(h+c) \subseteq \text{dom } f$  and for every  $n$  holds  $h(n) < 0$  holds  $h^{-1}(f \cdot (h+c) - f \cdot c)$  is convergent.

We now state four propositions:

- (5) If  $f$  is left differentiable in  $x_0$ , then  $f$  is left continuous in  $x_0$ .  
 (6) Suppose  $f$  is left continuous in  $x_0$  and  $f(x_0) \neq g_2$  and there exists  $r$  such that  $r > 0$  and  $[x_0 - r, x_0] \subseteq \text{dom } f$ . Then there exists  $r_1$  such that  $r_1 > 0$  and  $[x_0 - r_1, x_0] \subseteq \text{dom } f$  and for every  $g$  such that  $g \in [x_0 - r_1, x_0]$  holds  $f(g) \neq g_2$ .  
 (7) If  $f$  is right differentiable in  $x_0$ , then  $f$  is right continuous in  $x_0$ .  
 (8) Suppose  $f$  is right continuous in  $x_0$  and  $f(x_0) \neq g_2$  and there exists  $r$  such that  $r > 0$  and  $[x_0, x_0 + r] \subseteq \text{dom } f$ . Then there exists  $r_1$  such that  $r_1 > 0$  and  $[x_0, x_0 + r_1] \subseteq \text{dom } f$  and for every  $g$  such that  $g \in [x_0, x_0 + r_1]$  holds  $f(g) \neq g_2$ .

Let us consider  $x_0, f$ . Let us assume that  $f$  is left differentiable in  $x_0$ . The functor  $f'_-(x_0)$  yields a real number and is defined by:

- (Def. 5) For all  $h, c$  such that  $\text{rng } c = \{x_0\}$  and  $\text{rng}(h+c) \subseteq \text{dom } f$  and for every  $n$  holds  $h(n) < 0$  holds  $f'_-(x_0) = \lim(h^{-1}(f \cdot (h+c) - f \cdot c))$ .

Let us consider  $x_0, f$ . Let us assume that  $f$  is right differentiable in  $x_0$ . The functor  $f'_+(x_0)$  yielding a real number is defined by:

- (Def. 6) For all  $h, c$  such that  $\text{rng } c = \{x_0\}$  and  $\text{rng}(h+c) \subseteq \text{dom } f$  and for every  $n$  holds  $h(n) > 0$  holds  $f'_+(x_0) = \lim(h^{-1}(f \cdot (h+c) - f \cdot c))$ .

The following propositions are true:

- (9)  $f$  is left differentiable in  $x_0$  and  $f'_-(x_0) = g$  if and only if the following conditions are satisfied:  
 (i) there exists  $r$  such that  $0 < r$  and  $[x_0 - r, x_0] \subseteq \text{dom } f$ , and  
 (ii) for all  $h, c$  such that  $\text{rng } c = \{x_0\}$  and  $\text{rng}(h+c) \subseteq \text{dom } f$  and for every  $n$  holds  $h(n) < 0$  holds  $h^{-1}(f \cdot (h+c) - f \cdot c)$  is convergent and  $\lim(h^{-1}(f \cdot (h+c) - f \cdot c)) = g$ .  
 (10) Suppose  $f_1$  is left differentiable in  $x_0$  and  $f_2$  is left differentiable in  $x_0$ . Then  $f_1 + f_2$  is left differentiable in  $x_0$  and  $(f_1 + f_2)'_-(x_0) = f_1'_-(x_0) + f_2'_-(x_0)$ .  
 (11) Suppose  $f_1$  is left differentiable in  $x_0$  and  $f_2$  is left differentiable in  $x_0$ . Then  $f_1 - f_2$  is left differentiable in  $x_0$  and  $(f_1 - f_2)'_-(x_0) = f_1'_-(x_0) - f_2'_-(x_0)$ .  
 (12) Suppose  $f_1$  is left differentiable in  $x_0$  and  $f_2$  is left differentiable in  $x_0$ . Then  $f_1 f_2$  is left differentiable in  $x_0$  and  $(f_1 f_2)'_-(x_0) = f_1'_-(x_0) \cdot f_2(x_0) + f_2'_-(x_0) \cdot f_1(x_0)$ .  
 (13) Suppose  $f_1$  is left differentiable in  $x_0$  and  $f_2$  is left differentiable in  $x_0$  and  $f_2(x_0) \neq 0$ . Then  $\frac{f_1}{f_2}$  is left differentiable in  $x_0$  and  $(\frac{f_1}{f_2})'_-(x_0) = \frac{f_1'_-(x_0) \cdot f_2(x_0) - f_2'_-(x_0) \cdot f_1(x_0)}{f_2(x_0)^2}$ .  
 (14) If  $f$  is left differentiable in  $x_0$  and  $f(x_0) \neq 0$ , then  $\frac{1}{f}$  is left differentiable in  $x_0$  and  $(\frac{1}{f})'_-(x_0) = -\frac{f'_-(x_0)}{f(x_0)^2}$ .

- (15)  $f$  is right differentiable in  $x_0$  and  $f'_+(x_0) = g_1$  if and only if the following conditions are satisfied:
- there exists  $r$  such that  $r > 0$  and  $[x_0, x_0 + r] \subseteq \text{dom } f$ , and
  - for all  $h, c$  such that  $\text{rng } c = \{x_0\}$  and  $\text{rng}(h + c) \subseteq \text{dom } f$  and for every  $n$  holds  $h(n) > 0$  holds  $h^{-1}(f \cdot (h + c) - f \cdot c)$  is convergent and  $\lim(h^{-1}(f \cdot (h + c) - f \cdot c)) = g_1$ .
- (16) Suppose  $f_1$  is right differentiable in  $x_0$  and  $f_2$  is right differentiable in  $x_0$ . Then  $f_1 + f_2$  is right differentiable in  $x_0$  and  $(f_1 + f_2)'_+(x_0) = f'_{1+}(x_0) + f'_{2+}(x_0)$ .
- (17) Suppose  $f_1$  is right differentiable in  $x_0$  and  $f_2$  is right differentiable in  $x_0$ . Then  $f_1 - f_2$  is right differentiable in  $x_0$  and  $(f_1 - f_2)'_+(x_0) = f'_{1+}(x_0) - f'_{2+}(x_0)$ .
- (18) Suppose  $f_1$  is right differentiable in  $x_0$  and  $f_2$  is right differentiable in  $x_0$ . Then  $f_1 f_2$  is right differentiable in  $x_0$  and  $(f_1 f_2)'_+(x_0) = f'_{1+}(x_0) \cdot f_2(x_0) + f'_{2+}(x_0) \cdot f_1(x_0)$ .
- (19) Suppose  $f_1$  is right differentiable in  $x_0$  and  $f_2$  is right differentiable in  $x_0$  and  $f_2(x_0) \neq 0$ . Then  $\frac{f_1}{f_2}$  is right differentiable in  $x_0$  and  $(\frac{f_1}{f_2})'_+(x_0) = \frac{f'_{1+}(x_0) \cdot f_2(x_0) - f'_{2+}(x_0) \cdot f_1(x_0)}{f_2(x_0)^2}$ .
- (20) If  $f$  is right differentiable in  $x_0$  and  $f(x_0) \neq 0$ , then  $\frac{1}{f}$  is right differentiable in  $x_0$  and  $(\frac{1}{f})'_+(x_0) = -\frac{f'_+(x_0)}{f(x_0)^2}$ .
- (21) Suppose  $f$  is right differentiable in  $x_0$  and left differentiable in  $x_0$  and  $f'_+(x_0) = f'_-(x_0)$ . Then  $f$  is differentiable in  $x_0$  and  $f'(x_0) = f'_+(x_0)$  and  $f'(x_0) = f'_-(x_0)$ .
- (22) Suppose  $f$  is differentiable in  $x_0$ . Then  $f$  is right differentiable in  $x_0$  and left differentiable in  $x_0$  and  $f'(x_0) = f'_+(x_0)$  and  $f'(x_0) = f'_-(x_0)$ .

## REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinal1.html>.
- [2] Krzysztof Hryniewiecki. Basic properties of real numbers. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/real\\_1.html](http://mizar.org/JFM/Vol1/real_1.html).
- [3] Jarosław Kotowicz. Convergent sequences and the limit of sequences. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/seq\\_2.html](http://mizar.org/JFM/Vol1/seq_2.html).
- [4] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/seqm\\_3.html](http://mizar.org/JFM/Vol1/seqm_3.html).
- [5] Jarosław Kotowicz. Real sequences and basic operations on them. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/seq\\_1.html](http://mizar.org/JFM/Vol1/seq_1.html).
- [6] Jarosław Kotowicz. The limit of a real function at infinity. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol2/limfunc1.html>.
- [7] Jarosław Kotowicz. Partial functions from a domain to the set of real numbers. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol2/rfunc1.html>.
- [8] Jarosław Kotowicz. Properties of real functions. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol2/rfunc2.html>.
- [9] Konrad Raczkowski and Paweł Sadowski. Real function differentiability. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/fdiff\\_1.html](http://mizar.org/JFM/Vol2/fdiff_1.html).
- [10] Konrad Raczkowski and Paweł Sadowski. Topological properties of subsets in real numbers. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/rcomp\\_1.html](http://mizar.org/JFM/Vol2/rcomp_1.html).
- [11] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [12] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [13] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers operations: min, max, square, and square root. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/square\\_1.html](http://mizar.org/JFM/Vol1/square_1.html).

- [14] Edmund Woronowicz. Relations defined on sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll/relset\\_1.html](http://mizar.org/JFM/Voll/relset_1.html).

*Received December 12, 1991*

*Published January 2, 2004*

---