

Free Universal Algebra Construction

Beata Perkowski
Warsaw University
Białystok

Summary. A construction of the free universal algebra with fixed signature and a given set of generators.

MML Identifier: FREEALG.

WWW: <http://mizar.org/JFM/Vol5/freealg.html>

The articles [15], [18], [16], [19], [7], [9], [4], [8], [14], [10], [1], [2], [3], [17], [13], [6], [11], [5], and [12] provide the notation and terminology for this paper.

1. PRELIMINARIES

In this paper n denotes a natural number.

Let I_1 be a set. We say that I_1 is missing \mathbb{N} if and only if:

(Def. 1) I_1 misses \mathbb{N} .

One can check that there exists a set which is non empty and missing \mathbb{N} .

Let I_1 be a finite sequence. We say that I_1 has zero if and only if:

(Def. 2) $0 \in \text{rng } I_1$.

We introduce I_1 is without zero as an antonym of I_1 has zero.

Let us note that there exists a finite sequence of elements of \mathbb{N} which is non empty and has zero and there exists a finite sequence of elements of \mathbb{N} which is non empty and without zero.

2. FREE UNIVERSAL ALGEBRA — GENERAL NOTIONS

Let U_1 be a universal algebra and let n be a natural number. Let us assume that $n \in \text{dom}$ (the characteristic of U_1). The functor $\text{oper}(n, U_1)$ yields an operation of U_1 and is defined by:

(Def. 4)¹ $\text{oper}(n, U_1) = (\text{the characteristic of } U_1)(n)$.

Let U_0 be a universal algebra. A subset of U_0 is called a generator set of U_0 if:

(Def. 5) The carrier of $\text{Gen}^{\text{UA}}(\text{it}) = \text{the carrier of } U_0$.

Let U_0 be a universal algebra and let I_1 be a generator set of U_0 . We say that I_1 is free if and only if the condition (Def. 6) is satisfied.

¹ The definition (Def. 3) has been removed.

(Def. 6) Let U_1 be a universal algebra. Suppose U_0 and U_1 are similar. Let f be a function from I_1 into the carrier of U_1 . Then there exists a function h from U_0 into U_1 such that h is a homomorphism of U_0 into U_1 and $h|I_1 = f$.

Let I_1 be a universal algebra. We say that I_1 is free if and only if:

(Def. 7) There exists a generator set of I_1 which is free.

Let us observe that there exists a universal algebra which is free and strict.

Let U_0 be a free universal algebra. Observe that there exists a generator set of U_0 which is free.

We now state the proposition

(1) Let U_0 be a strict universal algebra and A be a subset of U_0 . Then A is a generator set of U_0 if and only if $\text{Gen}^{\text{UA}}(A) = U_0$.

3. CONSTRUCTION OF DECORATED TREE STRUCTURE FOR FREE UNIVERSAL ALGEBRA

Let f be a non empty finite sequence of elements of \mathbb{N} and let X be a set. The functor $\text{REL}(f, X)$ yielding a relation between $\text{dom } f \cup X$ and $(\text{dom } f \cup X)^*$ is defined by:

(Def. 8) For every element a of $\text{dom } f \cup X$ and for every element b of $(\text{dom } f \cup X)^*$ holds $\langle a, b \rangle \in \text{REL}(f, X)$ iff $a \in \text{dom } f$ and $f(a) = \text{len } b$.

Let f be a non empty finite sequence of elements of \mathbb{N} and let X be a set. The functor $\text{DTConUA}(f, X)$ yielding a strict tree construction structure is defined as follows:

(Def. 9) $\text{DTConUA}(f, X) = \langle \text{dom } f \cup X, \text{REL}(f, X) \rangle$.

Let f be a non empty finite sequence of elements of \mathbb{N} and let X be a set. One can check that $\text{DTConUA}(f, X)$ is non empty.

The following propositions are true:

(2) Let f be a non empty finite sequence of elements of \mathbb{N} and X be a set. Then the terminals of $\text{DTConUA}(f, X) \subseteq X$ and the nonterminals of $\text{DTConUA}(f, X) = \text{dom } f$.

(3) Let f be a non empty finite sequence of elements of \mathbb{N} and X be a missing \mathbb{N} set. Then the terminals of $\text{DTConUA}(f, X) = X$.

Let f be a non empty finite sequence of elements of \mathbb{N} and let X be a set. Note that $\text{DTConUA}(f, X)$ has nonterminals.

Let f be a non empty finite sequence of elements of \mathbb{N} with zero and let X be a set. Note that $\text{DTConUA}(f, X)$ has nonterminals and useful nonterminals.

Let f be a non empty finite sequence of elements of \mathbb{N} and let D be a missing \mathbb{N} non empty set. Note that $\text{DTConUA}(f, D)$ has terminals, nonterminals, and useful nonterminals.

Let f be a non empty finite sequence of elements of \mathbb{N} , let X be a set, and let n be a natural number. Let us assume that $n \in \text{dom } f$. The functor $\text{Sym}(n, f, X)$ yields a symbol of $\text{DTConUA}(f, X)$ and is defined as follows:

(Def. 10) $\text{Sym}(n, f, X) = n$.

4. CONSTRUCTION OF FREE UNIVERSAL ALGEBRA FOR NON-EMPTY SET OF GENERATORS AND GIVEN SIGNATURE

Let f be a non empty finite sequence of elements of \mathbb{N} **qua** non empty set, let D be a missing \mathbb{N} non empty set, and let n be a natural number. Let us assume that $n \in \text{dom } f$. The functor $\text{FreeOpNSG}(n, f, D)$ yielding a homogeneous quasi total non empty partial function from $\text{TS}(\text{DTConUA}(f, D))^*$ to $\text{TS}(\text{DTConUA}(f, D))$ is defined by the conditions (Def. 11).

(Def. 11)(i) $\text{dom FreeOpNSG}(n, f, D) = \text{TS}(\text{DTConUA}(f, D))^{f_n}$, and

(ii) for every finite sequence p of elements of $\text{TS}(\text{DTConUA}(f, D))$ such that $p \in \text{dom FreeOpNSG}(n, f, D)$ holds $(\text{FreeOpNSG}(n, f, D))(p) = \text{Sym}(n, f, D)\text{-tree}(p)$.

Let f be a non empty finite sequence of elements of \mathbb{N} and let D be a missing \mathbb{N} non empty set. The functor $\text{FreeOpSeqNSG}(f, D)$ yields a finite sequence of operational functions of $\text{TS}(\text{DTConUA}(f, D))$ and is defined by:

(Def. 12) $\text{lenFreeOpSeqNSG}(f, D) = \text{len } f$ and for every n such that $n \in \text{domFreeOpSeqNSG}(f, D)$ holds $(\text{FreeOpSeqNSG}(f, D))(n) = \text{FreeOpNSG}(n, f, D)$.

Let f be a non empty finite sequence of elements of \mathbb{N} and let D be a missing \mathbb{N} non empty set. The functor $\text{FreeUnivAlgNSG}(f, D)$ yields a strict universal algebra and is defined as follows:

(Def. 13) $\text{FreeUnivAlgNSG}(f, D) = \langle \text{TS}(\text{DTConUA}(f, D)), \text{FreeOpSeqNSG}(f, D) \rangle$.

The following proposition is true

(4) For every non empty finite sequence f of elements of \mathbb{N} and for every missing \mathbb{N} non empty set D holds signature $\text{FreeUnivAlgNSG}(f, D) = f$.

Let f be a non empty finite sequence of elements of \mathbb{N} and let D be a non empty missing \mathbb{N} set. The functor $\text{FreeGenSetNSG}(f, D)$ yielding a subset of $\text{FreeUnivAlgNSG}(f, D)$ is defined as follows:

(Def. 14) $\text{FreeGenSetNSG}(f, D) = \{\text{the root tree of } s; s \text{ ranges over symbols of } \text{DTConUA}(f, D); s \in \text{the terminals of } \text{DTConUA}(f, D)\}$.

One can prove the following proposition

(5) Let f be a non empty finite sequence of elements of \mathbb{N} and D be a non empty missing \mathbb{N} set. Then $\text{FreeGenSetNSG}(f, D)$ is non empty.

Let f be a non empty finite sequence of elements of \mathbb{N} **qua** non empty set and let D be a non empty missing \mathbb{N} set. Then $\text{FreeGenSetNSG}(f, D)$ is a generator set of $\text{FreeUnivAlgNSG}(f, D)$.

Let f be a non empty finite sequence of elements of \mathbb{N} , let D be a non empty missing \mathbb{N} set, let C be a non empty set, let s be a symbol of $\text{DTConUA}(f, D)$, and let F be a function from $\text{FreeGenSetNSG}(f, D)$ into C . Let us assume that $s \in \text{the terminals of } \text{DTConUA}(f, D)$. The functor $\pi_s F$ yields an element of C and is defined as follows:

(Def. 15) $\pi_s F = F(\text{the root tree of } s)$.

Let f be a non empty finite sequence of elements of \mathbb{N} , let D be a set, and let s be a symbol of $\text{DTConUA}(f, D)$. Let us assume that there exists a finite sequence p such that $s \Rightarrow p$. The functor $@_s$ yields a natural number and is defined by:

(Def. 16) $@_s = s$.

The following proposition is true

(6) For every non empty finite sequence f of elements of \mathbb{N} and for every non empty missing \mathbb{N} set D holds $\text{FreeGenSetNSG}(f, D)$ is free.

Let f be a non empty finite sequence of elements of \mathbb{N} and let D be a non empty missing \mathbb{N} set. Note that $\text{FreeUnivAlgNSG}(f, D)$ is free.

Let f be a non empty finite sequence of elements of \mathbb{N} and let D be a non empty missing \mathbb{N} set. Then $\text{FreeGenSetNSG}(f, D)$ is a free generator set of $\text{FreeUnivAlgNSG}(f, D)$.

5. CONSTRUCTION OF FREE UNIVERSAL ALGEBRA AND SET OF GENERATORS

Let f be a non empty finite sequence of elements of \mathbb{N} **qua** non empty set with zero, let D be a missing \mathbb{N} set, and let n be a natural number. Let us assume that $n \in \text{dom } f$. The functor $\text{FreeOpZAO}(n, f, D)$ yields a homogeneous quasi total non empty partial function from $\text{TS}(\text{DTConUA}(f, D))^*$ to $\text{TS}(\text{DTConUA}(f, D))$ and is defined by the conditions (Def. 17).

- (Def. 17)(i) $\text{domFreeOpZAO}(n, f, D) = \text{TS}(\text{DTConUA}(f, D))^{f^n}$, and
(ii) for every finite sequence p of elements of $\text{TS}(\text{DTConUA}(f, D))$ such that $p \in \text{domFreeOpZAO}(n, f, D)$ holds $(\text{FreeOpZAO}(n, f, D))(p) = \text{Sym}(n, f, D)\text{-tree}(p)$.

Let f be a non empty finite sequence of elements of \mathbb{N} with zero and let D be a missing \mathbb{N} set. The functor $\text{FreeOpSeqZAO}(f, D)$ yields a finite sequence of operational functions of $\text{TS}(\text{DTConUA}(f, D))$ and is defined by:

- (Def. 18) $\text{lenFreeOpSeqZAO}(f, D) = \text{len } f$ and for every n such that $n \in \text{domFreeOpSeqZAO}(f, D)$ holds $(\text{FreeOpSeqZAO}(f, D))(n) = \text{FreeOpZAO}(n, f, D)$.

Let f be a non empty finite sequence of elements of \mathbb{N} with zero and let D be a missing \mathbb{N} set. The functor $\text{FreeUnivAlgZAO}(f, D)$ yields a strict universal algebra and is defined by:

- (Def. 19) $\text{FreeUnivAlgZAO}(f, D) = \langle \text{TS}(\text{DTConUA}(f, D)), \text{FreeOpSeqZAO}(f, D) \rangle$.

Next we state three propositions:

- (7) For every non empty finite sequence f of elements of \mathbb{N} with zero and for every missing \mathbb{N} set D holds $\text{signatureFreeUnivAlgZAO}(f, D) = f$.
(8) Let f be a non empty finite sequence of elements of \mathbb{N} with zero and D be a missing \mathbb{N} set. Then $\text{FreeUnivAlgZAO}(f, D)$ has constants.
(9) For every non empty finite sequence f of elements of \mathbb{N} with zero and for every missing \mathbb{N} set D holds $\text{Constants}(\text{FreeUnivAlgZAO}(f, D)) \neq \emptyset$.

Let f be a non empty finite sequence of elements of \mathbb{N} with zero and let D be a missing \mathbb{N} set. The functor $\text{FreeGenSetZAO}(f, D)$ yielding a subset of $\text{FreeUnivAlgZAO}(f, D)$ is defined by:

- (Def. 20) $\text{FreeGenSetZAO}(f, D) = \{\text{the root tree of } s; s \text{ ranges over symbols of } \text{DTConUA}(f, D); s \in \text{the terminals of } \text{DTConUA}(f, D)\}$.

Let f be a non empty finite sequence of elements of \mathbb{N} **qua** non empty set with zero and let D be a missing \mathbb{N} set. Then $\text{FreeGenSetZAO}(f, D)$ is a generator set of $\text{FreeUnivAlgZAO}(f, D)$.

Let f be a non empty finite sequence of elements of \mathbb{N} with zero, let D be a missing \mathbb{N} set, let C be a non empty set, let s be a symbol of $\text{DTConUA}(f, D)$, and let F be a function from $\text{FreeGenSetZAO}(f, D)$ into C . Let us assume that $s \in \text{the terminals of } \text{DTConUA}(f, D)$. The functor $\pi_s F$ yields an element of C and is defined by:

- (Def. 21) $\pi_s F = F(\text{the root tree of } s)$.

We now state the proposition

- (10) For every non empty finite sequence f of elements of \mathbb{N} with zero and for every missing \mathbb{N} set D holds $\text{FreeGenSetZAO}(f, D)$ is free.

Let f be a non empty finite sequence of elements of \mathbb{N} with zero and let D be a missing \mathbb{N} set. One can check that $\text{FreeUnivAlgZAO}(f, D)$ is free.

Let f be a non empty finite sequence of elements of \mathbb{N} with zero and let D be a missing \mathbb{N} set. Then $\text{FreeGenSetZAO}(f, D)$ is a free generator set of $\text{FreeUnivAlgZAO}(f, D)$.

Let us observe that there exists a universal algebra which is strict and free and has constants.

REFERENCES

- [1] Grzegorz Bancerek. König's Lemma. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/trees_2.html.
[2] Grzegorz Bancerek. Sets and functions of trees and joining operations of trees. *Journal of Formalized Mathematics*, 4, 1992. http://mizar.org/JFM/Vol4/trees_3.html.
[3] Grzegorz Bancerek. Joining of decorated trees. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/trees_4.html.

- [4] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.
- [5] Grzegorz Bancerek and Piotr Rudnicki. On defining functions on trees. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/dtconstr.html>.
- [6] Ewa Burakowska. Subalgebras of the universal algebra. Lattices of subalgebras. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/unialg_2.html.
- [7] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [8] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [9] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/partfun1.html>.
- [10] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_2.html.
- [11] Patricia L. Carlson and Grzegorz Bancerek. Context-free grammar — part I. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/lang1.html>.
- [12] Małgorzata Korolkiewicz. Homomorphisms of algebras. Quotient universal algebra. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/alg_1.html.
- [13] Jarosław Kotowicz, Beata Madras, and Małgorzata Korolkiewicz. Basic notation of universal algebra. *Journal of Formalized Mathematics*, 4, 1992. http://mizar.org/JFM/Vol4/unialg_1.html.
- [14] Andrzej Trybulec. Binary operations applied to functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funcop_1.html.
- [15] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [16] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [17] Wojciech A. Trybulec. Pigeon hole principle. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_4.html.
- [18] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [19] Edmund Woronowicz. Relations defined on sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relset_1.html.

Received October 20, 1993

Published January 2, 2004
