

# Basic Functions and Operations on Functions

Czesław Byliński  
Warsaw University  
Białystok

**Summary.** We define the following mappings: the characteristic function of a subset of a set, the inclusion function (injection or embedding), the projections from a Cartesian product onto its arguments and diagonal function (inclusion of a set into its Cartesian square). Some operations on functions are also defined: the products of two functions (the complex function and the more general product-function), the function induced on power sets by the image and inverse-image. Some simple propositions related to the introduced notions are proved.

MML Identifier: FUNCT\_3.

WWW: [http://mizar.org/JFM/Vol1/funct\\_3.html](http://mizar.org/JFM/Vol1/funct_3.html)

The articles [4], [3], [5], [6], [7], [1], and [2] provide the notation and terminology for this paper.

We use the following convention:  $x, y, z, z_1, z_2$  denote sets,  $A, B, V, X, X_1, X_2, Y, Y_1, Y_2, Z$  denote sets, and  $C, C_1, C_2, D, D_1, D_2$  denote non empty sets.

One can prove the following propositions:

- (1) If  $A \subseteq Y$ , then  $\text{id}_A = \text{id}_Y \upharpoonright A$ .
- (2) For all functions  $f, g$  such that  $X \subseteq \text{dom}(g \cdot f)$  holds  $f^\circ X \subseteq \text{dom } g$ .
- (3) For all functions  $f, g$  such that  $X \subseteq \text{dom } f$  and  $f^\circ X \subseteq \text{dom } g$  holds  $X \subseteq \text{dom}(g \cdot f)$ .
- (4) For all functions  $f, g$  such that  $Y \subseteq \text{rng}(g \cdot f)$  and  $g$  is one-to-one holds  $g^{-1}(Y) \subseteq \text{rng } f$ .
- (5) For all functions  $f, g$  such that  $Y \subseteq \text{rng } g$  and  $g^{-1}(Y) \subseteq \text{rng } f$  holds  $Y \subseteq \text{rng}(g \cdot f)$ .

In this article we present several logical schemes. The scheme *FuncEx 3* deals with a set  $\mathcal{A}$ , a set  $\mathcal{B}$ , and a ternary predicate  $\mathcal{P}$ , and states that:

There exists a function  $f$  such that  $\text{dom } f = [:\mathcal{A}, \mathcal{B}:]$  and for all  $x, y$  such that  $x \in \mathcal{A}$  and  $y \in \mathcal{B}$  holds  $\mathcal{P}[x, y, f(\langle x, y \rangle)]$

provided the parameters satisfy the following conditions:

- For all  $x, y, z_1, z_2$  such that  $x \in \mathcal{A}$  and  $y \in \mathcal{B}$  and  $\mathcal{P}[x, y, z_1]$  and  $\mathcal{P}[x, y, z_2]$  holds  $z_1 = z_2$ , and
- For all  $x, y$  such that  $x \in \mathcal{A}$  and  $y \in \mathcal{B}$  there exists  $z$  such that  $\mathcal{P}[x, y, z]$ .

The scheme *Lambda 3* deals with a set  $\mathcal{A}$ , a set  $\mathcal{B}$ , and a binary functor  $\mathcal{F}$  yielding a set, and states that:

There exists a function  $f$  such that  $\text{dom } f = [:\mathcal{A}, \mathcal{B}:]$  and for all  $x, y$  such that  $x \in \mathcal{A}$  and  $y \in \mathcal{B}$  holds  $f(\langle x, y \rangle) = \mathcal{F}(x, y)$

for all values of the parameters.

The following proposition is true

- (6) Let  $f, g$  be functions. Suppose  $\text{dom } f = [X, Y]$  and  $\text{dom } g = [X, Y]$  and for all  $x, y$  such that  $x \in X$  and  $y \in Y$  holds  $f(\langle x, y \rangle) = g(\langle x, y \rangle)$ . Then  $f = g$ .

Let  $f$  be a function. The functor  ${}^\circ f$  yields a function and is defined as follows:

(Def. 1)  $\text{dom } {}^\circ f = 2^{\text{dom } f}$  and for every  $X$  such that  $X \subseteq \text{dom } f$  holds  $({}^\circ f)(X) = f^\circ X$ .

Next we state a number of propositions:

- (8)<sup>1</sup> For every function  $f$  such that  $X \in \text{dom } {}^\circ f$  holds  $({}^\circ f)(X) = f^\circ X$ .
- (9) For every function  $f$  holds  $({}^\circ f)(\emptyset) = \emptyset$ .
- (10) For every function  $f$  holds  $\text{rng } {}^\circ f \subseteq 2^{\text{rng } f}$ .
- (12)<sup>2</sup> For every function  $f$  holds  $({}^\circ f)^\circ A \subseteq 2^{\text{rng } f}$ .
- (13) For every function  $f$  holds  $({}^\circ f)^{-1}(B) \subseteq 2^{\text{dom } f}$ .
- (14) For every function  $f$  from  $X$  into  $D$  holds  $({}^\circ f)^{-1}(B) \subseteq 2^X$ .
- (15) For every function  $f$  holds  $\bigcup(({}^\circ f)^\circ A) \subseteq f^\circ \bigcup A$ .
- (16) For every function  $f$  such that  $A \subseteq 2^{\text{dom } f}$  holds  $f^\circ \bigcup A = \bigcup(({}^\circ f)^\circ A)$ .
- (17) For every function  $f$  from  $X$  into  $D$  such that  $A \subseteq 2^X$  holds  $f^\circ \bigcup A = \bigcup(({}^\circ f)^\circ A)$ .
- (18) For every function  $f$  holds  $\bigcup(({}^\circ f)^{-1}(B)) \subseteq f^{-1}(\bigcup B)$ .
- (19) For every function  $f$  such that  $B \subseteq 2^{\text{rng } f}$  holds  $f^{-1}(\bigcup B) = \bigcup(({}^\circ f)^{-1}(B))$ .
- (20) For all functions  $f, g$  holds  ${}^\circ(g \cdot f) = {}^\circ g \cdot {}^\circ f$ .
- (21) For every function  $f$  holds  ${}^\circ f$  is a function from  $2^{\text{dom } f}$  into  $2^{\text{rng } f}$ .
- (22) For every function  $f$  from  $X$  into  $Y$  such that if  $Y = \emptyset$ , then  $X = \emptyset$  holds  ${}^\circ f$  is a function from  $2^X$  into  $2^Y$ .

Let us consider  $X, D$  and let  $f$  be a function from  $X$  into  $D$ . Then  ${}^\circ f$  is a function from  $2^X$  into  $2^D$ .

Let  $f$  be a function. The functor  ${}^{-1}f$  yields a function and is defined as follows:

(Def. 2)  $\text{dom } {}^{-1}f = 2^{\text{rng } f}$  and for every  $Y$  such that  $Y \subseteq \text{rng } f$  holds  $({}^{-1}f)(Y) = f^{-1}(Y)$ .

One can prove the following propositions:

- (24)<sup>3</sup> For every function  $f$  such that  $Y \in \text{dom } {}^{-1}f$  holds  $({}^{-1}f)(Y) = f^{-1}(Y)$ .
- (25) For every function  $f$  holds  $\text{rng } {}^{-1}f \subseteq 2^{\text{dom } f}$ .
- (27)<sup>4</sup> For every function  $f$  holds  $({}^{-1}f)^\circ B \subseteq 2^{\text{dom } f}$ .
- (28) For every function  $f$  holds  $({}^{-1}f)^{-1}(A) \subseteq 2^{\text{rng } f}$ .
- (29) For every function  $f$  holds  $\bigcup(({}^{-1}f)^\circ B) \subseteq f^{-1}(\bigcup B)$ .
- (30) For every function  $f$  such that  $B \subseteq 2^{\text{rng } f}$  holds  $\bigcup(({}^{-1}f)^\circ B) = f^{-1}(\bigcup B)$ .
- (31) For every function  $f$  holds  $\bigcup(({}^{-1}f)^{-1}(A)) \subseteq f^\circ \bigcup A$ .

<sup>1</sup> The proposition (7) has been removed.

<sup>2</sup> The proposition (11) has been removed.

<sup>3</sup> The proposition (23) has been removed.

<sup>4</sup> The proposition (26) has been removed.

- (32) For every function  $f$  such that  $A \subseteq 2^{\text{dom}f}$  and  $f$  is one-to-one holds  $\bigcup((^{-1}f)^{-1}(A)) = f^{\circ} \bigcup A$ .
- (33) For every function  $f$  holds  $(^{-1}f)^{\circ}B \subseteq (^{\circ}f)^{-1}(B)$ .
- (34) For every function  $f$  such that  $f$  is one-to-one holds  $(^{-1}f)^{\circ}B = (^{\circ}f)^{-1}(B)$ .
- (35) For every function  $f$  and for every set  $A$  such that  $A \subseteq 2^{\text{dom}f}$  holds  $(^{-1}f)^{-1}(A) \subseteq (^{\circ}f)^{\circ}A$ .
- (36) For every function  $f$  and for every set  $A$  such that  $f$  is one-to-one holds  $(^{\circ}f)^{\circ}A \subseteq (^{-1}f)^{-1}(A)$ .
- (37) For every function  $f$  and for every set  $A$  such that  $f$  is one-to-one and  $A \subseteq 2^{\text{dom}f}$  holds  $(^{-1}f)^{-1}(A) = (^{\circ}f)^{\circ}A$ .
- (38) For all functions  $f, g$  such that  $g$  is one-to-one holds  $(^{-1}(g \cdot f)) = (^{-1}f) \cdot (^{-1}g)$ .
- (39) For every function  $f$  holds  $(^{-1}f)$  is a function from  $2^{\text{rng}f}$  into  $2^{\text{dom}f}$ .

Let us consider  $A, X$ . The functor  $\chi_{A,X}$  yields a function and is defined as follows:

(Def. 3)  $\text{dom}(\chi_{A,X}) = X$  and for every  $x$  such that  $x \in X$  holds if  $x \in A$ , then  $\chi_{A,X}(x) = 1$  and if  $x \notin A$ , then  $\chi_{A,X}(x) = 0$ .

We now state several propositions:

- (42)<sup>5</sup> If  $\chi_{A,X}(x) = 1$ , then  $x \in A$ .
- (43) If  $x \in X \setminus A$ , then  $\chi_{A,X}(x) = 0$ .
- (47)<sup>6</sup> If  $A \subseteq X$  and  $B \subseteq X$  and  $\chi_{A,X} = \chi_{B,X}$ , then  $A = B$ .
- (48)  $\text{rng}(\chi_{A,X}) \subseteq \{0, 1\}$ .
- (49) For every function  $f$  from  $X$  into  $\{0, 1\}$  holds  $f = \chi_{f^{-1}(\{1\}),X}$ .

Let us consider  $A, X$ . Then  $\chi_{A,X}$  is a function from  $X$  into  $\{0, 1\}$ .

Let us consider  $Y$  and let  $A$  be a subset of  $Y$ . The functor  $\underset{\hookrightarrow}{A}$  yields a function from  $A$  into  $Y$  and is defined by:

(Def. 4)  $\underset{\hookrightarrow}{A} = \text{id}_A$ .

The following four propositions are true:

- (53)<sup>7</sup> For every subset  $A$  of  $Y$  holds  $\underset{\hookrightarrow}{A} = \text{id}_Y \upharpoonright A$ .
- (54) For every subset  $A$  of  $Y$  holds  $\text{dom}(\underset{\hookrightarrow}{A}) = A$  and  $\text{rng}(\underset{\hookrightarrow}{A}) = A$ .
- (55) For every subset  $A$  of  $Y$  such that  $x \in A$  holds  $(\underset{\hookrightarrow}{A})(x) = x$ .
- (56) For every subset  $A$  of  $Y$  such that  $x \in A$  holds  $(\underset{\hookrightarrow}{A})(x) \in Y$ .

Let us consider  $X, Y$ . The functor  $\pi_1(X \times Y)$  yields a function and is defined as follows:

(Def. 5)  $\text{dom}\pi_1(X \times Y) = [X, Y:]$  and for all  $x, y$  such that  $x \in X$  and  $y \in Y$  holds  $\pi_1(X \times Y)(\langle x, y \rangle) = x$ .

The functor  $\pi_2(X \times Y)$  yields a function and is defined as follows:

(Def. 6)  $\text{dom}\pi_2(X \times Y) = [X, Y:]$  and for all  $x, y$  such that  $x \in X$  and  $y \in Y$  holds  $\pi_2(X \times Y)(\langle x, y \rangle) = y$ .

<sup>5</sup> The propositions (40) and (41) have been removed.

<sup>6</sup> The propositions (44)–(46) have been removed.

<sup>7</sup> The propositions (50)–(52) have been removed.

We now state four propositions:

$$(59)^8 \quad \text{rng } \pi_1(X \times Y) \subseteq X.$$

$$(60) \quad \text{If } Y \neq \emptyset, \text{ then } \text{rng } \pi_1(X \times Y) = X.$$

$$(61) \quad \text{rng } \pi_2(X \times Y) \subseteq Y.$$

$$(62) \quad \text{If } X \neq \emptyset, \text{ then } \text{rng } \pi_2(X \times Y) = Y.$$

Let us consider  $X, Y$ . Then  $\pi_1(X \times Y)$  is a function from  $[:X, Y:]$  into  $X$ . Then  $\pi_2(X \times Y)$  is a function from  $[:X, Y:]$  into  $Y$ .

Let us consider  $X$ . The functor  $\delta_X$  yielding a function is defined as follows:

$$(\text{Def. 7}) \quad \text{dom}(\delta_X) = X \text{ and for every } x \text{ such that } x \in X \text{ holds } \delta_X(x) = \langle x, x \rangle.$$

The following proposition is true

$$(66)^9 \quad \text{rng}(\delta_X) \subseteq [:X, X:].$$

Let us consider  $X$ . Then  $\delta_X$  is a function from  $X$  into  $[:X, X:]$ .

Let  $f, g$  be functions. The functor  $\langle f, g \rangle$  yielding a function is defined as follows:

$$(\text{Def. 8}) \quad \text{dom}\langle f, g \rangle = \text{dom } f \cap \text{dom } g \text{ and for every } x \text{ such that } x \in \text{dom}\langle f, g \rangle \text{ holds } \langle f, g \rangle(x) = \langle f(x), g(x) \rangle.$$

One can prove the following propositions:

$$(68)^{10} \quad \text{For all functions } f, g \text{ such that } x \in \text{dom } f \cap \text{dom } g \text{ holds } \langle f, g \rangle(x) = \langle f(x), g(x) \rangle.$$

$$(69) \quad \text{For all functions } f, g \text{ such that } \text{dom } f = X \text{ and } \text{dom } g = X \text{ and } x \in X \text{ holds } \langle f, g \rangle(x) = \langle f(x), g(x) \rangle.$$

$$(70) \quad \text{For all functions } f, g \text{ such that } \text{dom } f = X \text{ and } \text{dom } g = X \text{ holds } \text{dom}\langle f, g \rangle = X.$$

$$(71) \quad \text{For all functions } f, g \text{ holds } \text{rng}\langle f, g \rangle \subseteq [: \text{rng } f, \text{rng } g :].$$

$$(72) \quad \text{For all functions } f, g \text{ such that } \text{dom } f = \text{dom } g \text{ and } \text{rng } f \subseteq Y \text{ and } \text{rng } g \subseteq Z \text{ holds } \pi_1(Y \times Z) \cdot \langle f, g \rangle = f \text{ and } \pi_2(Y \times Z) \cdot \langle f, g \rangle = g.$$

$$(73) \quad \langle \pi_1(X \times Y), \pi_2(X \times Y) \rangle = \text{id}_{[:X, Y:]}$$

$$(74) \quad \text{For all functions } f, g, h, k \text{ such that } \text{dom } f = \text{dom } g \text{ and } \text{dom } k = \text{dom } h \text{ and } \langle f, g \rangle = \langle k, h \rangle \text{ holds } f = k \text{ and } g = h.$$

$$(75) \quad \text{For all functions } f, g, h \text{ holds } \langle f \cdot h, g \cdot h \rangle = \langle f, g \rangle \cdot h.$$

$$(76) \quad \text{For all functions } f, g \text{ holds } \langle f, g \rangle^\circ A \subseteq [: f^\circ A, g^\circ A :].$$

$$(77) \quad \text{For all functions } f, g \text{ holds } \langle f, g \rangle^{-1}([: B, C :]) = f^{-1}(B) \cap g^{-1}(C).$$

$$(78) \quad \text{Let } f \text{ be a function from } X \text{ into } Y \text{ and } g \text{ be a function from } X \text{ into } Z. \text{ Suppose if } Y = \emptyset, \text{ then } X = \emptyset \text{ and if } Z = \emptyset, \text{ then } X = \emptyset. \text{ Then } \langle f, g \rangle \text{ is a function from } X \text{ into } [: Y, Z :].$$

Let us consider  $X, D_1, D_2$ , let  $f_1$  be a function from  $X$  into  $D_1$ , and let  $f_2$  be a function from  $X$  into  $D_2$ . Then  $\langle f_1, f_2 \rangle$  is a function from  $X$  into  $[: D_1, D_2 :]$ .

Next we state several propositions:

$$(79) \quad \text{Let } f_1 \text{ be a function from } C \text{ into } D_1, f_2 \text{ be a function from } C \text{ into } D_2, \text{ and } c \text{ be an element of } C. \text{ Then } \langle f_1, f_2 \rangle(c) = \langle f_1(c), f_2(c) \rangle.$$

<sup>8</sup> The propositions (57) and (58) have been removed.

<sup>9</sup> The propositions (63)–(65) have been removed.

<sup>10</sup> The proposition (67) has been removed.

- (80) For every function  $f$  from  $X$  into  $Y$  and for every function  $g$  from  $X$  into  $Z$  holds  $\text{rng}\langle f, g \rangle \subseteq [Y, Z]$ .
- (81) Let  $f$  be a function from  $X$  into  $Y$  and  $g$  be a function from  $X$  into  $Z$ . Suppose if  $Y = \emptyset$ , then  $X = \emptyset$  and if  $Z = \emptyset$ , then  $X = \emptyset$ . Then  $\pi_1(Y \times Z) \cdot \langle f, g \rangle = f$  and  $\pi_2(Y \times Z) \cdot \langle f, g \rangle = g$ .
- (82) For every function  $f$  from  $X$  into  $D_1$  and for every function  $g$  from  $X$  into  $D_2$  holds  $\pi_1(D_1 \times D_2) \cdot \langle f, g \rangle = f$  and  $\pi_2(D_1 \times D_2) \cdot \langle f, g \rangle = g$ .
- (83) Let  $f_1, f_2$  be functions from  $X$  into  $Y$  and  $g_1, g_2$  be functions from  $X$  into  $Z$ . Suppose if  $Y = \emptyset$ , then  $X = \emptyset$  and if  $Z = \emptyset$ , then  $X = \emptyset$  and  $\langle f_1, g_1 \rangle = \langle f_2, g_2 \rangle$ . Then  $f_1 = f_2$  and  $g_1 = g_2$ .
- (84) Let  $f_1, f_2$  be functions from  $X$  into  $D_1$  and  $g_1, g_2$  be functions from  $X$  into  $D_2$ . If  $\langle f_1, g_1 \rangle = \langle f_2, g_2 \rangle$ , then  $f_1 = f_2$  and  $g_1 = g_2$ .

Let  $f, g$  be functions. The functor  $[f, g]$  yielding a function is defined by:

(Def. 9)  $\text{dom}[f, g] = [\text{dom } f, \text{dom } g]$  and for all  $x, y$  such that  $x \in \text{dom } f$  and  $y \in \text{dom } g$  holds  $[f, g](\langle x, y \rangle) = \langle f(x), g(y) \rangle$ .

We now state a number of propositions:

- (86)<sup>11</sup> For all functions  $f, g$  and for all  $x, y$  such that  $\langle x, y \rangle \in [\text{dom } f, \text{dom } g]$  holds  $[f, g](\langle x, y \rangle) = \langle f(x), g(y) \rangle$ .
- (87) For all functions  $f, g$  holds  $[f, g] = \langle f \cdot \pi_1(\text{dom } f \times \text{dom } g), g \cdot \pi_2(\text{dom } f \times \text{dom } g) \rangle$ .
- (88) For all functions  $f, g$  holds  $\text{rng}[f, g] = [\text{rng } f, \text{rng } g]$ .
- (89) For all functions  $f, g$  such that  $\text{dom } f = X$  and  $\text{dom } g = X$  holds  $\langle f, g \rangle = [f, g] \cdot \delta_X$ .
- (90)  $[id_X, id_Y] = id_{[X, Y]}$ .
- (91) For all functions  $f, g, h, k$  holds  $[f, h] \cdot \langle g, k \rangle = \langle f \cdot g, h \cdot k \rangle$ .
- (92) For all functions  $f, g, h, k$  holds  $[f, h] \cdot [g, k] = [f \cdot g, h \cdot k]$ .
- (93) For all functions  $f, g$  holds  $[f, g]^\circ [B, A] = [f^\circ B, g^\circ A]$ .
- (94) For all functions  $f, g$  holds  $[f, g]^{-1}([B, A]) = [f^{-1}(B), g^{-1}(A)]$ .
- (95) Let  $f$  be a function from  $X$  into  $Y$  and  $g$  be a function from  $V$  into  $Z$ . Then  $[f, g]$  is a function from  $[X, V]$  into  $[Y, Z]$ .

Let us consider  $X_1, X_2, Y_1, Y_2$ , let  $f_1$  be a function from  $X_1$  into  $Y_1$ , and let  $f_2$  be a function from  $X_2$  into  $Y_2$ . Then  $[f_1, f_2]$  is a function from  $[X_1, X_2]$  into  $[Y_1, Y_2]$ .

Next we state several propositions:

- (96) Let  $f_1$  be a function from  $C_1$  into  $D_1$ ,  $f_2$  be a function from  $C_2$  into  $D_2$ ,  $c_1$  be an element of  $C_1$ , and  $c_2$  be an element of  $C_2$ . Then  $[f_1, f_2](\langle c_1, c_2 \rangle) = \langle f_1(c_1), f_2(c_2) \rangle$ .
- (97) Let  $f_1$  be a function from  $X_1$  into  $Y_1$  and  $f_2$  be a function from  $X_2$  into  $Y_2$ . If if  $Y_1 = \emptyset$ , then  $X_1 = \emptyset$  and if  $Y_2 = \emptyset$ , then  $X_2 = \emptyset$ , then  $[f_1, f_2] = \langle f_1 \cdot \pi_1(X_1 \times X_2), f_2 \cdot \pi_2(X_1 \times X_2) \rangle$ .
- (98) For every function  $f_1$  from  $X_1$  into  $D_1$  and for every function  $f_2$  from  $X_2$  into  $D_2$  holds  $[f_1, f_2] = \langle f_1 \cdot \pi_1(X_1 \times X_2), f_2 \cdot \pi_2(X_1 \times X_2) \rangle$ .
- (99) Let  $f_1$  be a function from  $X$  into  $Y_1$  and  $f_2$  be a function from  $X$  into  $Y_2$ . If if  $Y_1 = \emptyset$ , then  $X = \emptyset$  and if  $Y_2 = \emptyset$ , then  $X = \emptyset$ , then  $\langle f_1, f_2 \rangle = [f_1, f_2] \cdot \delta_X$ .
- (100) For every function  $f_1$  from  $X$  into  $D_1$  and for every function  $f_2$  from  $X$  into  $D_2$  holds  $\langle f_1, f_2 \rangle = [f_1, f_2] \cdot \delta_X$ .

<sup>11</sup> The proposition (85) has been removed.

## REFERENCES

- [1] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll/funct\\_1.html](http://mizar.org/JFM/Voll/funct_1.html).
- [2] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll/funct\\_2.html](http://mizar.org/JFM/Voll/funct_2.html).
- [3] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll/zfmisc\\_1.html](http://mizar.org/JFM/Voll/zfmisc_1.html).
- [4] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [5] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll/subset\\_1.html](http://mizar.org/JFM/Voll/subset_1.html).
- [6] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll/relat\\_1.html](http://mizar.org/JFM/Voll/relat_1.html).
- [7] Edmund Woronowicz. Relations defined on sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Voll/relset\\_1.html](http://mizar.org/JFM/Voll/relset_1.html).

*Received May 9, 1989*

*Published January 2, 2004*

---