

Cartesian Product of Functions

Grzegorz Bancerek
Warsaw University
Białystok

Summary. A supplement of [3] and [2], i.e. some useful and explanatory properties of the product and also the curried and uncurried functions are shown. Besides, the functions yielding functions are considered: two different products and other operation of such functions are introduced. Finally, two facts are presented: quasi-distributivity of the power of the set to other one w.r.t. the union ($X^{\bigcup_x f(x)} \approx \prod_x X^{f(x)}$) and quasi-distributivity of the product w.r.t. the raising to the power ($\prod_x f(x)^X \approx (\prod_x f(x))^X$).

MML Identifier: FUNCT_6.

WWW: http://mizar.org/JFM/Vol3/funct_6.html

The articles [16], [15], [9], [17], [18], [6], [4], [13], [7], [8], [5], [1], [14], [10], [11], [2], [12], and [3] provide the notation and terminology for this paper.

1. PROPERTIES OF CARTESIAN PRODUCT

For simplicity, we use the following convention: $x, y, y_1, y_2, z, a, X, Y, Z, V_1, V_2$ are sets, f, g, h, h', f_1, f_2 are functions, i is a natural number, P is a permutation of X , D, D_1, D_2, D_3 are non empty sets, d_1 is an element of D_1 , d_2 is an element of D_2 , and d_3 is an element of D_3 .

We now state a number of propositions:

- (1) $x \in \prod\langle X \rangle$ iff there exists y such that $y \in X$ and $x = \langle y \rangle$.
- (2) $z \in \prod\langle X, Y \rangle$ iff there exist x, y such that $x \in X$ and $y \in Y$ and $z = \langle x, y \rangle$.
- (3) $a \in \prod\langle X, Y, Z \rangle$ iff there exist x, y, z such that $x \in X$ and $y \in Y$ and $z \in Z$ and $a = \langle x, y, z \rangle$.
- (4) $\prod\langle D \rangle = D^1$.
- (5) $\prod\langle D_1, D_2 \rangle = \{\langle d_1, d_2 \rangle\}$.
- (6) $\prod\langle D, D \rangle = D^2$.
- (7) $\prod\langle D_1, D_2, D_3 \rangle = \{\langle d_1, d_2, d_3 \rangle\}$.
- (8) $\prod\langle D, D, D \rangle = D^3$.
- (9) $\prod(i \mapsto D) = D^i$.
- (10) $\prod f \subseteq (\bigcup f)^{\text{dom } f}$.

2. CURRIED AND UNCURRIED FUNCTIONS OF SOME FUNCTIONS

We now state a number of propositions:

- (11) If $x \in \text{dom} \curvearrowright f$, then there exist y, z such that $x = \langle y, z \rangle$.
- (12) $\curvearrowright([X, Y] \rightarrow z) = [Y, X] \rightarrow z$.
- (13) $\text{curry } f = \text{curry}' \curvearrowright f$ and $\text{uncurry } f = \curvearrowright \text{uncurry}' f$.
- (14) If $[X, Y] \neq \emptyset$, then $\text{curry}([X, Y] \rightarrow z) = X \rightarrow (Y \rightarrow z)$ and $\text{curry}'([X, Y] \rightarrow z) = Y \rightarrow (X \rightarrow z)$.
- (15) $\text{uncurry}(X \rightarrow (Y \rightarrow z)) = [X, Y] \rightarrow z$ and $\text{uncurry}'(X \rightarrow (Y \rightarrow z)) = [Y, X] \rightarrow z$.
- (16) If $x \in \text{dom } f$ and $g = f(x)$, then $\text{rng } g \subseteq \text{rng uncurry } f$ and $\text{rng } g \subseteq \text{rng uncurry}' f$.
- (17) $\text{dom uncurry}(X \rightarrow f) = [X, \text{dom } f]$ and $\text{rng uncurry}(X \rightarrow f) \subseteq \text{rng } f$ and $\text{dom uncurry}'(X \rightarrow f) = [\text{dom } f, X]$ and $\text{rng uncurry}'(X \rightarrow f) \subseteq \text{rng } f$.
- (18) If $X \neq \emptyset$, then $\text{rng uncurry}(X \rightarrow f) = \text{rng } f$ and $\text{rng uncurry}'(X \rightarrow f) = \text{rng } f$.
- (19) If $[X, Y] \neq \emptyset$ and $f \in Z^{[X, Y]}$, then $\text{curry } f \in (Z^Y)^X$ and $\text{curry}' f \in (Z^X)^Y$.
- (20) If $f \in (Z^Y)^X$, then $\text{uncurry } f \in Z^{[X, Y]}$ and $\text{uncurry}' f \in Z^{[Y, X]}$.
- (21) If $\text{curry } f \in (Z^Y)^X$ or $\text{curry}' f \in (Z^X)^Y$ and if $\text{dom } f \subseteq [V_1, V_2]$, then $f \in Z^{[X, Y]}$.
- (22) If $\text{uncurry } f \in Z^{[X, Y]}$ or $\text{uncurry}' f \in Z^{[Y, X]}$ and if $\text{rng } f \subseteq V_1 \dot{\rightarrow} V_2$ and if $\text{dom } f = X$, then $f \in (Z^Y)^X$.
- (23) If $f \in [X, Y] \dot{\rightarrow} Z$, then $\text{curry } f \in X \dot{\rightarrow} (Y \dot{\rightarrow} Z)$ and $\text{curry}' f \in Y \dot{\rightarrow} (X \dot{\rightarrow} Z)$.
- (24) If $f \in X \dot{\rightarrow} (Y \dot{\rightarrow} Z)$, then $\text{uncurry } f \in [X, Y] \dot{\rightarrow} Z$ and $\text{uncurry}' f \in [Y, X] \dot{\rightarrow} Z$.
- (25) If $\text{curry } f \in X \dot{\rightarrow} (Y \dot{\rightarrow} Z)$ or $\text{curry}' f \in Y \dot{\rightarrow} (X \dot{\rightarrow} Z)$ and if $\text{dom } f \subseteq [V_1, V_2]$, then $f \in [X, Y] \dot{\rightarrow} Z$.
- (26) If $\text{uncurry } f \in [X, Y] \dot{\rightarrow} Z$ or $\text{uncurry}' f \in [Y, X] \dot{\rightarrow} Z$ and if $\text{rng } f \subseteq V_1 \dot{\rightarrow} V_2$ and if $\text{dom } f \subseteq X$, then $f \in X \dot{\rightarrow} (Y \dot{\rightarrow} Z)$.

3. FUNCTIONS YIELDING FUNCTIONS

Let X be a set. The functor $\text{Sub}_f X$ is defined as follows:

- (Def. 1) $x \in \text{Sub}_f X$ iff $x \in X$ and x is a function.

Next we state four propositions:

- (27) $\text{Sub}_f X \subseteq X$.
- (28) $x \in f^{-1}(\text{Sub}_f \text{rng } f)$ iff $x \in \text{dom } f$ and $f(x)$ is a function.
- (29) $\text{Sub}_f \emptyset = \emptyset$ and $\text{Sub}_f \{f\} = \{f\}$ and $\text{Sub}_f \{f, g\} = \{f, g\}$ and $\text{Sub}_f \{f, g, h\} = \{f, g, h\}$.
- (30) If $Y \subseteq \text{Sub}_f X$, then $\text{Sub}_f Y = Y$.

Let f be a function. The functor $\text{dom}_K f(K)$ yielding a function is defined as follows:

- (Def. 2) $\text{dom}(\text{dom}_K f(K)) = f^{-1}(\text{Sub}_f \text{rng } f)$ and for every x such that $x \in f^{-1}(\text{Sub}_f \text{rng } f)$ holds $(\text{dom}_K f(K))(x) = \pi_1(f(x))$.

The functor $\text{rng}_K f(K)$ yielding a function is defined by:

(Def. 3) $\text{dom}(\text{rng}_\kappa f(\kappa)) = f^{-1}(\text{Sub}_{\text{frng}} f)$ and for every x such that $x \in f^{-1}(\text{Sub}_{\text{frng}} f)$ holds $(\text{rng}_\kappa f(\kappa))(x) = \pi_2(f(x))$.

The functor $\cap f$ is defined as follows:

(Def. 4) $\cap f = \cap \text{rng } f$.

One can prove the following propositions:

- (31) If $x \in \text{dom } f$ and $g = f(x)$, then $x \in \text{dom}(\text{dom}_\kappa f(\kappa))$ and $(\text{dom}_\kappa f(\kappa))(x) = \text{dom } g$ and $x \in \text{dom}(\text{rng}_\kappa f(\kappa))$ and $(\text{rng}_\kappa f(\kappa))(x) = \text{rng } g$.
- (32) $\text{dom}_\kappa \emptyset(\kappa) = \emptyset$ and $\text{rng}_\kappa \emptyset(\kappa) = \emptyset$.
- (33) $\text{dom}_\kappa \langle f \rangle(\kappa) = \langle \text{dom } f \rangle$ and $\text{rng}_\kappa \langle f \rangle(\kappa) = \langle \text{rng } f \rangle$.
- (34) $\text{dom}_\kappa \langle f, g \rangle(\kappa) = \langle \text{dom } f, \text{dom } g \rangle$ and $\text{rng}_\kappa \langle f, g \rangle(\kappa) = \langle \text{rng } f, \text{rng } g \rangle$.
- (35) $\text{dom}_\kappa \langle f, g, h \rangle(\kappa) = \langle \text{dom } f, \text{dom } g, \text{dom } h \rangle$ and $\text{rng}_\kappa \langle f, g, h \rangle(\kappa) = \langle \text{rng } f, \text{rng } g, \text{rng } h \rangle$.
- (36) $\text{dom}_\kappa(X \longmapsto f)(\kappa) = X \longmapsto \text{dom } f$ and $\text{rng}_\kappa(X \longmapsto f)(\kappa) = X \longmapsto \text{rng } f$.
- (37) If $f \neq \emptyset$, then $x \in \cap f$ iff for every y such that $y \in \text{dom } f$ holds $x \in f(y)$.
- (38) $\cap \emptyset = \emptyset$.
- (39) $\cup \langle X \rangle = X$ and $\cap \langle X \rangle = X$.
- (40) $\cup \langle X, Y \rangle = X \cup Y$ and $\cap \langle X, Y \rangle = X \cap Y$.
- (41) $\cup \langle X, Y, Z \rangle = X \cup Y \cup Z$ and $\cap \langle X, Y, Z \rangle = X \cap Y \cap Z$.
- (42) $\cup(\emptyset \longmapsto Y) = \emptyset$ and $\cap(\emptyset \longmapsto Y) = \emptyset$.
- (43) If $X \neq \emptyset$, then $\cup(X \longmapsto Y) = Y$ and $\cap(X \longmapsto Y) = Y$.

Let f be a function and let x, y be sets. The functor $f(x)(y)$ yields a set and is defined by:

(Def. 5) $f(x)(y) = (\text{uncurry } f)(\langle x, y \rangle)$.

Next we state several propositions:

- (44) If $x \in \text{dom } f$ and $g = f(x)$ and $y \in \text{dom } g$, then $f(x)(y) = g(y)$.
- (45) If $x \in \text{dom } f$, then $\langle f \rangle(1)(x) = f(x)$ and $\langle f, g \rangle(1)(x) = f(x)$ and $\langle f, g, h \rangle(1)(x) = f(x)$.
- (46) If $x \in \text{dom } g$, then $\langle f, g \rangle(2)(x) = g(x)$ and $\langle f, g, h \rangle(2)(x) = g(x)$.
- (47) If $x \in \text{dom } h$, then $\langle f, g, h \rangle(3)(x) = h(x)$.
- (48) If $x \in X$ and $y \in \text{dom } f$, then $(X \longmapsto f)(x)(y) = f(y)$.

4. CARTESIAN PRODUCT OF FUNCTIONS WITH THE SAME DOMAIN

Let f be a function. The functor $\Pi^* f$ yields a function and is defined by:

(Def. 6) $\Pi^* f = \text{curry}(\text{uncurry}' f | [: \cap(\text{dom}_\kappa f(\kappa)), \text{dom } f :])$.

One can prove the following propositions:

- (49) $\text{dom } \Pi^* f = \cap(\text{dom}_\kappa f(\kappa))$ and $\text{rng } \Pi^* f \subseteq \Pi(\text{rng}_\kappa f(\kappa))$.
- (50) If $x \in \text{dom } \Pi^* f$, then $(\Pi^* f)(x)$ is a function.
- (51) If $x \in \text{dom } \Pi^* f$ and $g = (\Pi^* f)(x)$, then $\text{dom } g = f^{-1}(\text{Sub}_{\text{frng}} f)$ and for every y such that $y \in \text{dom } g$ holds $\langle y, x \rangle \in \text{dom } \text{uncurry } f$ and $g(y) = (\text{uncurry } f)(\langle y, x \rangle)$.

- (52) If $x \in \text{dom } \Pi^* f$, then for every g such that $g \in \text{rng } f$ holds $x \in \text{dom } g$.
- (53) If $g \in \text{rng } f$ and for every g such that $g \in \text{rng } f$ holds $x \in \text{dom } g$, then $x \in \text{dom } \Pi^* f$.
- (54) If $x \in \text{dom } f$ and $g = f(x)$ and $y \in \text{dom } \Pi^* f$ and $h = (\Pi^* f)(y)$, then $g(y) = h(x)$.
- (55) If $x \in \text{dom } f$ and $f(x)$ is a function and $y \in \text{dom } \Pi^* f$, then $f(x)(y) = (\Pi^* f)(y)(x)$.

Let f be a function. The functor $\text{Frege}(f)$ yielding a function is defined by the conditions (Def. 7).

- (Def. 7)(i) $\text{dom } \text{Frege}(f) = \Pi(\text{dom}_K f(K))$, and
- (ii) for every g such that $g \in \Pi(\text{dom}_K f(K))$ there exists h such that $(\text{Frege}(f))(g) = h$ and $\text{dom } h = f^{-1}(\text{Sub}_f \text{rng } f)$ and for every x such that $x \in \text{dom } h$ holds $h(x) = (\text{uncurry } f)(\langle x, g(x) \rangle)$.

One can prove the following propositions:

- (56) If $g \in \Pi(\text{dom}_K f(K))$ and $x \in \text{dom } g$, then $(\text{Frege}(f))(g)(x) = f(x)(g(x))$.
- (57) If $x \in \text{dom } f$ and $g = f(x)$ and $h \in \Pi(\text{dom}_K f(K))$ and $h' = (\text{Frege}(f))(h)$, then $h(x) \in \text{dom } g$ and $h'(x) = g(h(x))$ and $h' \in \Pi(\text{rng}_K f(K))$.
- (58) $\text{rng } \text{Frege}(f) = \Pi(\text{rng}_K f(K))$.
- (59) If $\emptyset \notin \text{rng } f$, then $\text{Frege}(f)$ is one-to-one iff for every g such that $g \in \text{rng } f$ holds g is one-to-one.

5. CARTESIAN PRODUCT OF FUNCTIONS

One can prove the following propositions:

- (60) $\Pi^* \emptyset = \emptyset$ and $\text{Frege}(\emptyset) = \{\emptyset\} \longmapsto \emptyset$.
- (61) $\text{dom } \Pi^* \langle h \rangle = \text{dom } h$ and for every x such that $x \in \text{dom } h$ holds $(\Pi^* \langle h \rangle)(x) = \langle h(x) \rangle$.
- (62) $\text{dom } \Pi^* \langle f_1, f_2 \rangle = \text{dom } f_1 \cap \text{dom } f_2$ and for every x such that $x \in \text{dom } f_1 \cap \text{dom } f_2$ holds $(\Pi^* \langle f_1, f_2 \rangle)(x) = \langle f_1(x), f_2(x) \rangle$.
- (63) If $X \neq \emptyset$, then $\text{dom } \Pi^*(X \longmapsto f) = \text{dom } f$ and for every x such that $x \in \text{dom } f$ holds $(\Pi^*(X \longmapsto f))(x) = X \longmapsto f(x)$.
- (64) $\text{dom } \text{Frege}(\langle h \rangle) = \Pi \langle \text{dom } h \rangle$ and $\text{rng } \text{Frege}(\langle h \rangle) = \Pi \langle \text{rng } h \rangle$ and for every x such that $x \in \text{dom } h$ holds $(\text{Frege}(\langle h \rangle))(\langle x \rangle) = \langle h(x) \rangle$.
- (65) $\text{dom } \text{Frege}(\langle f_1, f_2 \rangle) = \Pi \langle \text{dom } f_1, \text{dom } f_2 \rangle$ and $\text{rng } \text{Frege}(\langle f_1, f_2 \rangle) = \Pi \langle \text{rng } f_1, \text{rng } f_2 \rangle$ and for all x, y such that $x \in \text{dom } f_1$ and $y \in \text{dom } f_2$ holds $(\text{Frege}(\langle f_1, f_2 \rangle))(\langle x, y \rangle) = \langle f_1(x), f_2(y) \rangle$.
- (66) $\text{dom } \text{Frege}(X \longmapsto f) = (\text{dom } f)^X$ and $\text{rng } \text{Frege}(X \longmapsto f) = (\text{rng } f)^X$ and for every g such that $g \in (\text{dom } f)^X$ holds $(\text{Frege}(X \longmapsto f))(g) = f \cdot g$.
- (67) If $x \in \text{dom } f_1$ and $x \in \text{dom } f_2$, then for all y_1, y_2 holds $\langle f_1, f_2 \rangle(x) = \langle y_1, y_2 \rangle$ iff $(\Pi^* \langle f_1, f_2 \rangle)(x) = \langle y_1, y_2 \rangle$.
- (68) If $x \in \text{dom } f_1$ and $y \in \text{dom } f_2$, then for all y_1, y_2 holds $[f_1, f_2](\langle x, y \rangle) = \langle y_1, y_2 \rangle$ iff $(\text{Frege}(\langle f_1, f_2 \rangle))(\langle x, y \rangle) = \langle y_1, y_2 \rangle$.
- (69) If $\text{dom } f = X$ and $\text{dom } g = X$ and for every x such that $x \in X$ holds $f(x) \approx g(x)$, then $\Pi f \approx \Pi g$.
- (70) If $\text{dom } f = \text{dom } h$ and $\text{dom } g = \text{rng } h$ and h is one-to-one and for every x such that $x \in \text{dom } h$ holds $f(x) \approx g(h(x))$, then $\Pi f \approx \Pi g$.
- (71) If $\text{dom } f = X$, then $\Pi f \approx \Pi(f \cdot P)$.

6. FUNCTION YIELDING POWERS

Let us consider f, X . The functor X^f yielding a function is defined as follows:

(Def. 8) $\text{dom}(X^f) = \text{dom } f$ and for every x such that $x \in \text{dom } f$ holds $X^f(x) = X^{f(x)}$.

We now state several propositions:

(72) If $\emptyset \notin \text{rng } f$, then $\emptyset^f = \text{dom } f \longmapsto \emptyset$.

(73) $X^\emptyset = \emptyset$.

(74) $Y^{\langle X \rangle} = \langle Y^X \rangle$.

(75) $Z^{\langle X, Y \rangle} = \langle Z^X, Z^Y \rangle$.

(76) $Z^{X \longmapsto Y} = X \longmapsto Z^Y$.

(77) $X^{\cup \text{disjoint}^f} \approx \prod(X^f)$.

Let us consider X, f . The functor f^X yielding a function is defined by:

(Def. 9) $\text{dom}(f^X) = \text{dom } f$ and for every x such that $x \in \text{dom } f$ holds $f^X(x) = f(x)^X$.

The following propositions are true:

(78) $f^\emptyset = \text{dom } f \longmapsto \{\emptyset\}$.

(79) $\emptyset^X = \emptyset$.

(80) $\langle Y \rangle^X = \langle Y^X \rangle$.

(81) $\langle Y, Z \rangle^X = \langle Y^X, Z^X \rangle$.

(82) $(Y \longmapsto Z)^X = Y \longmapsto Z^X$.

(83) $\prod(f^X) \approx (\prod f)^X$.

REFERENCES

- [1] Grzegorz Bancerek. Zermelo theorem and axiom of choice. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/wellord2.html>.
- [2] Grzegorz Bancerek. Curried and uncurried functions. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/funct_5.html.
- [3] Grzegorz Bancerek. König's theorem. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/card_3.html.
- [4] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.
- [5] Czesław Byliński. Basic functions and operations on functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_3.html.
- [6] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [7] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [8] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/partfun1.html>.
- [9] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [10] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_2.html.
- [11] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/funct_4.html.
- [12] Andrzej Nędzusiak. σ -fields and probability. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/prob_1.html.

- [13] Beata Padlewska. Families of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/setfam_1.html.
- [14] Andrzej Trybulec. Binary operations applied to functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funcop_1.html.
- [15] Andrzej Trybulec. Enumerated sets. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/enumset1.html>.
- [16] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [17] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [18] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received September 30, 1991

Published January 2, 2004
