

Category of Functors between Alternative Categories

Robert Nieszczerzewski
Warsaw University
Białystok

MML Identifier: FUNCTOR2.

WWW: <http://mizar.org/JFM/Vol9/functor2.html>

The articles [7], [5], [13], [12], [14], [1], [3], [4], [2], [6], [8], [9], [10], and [11] provide the notation and terminology for this paper.

1. PRELIMINARIES

Let A be a transitive non empty category structure with units and let B be a non empty category structure with units. Note that every functor from A to B is feasible and id-preserving.

Let A be a transitive non empty category structure with units and let B be a non empty category structure with units. One can verify the following observations:

- * every functor from A to B which is covariant is also precovariant and comp-preserving,
- * every functor from A to B which is precovariant and comp-preserving is also covariant,
- * every functor from A to B which is contravariant is also precontravariant and comp-reversing, and
- * every functor from A to B which is precontravariant and comp-reversing is also contravariant.

Next we state the proposition

- (2)¹ Let A, B be transitive non empty category structures with units, F be a covariant functor from A to B , and a be an object of A . Then $F(\text{id}_a) = \text{id}_{F(a)}$.

2. TRANSFORMATIONS

Let A, B be transitive non empty category structures with units and let F_1, F_2 be covariant functors from A to B . We say that F_1 is transformable to F_2 if and only if:

(Def. 1) For every object a of A holds $\langle F_1(a), F_2(a) \rangle \neq \emptyset$.

Let us note that the predicate F_1 is transformable to F_2 is reflexive.

The following proposition is true

¹ The proposition (1) has been removed.

(4)² Let A, B be transitive non empty category structures with units and F, F_1, F_2 be covariant functors from A to B . Suppose F is transformable to F_1 and F_1 is transformable to F_2 . Then F is transformable to F_2 .

Let A, B be transitive non empty category structures with units and let F_1, F_2 be covariant functors from A to B . Let us assume that F_1 is transformable to F_2 . A many sorted set indexed by the carrier of A is said to be a transformation from F_1 to F_2 if:

(Def. 2) For every object a of A holds $it(a)$ is a morphism from $F_1(a)$ to $F_2(a)$.

Let A, B be transitive non empty category structures with units and let F be a covariant functor from A to B . The functor id_F yields a transformation from F to F and is defined by:

(Def. 3) For every object a of A holds $id_F(a) = id_{F(a)}$.

Let A, B be transitive non empty category structures with units and let F_1, F_2 be covariant functors from A to B . Let us assume that F_1 is transformable to F_2 . Let t be a transformation from F_1 to F_2 and let a be an object of A . The functor $t[a]$ yielding a morphism from $F_1(a)$ to $F_2(a)$ is defined as follows:

(Def. 4) $t[a] = t(a)$.

Let A, B be transitive non empty category structures with units and let F, F_1, F_2 be covariant functors from A to B . Let us assume that F is transformable to F_1 and F_1 is transformable to F_2 . Let t_1 be a transformation from F to F_1 and let t_2 be a transformation from F_1 to F_2 . The functor $t_2 \circ t_1$ yields a transformation from F to F_2 and is defined by:

(Def. 5) For every object a of A holds $(t_2 \circ t_1)[a] = t_2[a] \cdot t_1[a]$.

The following four propositions are true:

(5) Let A, B be transitive non empty category structures with units and F_1, F_2 be covariant functors from A to B . Suppose F_1 is transformable to F_2 . Let t_1, t_2 be transformations from F_1 to F_2 . If for every object a of A holds $t_1[a] = t_2[a]$, then $t_1 = t_2$.

(6) Let A, B be transitive non empty category structures with units, F be a covariant functor from A to B , and a be an object of A . Then $id_F[a] = id_{F(a)}$.

(7) Let A, B be transitive non empty category structures with units and F_1, F_2 be covariant functors from A to B . Suppose F_1 is transformable to F_2 . Let t be a transformation from F_1 to F_2 . Then $id_{(F_2)} \circ t = t$ and $t \circ id_{(F_1)} = t$.

(8) Let A, B be categories and F, F_1, F_2, F_3 be covariant functors from A to B . Suppose F is transformable to F_1 and F_1 is transformable to F_2 and F_2 is transformable to F_3 . Let t_1 be a transformation from F to F_1 , t_2 be a transformation from F_1 to F_2 , and t_3 be a transformation from F_2 to F_3 . Then $(t_3 \circ t_2) \circ t_1 = t_3 \circ (t_2 \circ t_1)$.

3. NATURAL TRANSFORMATIONS

Let A, B be transitive non empty category structures with units and let F_1, F_2 be covariant functors from A to B . We say that F_1 is naturally transformable to F_2 if and only if the conditions (Def. 6) are satisfied.

(Def. 6)(i) F_1 is transformable to F_2 , and

(ii) there exists a transformation t from F_1 to F_2 such that for all objects a, b of A such that $\langle a, b \rangle \neq \emptyset$ and for every morphism f from a to b holds $t[b] \cdot F_1(f) = F_2(f) \cdot t[a]$.

Next we state two propositions:

² The proposition (3) has been removed.

(9) For all transitive non empty category structures A, B with units holds every covariant functor F from A to B is naturally transformable to F .

(10) Let A, B be categories and F, F_1, F_2 be covariant functors from A to B . Suppose F is naturally transformable to F_1 and F_1 is naturally transformable to F_2 . Then F is naturally transformable to F_2 .

Let A, B be transitive non empty category structures with units and let F_1, F_2 be covariant functors from A to B . Let us assume that F_1 is naturally transformable to F_2 . A transformation from F_1 to F_2 is said to be a natural transformation from F_1 to F_2 if:

(Def. 7) For all objects a, b of A such that $\langle a, b \rangle \neq \emptyset$ and for every morphism f from a to b holds $it[b] \cdot F_1(f) = F_2(f) \cdot it[a]$.

Let A, B be transitive non empty category structures with units and let F be a covariant functor from A to B . Then id_F is a natural transformation from F to F .

Let A, B be categories and let F, F_1, F_2 be covariant functors from A to B . Let us assume that F is naturally transformable to F_1 and F_1 is naturally transformable to F_2 . Let t_1 be a natural transformation from F to F_1 and let t_2 be a natural transformation from F_1 to F_2 . The functor $t_2 \circ t_1$ yields a natural transformation from F to F_2 and is defined as follows:

(Def. 8) $t_2 \circ t_1 = t_2 \circ t_1$.

Next we state three propositions:

(11) Let A, B be transitive non empty category structures with units and F_1, F_2 be covariant functors from A to B . Suppose F_1 is naturally transformable to F_2 . Let t be a natural transformation from F_1 to F_2 . Then $id_{(F_2)} \circ t = t$ and $t \circ id_{(F_1)} = t$.

(12) Let A, B be transitive non empty category structures with units and F, F_1, F_2 be covariant functors from A to B . Suppose F is naturally transformable to F_1 and F_1 is naturally transformable to F_2 . Let t_1 be a natural transformation from F to F_1 , t_2 be a natural transformation from F_1 to F_2 , and a be an object of A . Then $(t_2 \circ t_1)[a] = t_2[a] \cdot t_1[a]$.

(13) Let A, B be categories, F, F_1, F_2, F_3 be covariant functors from A to B , t be a natural transformation from F to F_1 , and t_1 be a natural transformation from F_1 to F_2 . Suppose F is naturally transformable to F_1 and F_1 is naturally transformable to F_2 and F_2 is naturally transformable to F_3 . Let t_3 be a natural transformation from F_2 to F_3 . Then $(t_3 \circ t_1) \circ t = t_3 \circ (t_1 \circ t)$.

4. CATEGORY OF FUNCTORS

Let I be a set and let A, B be many sorted sets indexed by I . The functor B^A yielding a set is defined as follows:

(Def. 9)(i) For every set x holds $x \in B^A$ iff x is a many sorted function from A into B if for every set i such that $i \in I$ holds if $B(i) = \emptyset$, then $A(i) = \emptyset$,

(ii) $B^A = \emptyset$, otherwise.

Let A, B be transitive non empty category structures with units. The functor $\text{Funct}(A, B)$ yields a set and is defined as follows:

(Def. 10) For every set x holds $x \in \text{Funct}(A, B)$ iff x is a covariant strict functor from A to B .

Let A, B be categories. The functor B^A yielding a strict non empty transitive category structure is defined by the conditions (Def. 11).

- (Def. 11)(i) The carrier of $B^A = \text{Funct}(A, B)$,
- (ii) for all strict covariant functors F, G from A to B and for every set x holds $x \in$ (the arrows of B^A)(F, G) iff F is naturally transformable to G and x is a natural transformation from F to G , and
- (iii) for all strict covariant functors F, G, H from A to B such that F is naturally transformable to G and G is naturally transformable to H and for every natural transformation t_1 from F to G and for every natural transformation t_2 from G to H there exists a function f such that $f =$ (the composition of B^A)(F, G, H) and $f(t_2, t_1) = t_2 \circ t_1$.

REFERENCES

- [1] Grzegorz Bancerek. König's theorem. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/card_3.html.
- [2] Czesław Byliński. Binary operations. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/binop_1.html.
- [3] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [4] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [5] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [6] Michał Muzalewski and Wojciech Skaba. Three-argument operations and four-argument operations. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/multop_1.html.
- [7] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [8] Andrzej Trybulec. Function domains and Fränkel operator. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol2/fraenkel.html>.
- [9] Andrzej Trybulec. Many-sorted sets. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/pboole.html>.
- [10] Andrzej Trybulec. Many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg_1.html.
- [11] Andrzej Trybulec. Categories without uniqueness of cod and dom. *Journal of Formalized Mathematics*, 7, 1995. http://mizar.org/JFM/Vol7/altcat_1.html.
- [12] Andrzej Trybulec. Functors for alternative categories. *Journal of Formalized Mathematics*, 8, 1996. <http://mizar.org/JFM/Vol8/functor0.html>.
- [13] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [14] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received June 12, 1997

Published January 2, 2004
