

## Some Topological Properties of Cells in $R^2$

Yatsuka Nakamura  
Shinshu University  
Nagano

Andrzej Trybulec  
Warsaw University  
Białystok

**Summary.** We examine the topological property of cells (rectangles) in a plane. First, some Fraenkel expressions of cells are shown. Second, it is proved that cells are closed. The last theorem asserts that the closure of the interior of a cell is the same as itself.

MML Identifier: GOBRD11.

WWW: <http://mizar.org/JFM/Vol8/gobrd11.html>

The articles [14], [15], [5], [8], [2], [1], [13], [3], [6], [7], [16], [12], [4], [9], [10], and [11] provide the notation and terminology for this paper.

We use the following convention:  $i, j$  denote natural numbers,  $r, s, r_2, s_1, s_2$  denote real numbers, and  $G_1$  denotes a non empty topological space.

One can prove the following propositions:

- (1) For every subset  $A$  of  $G_1$  and for every point  $p$  of  $G_1$  such that  $p \in A$  and  $A$  is connected holds  $A \subseteq \text{Component}(p)$ .
- (2) For all subsets  $A, B, C$  of  $G_1$  such that  $C$  is a component of  $G_1$  and  $A \subseteq C$  and  $B$  is connected and  $\bar{A}$  meets  $\bar{B}$  holds  $B \subseteq C$ .

In the sequel  $G_2$  is a non empty topological space.

One can prove the following three propositions:

- (3) Let  $A, B$  be subsets of  $G_2$ . Suppose  $A$  is a component of  $G_2$  and  $B$  is a component of  $G_2$ . Then  $A \cup B$  is a union of components of  $G_2$ .
- (4) For all subsets  $B_1, B_2, V$  of  $G_1$  holds  $\text{Down}(B_1 \cup B_2, V) = \text{Down}(B_1, V) \cup \text{Down}(B_2, V)$ .
- (5) For all subsets  $B_1, B_2, V$  of  $G_1$  holds  $\text{Down}(B_1 \cap B_2, V) = \text{Down}(B_1, V) \cap \text{Down}(B_2, V)$ .

In the sequel  $f$  is a non constant standard special circular sequence and  $G$  is a non empty yielding matrix over  $\mathcal{E}_T^2$ .

The following proposition is true

- (6)  $(\tilde{\mathcal{L}}(f))^c \neq \emptyset$ .

Let us consider  $f$ . Note that  $(\tilde{\mathcal{L}}(f))^c$  is non empty.

Next we state a number of propositions:

- (7) For every  $f$  holds the carrier of  $\mathcal{E}_T^2 = \bigcup \{\text{cell}(\text{the Go-board of } f, i, j) : i \leq \text{len the Go-board of } f \wedge j \leq \text{width the Go-board of } f\}$ .

- (8) For all subsets  $P_1, P_2$  of  $\mathcal{E}_T^2$  such that  $P_1 = \{[r, s] : s \leq s_1\}$  and  $P_2 = \{[r_2, s_2] : s_2 > s_1\}$  holds  $P_1 = P_2^c$ .
- (9) For all subsets  $P_1, P_2$  of  $\mathcal{E}_T^2$  such that  $P_1 = \{[r, s] : s \geq s_1\}$  and  $P_2 = \{[r_2, s_2] : s_2 < s_1\}$  holds  $P_1 = P_2^c$ .
- (10) For all subsets  $P_1, P_2$  of  $\mathcal{E}_T^2$  such that  $P_1 = \{[s, r] : s \geq s_1\}$  and  $P_2 = \{[s_2, r_2] : s_2 < s_1\}$  holds  $P_1 = P_2^c$ .
- (11) For all subsets  $P_1, P_2$  of  $\mathcal{E}_T^2$  such that  $P_1 = \{[s, r] : s \leq s_1\}$  and  $P_2 = \{[s_2, r_2] : s_2 > s_1\}$  holds  $P_1 = P_2^c$ .
- (12) For every subset  $P$  of  $\mathcal{E}_T^2$  and for every  $s_1$  such that  $P = \{[r, s] : s \leq s_1\}$  holds  $P$  is closed.
- (13) For every subset  $P$  of  $\mathcal{E}_T^2$  and for every  $s_1$  such that  $P = \{[r, s] : s_1 \leq s\}$  holds  $P$  is closed.
- (14) For every subset  $P$  of  $\mathcal{E}_T^2$  and for every  $s_1$  such that  $P = \{[s, r] : s \leq s_1\}$  holds  $P$  is closed.
- (15) For every subset  $P$  of  $\mathcal{E}_T^2$  and for every  $s_1$  such that  $P = \{[s, r] : s_1 \leq s\}$  holds  $P$  is closed.
- (16) For every matrix  $G$  over  $\mathcal{E}_T^2$  holds  $\text{hstrip}(G, j)$  is closed.
- (17) For every matrix  $G$  over  $\mathcal{E}_T^2$  holds  $\text{vstrip}(G, j)$  is closed.
- (18) If  $G$  is line  $\mathbf{X}$ -constant, then  $\text{vstrip}(G, 0) = \{[r, s] : r \leq (G \circ (1, 1))_1\}$ .
- (19) If  $G$  is line  $\mathbf{X}$ -constant, then  $\text{vstrip}(G, \text{len } G) = \{[r, s] : (G \circ (\text{len } G, 1))_1 \leq r\}$ .
- (20) If  $G$  is line  $\mathbf{X}$ -constant and  $1 \leq i$  and  $i < \text{len } G$ , then  $\text{vstrip}(G, i) = \{[r, s] : (G \circ (i, 1))_1 \leq r \wedge r \leq (G \circ (i+1, 1))_1\}$ .
- (21) If  $G$  is column  $\mathbf{Y}$ -constant, then  $\text{hstrip}(G, 0) = \{[r, s] : s \leq (G \circ (1, 1))_2\}$ .
- (22) If  $G$  is column  $\mathbf{Y}$ -constant, then  $\text{hstrip}(G, \text{width } G) = \{[r, s] : (G \circ (1, \text{width } G))_2 \leq s\}$ .
- (23) If  $G$  is column  $\mathbf{Y}$ -constant and  $1 \leq j$  and  $j < \text{width } G$ , then  $\text{hstrip}(G, j) = \{[r, s] : (G \circ (1, j))_2 \leq s \wedge s \leq (G \circ (1, j+1))_2\}$ .

In the sequel  $G$  is a non empty yielding line  $\mathbf{X}$ -constant column  $\mathbf{Y}$ -constant matrix over  $\mathcal{E}_T^2$ .

The following propositions are true:

- (24)  $\text{cell}(G, 0, 0) = \{[r, s] : r \leq (G \circ (1, 1))_1 \wedge s \leq (G \circ (1, 1))_2\}$ .
- (25)  $\text{cell}(G, 0, \text{width } G) = \{[r, s] : r \leq (G \circ (1, 1))_1 \wedge (G \circ (1, \text{width } G))_2 \leq s\}$ .
- (26) If  $1 \leq j$  and  $j < \text{width } G$ , then  $\text{cell}(G, 0, j) = \{[r, s] : r \leq (G \circ (1, 1))_1 \wedge (G \circ (1, j))_2 \leq s \wedge s \leq (G \circ (1, j+1))_2\}$ .
- (27)  $\text{cell}(G, \text{len } G, 0) = \{[r, s] : (G \circ (\text{len } G, 1))_1 \leq r \wedge s \leq (G \circ (1, 1))_2\}$ .
- (28)  $\text{cell}(G, \text{len } G, \text{width } G) = \{[r, s] : (G \circ (\text{len } G, 1))_1 \leq r \wedge (G \circ (1, \text{width } G))_2 \leq s\}$ .
- (29) If  $1 \leq j$  and  $j < \text{width } G$ , then  $\text{cell}(G, \text{len } G, j) = \{[r, s] : (G \circ (\text{len } G, 1))_1 \leq r \wedge (G \circ (1, j))_2 \leq s \wedge s \leq (G \circ (1, j+1))_2\}$ .
- (30) If  $1 \leq i$  and  $i < \text{len } G$ , then  $\text{cell}(G, i, 0) = \{[r, s] : (G \circ (i, 1))_1 \leq r \wedge r \leq (G \circ (i+1, 1))_1 \wedge s \leq (G \circ (1, 1))_2\}$ .
- (31) If  $1 \leq i$  and  $i < \text{len } G$ , then  $\text{cell}(G, i, \text{width } G) = \{[r, s] : (G \circ (i, 1))_1 \leq r \wedge r \leq (G \circ (i+1, 1))_1 \wedge (G \circ (1, \text{width } G))_2 \leq s\}$ .
- (32) If  $1 \leq i$  and  $i < \text{len } G$  and  $1 \leq j$  and  $j < \text{width } G$ , then  $\text{cell}(G, i, j) = \{[r, s] : (G \circ (i, 1))_1 \leq r \wedge r \leq (G \circ (i+1, 1))_1 \wedge (G \circ (1, j))_2 \leq s \wedge s \leq (G \circ (1, j+1))_2\}$ .
- (33) For every matrix  $G$  over  $\mathcal{E}_T^2$  holds  $\text{cell}(G, i, j)$  is closed.

- (34) For every non empty yielding matrix  $G$  over  $\mathcal{E}_T^2$  holds  $1 \leq \text{len } G$  and  $1 \leq \text{width } G$ .
- (35) For every Go-board  $G$  such that  $i \leq \text{len } G$  and  $j \leq \text{width } G$  holds  $\text{cell}(G, i, j) = \overline{\text{Int cell}(G, i, j)}$ .

## REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/nat\\_1.html](http://mizar.org/JFM/Vol1/nat_1.html).
- [2] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinal1.html>.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/finseq\\_1.html](http://mizar.org/JFM/Vol1/finseq_1.html).
- [4] Agata Darmochwał. The Euclidean space. *Journal of Formalized Mathematics*, 3, 1991. <http://mizar.org/JFM/Vol3/euclid.html>.
- [5] Agata Darmochwał and Yatsuka Nakamura. The topological space  $\mathcal{E}_T^2$ . Arcs, line segments and special polygonal arcs. *Journal of Formalized Mathematics*, 3, 1991. <http://mizar.org/JFM/Vol3/topreal1.html>.
- [6] Katarzyna Jankowska. Matrices. Abelian group of matrices. *Journal of Formalized Mathematics*, 3, 1991. [http://mizar.org/JFM/Vol3/matrix\\_1.html](http://mizar.org/JFM/Vol3/matrix_1.html).
- [7] Stanisława Kanas, Adam Lecko, and Mariusz Startek. Metric spaces. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/metric\\_1.html](http://mizar.org/JFM/Vol2/metric_1.html).
- [8] Jarosław Kotowicz and Yatsuka Nakamura. Introduction to Go-Board — part I. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/goboard1.html>.
- [9] Jarosław Kotowicz and Yatsuka Nakamura. Introduction to Go-Board — part II. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/goboard2.html>.
- [10] Yatsuka Nakamura and Andrzej Trybulec. Decomposing a Go-Board into cells. *Journal of Formalized Mathematics*, 7, 1995. <http://mizar.org/JFM/Vol7/goboard5.html>.
- [11] Yatsuka Nakamura and Andrzej Trybulec. Components and unions of components. *Journal of Formalized Mathematics*, 8, 1996. [http://mizar.org/JFM/Vol8/connsp\\_3.html](http://mizar.org/JFM/Vol8/connsp_3.html).
- [12] Beata Padlewska. Connected spaces. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/connsp\\_1.html](http://mizar.org/JFM/Vol1/connsp_1.html).
- [13] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/pre\\_topc.html](http://mizar.org/JFM/Vol1/pre_topc.html).
- [14] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [15] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/subset\\_1.html](http://mizar.org/JFM/Vol1/subset_1.html).
- [16] Mirosław Wysocki and Agata Darmochwał. Subsets of topological spaces. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/tops\\_1.html](http://mizar.org/JFM/Vol1/tops_1.html).

Received July 22, 1996

Published January 2, 2004

---