

Euler Circuits and Paths¹

Yatsuka Nakamura
Shinshu University
Nagano

Piotr Rudnicki
University of Alberta
Edmonton

Summary. We prove the Euler theorem on existence of Euler circuits and paths in multigraphs.

MML Identifier: GRAPH_3.

WWW: http://mizar.org/JFM/Vol9/graph_3.html

The articles [16], [7], [20], [17], [21], [4], [6], [5], [8], [10], [2], [18], [12], [3], [9], [19], [1], [13], [14], [11], and [15] provide the notation and terminology for this paper.

1. PRELIMINARIES

Let D be a set, let T be a non empty set of finite sequences of D , and let S be a non empty subset of T . We see that the element of S is a finite sequence of elements of D .

Let i, j be even integers. One can verify that $i - j$ is even.

One can prove the following propositions:

- (1) For all integers i, j holds i is even iff j is even iff $i - j$ is even.
- (2) Let p be a finite sequence and m, n, a be natural numbers. Suppose $a \in \text{dom}\langle p(m), \dots, p(n) \rangle$. Then there exists a natural number k such that $k \in \text{dom } p$ and $p(k) = \langle p(m), \dots, p(n) \rangle(a)$ and $k + 1 = m + a$ and $m \leq k$ and $k \leq n$.

Let G be a graph. A vertex of G is an element of the vertices of G .

For simplicity, we adopt the following rules: G is a graph, v, v_1, v_2 are vertices of G , c is a chain of G , p, p_1, p_2 are paths of G , v_3, v_4, v_5 are finite sequences of elements of the vertices of G , e, X are sets, and n, m are natural numbers.

The following propositions are true:

- (3) If v_3 is vertex sequence of c , then v_3 is non empty.
- (7)¹ If $e \in$ the edges of G , then $\langle e \rangle$ is a path of G .
- (8) $\langle p(m), \dots, p(n) \rangle$ is a path of G .
- (9) Suppose $\text{rng } p_1$ misses $\text{rng } p_2$ and v_4 is vertex sequence of p_1 and v_5 is vertex sequence of p_2 and $v_4(\text{len } v_4) = v_5(1)$. Then $p_1 \hat{\ } p_2$ is a path of G .
- (12)² If $c = \emptyset$, then c is cyclic.

¹This work was partially supported by NSERC Grant OGP9207 and Shinshu Endowment Fund for Information Science.

¹ The propositions (4)–(6) have been removed.

² The propositions (10) and (11) have been removed.

Let G be a graph. Observe that there exists a path of G which is cyclic.

Next we state several propositions:

- (13) For every cyclic path p of G holds $\langle p(m+1), \dots, p(\text{len } p) \rangle \frown \langle p(1), \dots, p(m) \rangle$ is a cyclic path of G .
- (14) If $m+1 \in \text{dom } p$, then $\text{len}(\langle p(m+1), \dots, p(\text{len } p) \rangle \frown \langle p(1), \dots, p(m) \rangle) = \text{len } p$ and $\text{rng}(\langle p(m+1), \dots, p(\text{len } p) \rangle \frown \langle p(1), \dots, p(m) \rangle) = \text{rng } p$ and $(\langle p(m+1), \dots, p(\text{len } p) \rangle \frown \langle p(1), \dots, p(m) \rangle)(1) = p(m+1)$.
- (15) For every cyclic path p of G such that $n \in \text{dom } p$ there exists a cyclic path p' of G such that $p'(1) = p(n)$ and $\text{len } p' = \text{len } p$ and $\text{rng } p' = \text{rng } p$.
- (16) Let s, t be vertices of G . Suppose $s = (\text{the source of } G)(e)$ and $t = (\text{the target of } G)(e)$. Then $\langle t, s \rangle$ is vertex sequence of $\langle e \rangle$.
- (17) Suppose $e \in$ the edges of G and v_3 is vertex sequence of c and $v_3(\text{len } v_3) = (\text{the source of } G)(e)$. Then
- (i) $c \frown \langle e \rangle$ is a chain of G , and
 - (ii) there exists a finite sequence v'_1 of elements of the vertices of G such that $v'_1 = v_3 \frown \langle (\text{the source of } G)(e), (\text{the target of } G)(e) \rangle$ and v'_1 is vertex sequence of $c \frown \langle e \rangle$ and $v'_1(1) = v_3(1)$ and $v'_1(\text{len } v'_1) = (\text{the target of } G)(e)$.
- (18) Suppose $e \in$ the edges of G and v_3 is vertex sequence of c and $v_3(\text{len } v_3) = (\text{the target of } G)(e)$. Then
- (i) $c \frown \langle e \rangle$ is a chain of G , and
 - (ii) there exists a finite sequence v'_1 of elements of the vertices of G such that $v'_1 = v_3 \frown \langle (\text{the target of } G)(e), (\text{the source of } G)(e) \rangle$ and v'_1 is vertex sequence of $c \frown \langle e \rangle$ and $v'_1(1) = v_3(1)$ and $v'_1(\text{len } v'_1) = (\text{the source of } G)(e)$.
- (19) Suppose v_3 is vertex sequence of c . Let n be a natural number. Suppose $n \in \text{dom } c$. Then
- (i) $v_3(n) = (\text{the target of } G)(c(n))$ and $v_3(n+1) = (\text{the source of } G)(c(n))$, or
 - (ii) $v_3(n) = (\text{the source of } G)(c(n))$ and $v_3(n+1) = (\text{the target of } G)(c(n))$.
- (20) If v_3 is vertex sequence of c and $e \in \text{rng } c$, then $(\text{the target of } G)(e) \in \text{rng } v_3$ and $(\text{the source of } G)(e) \in \text{rng } v_3$.

Let G be a graph and let X be a set. Then $G\text{-VSet}(X)$ is a subset of the vertices of G .

We now state four propositions:

- (21) $G\text{-VSet}(\emptyset) = \emptyset$.
- (22) If $e \in$ the edges of G and $e \in X$, then $G\text{-VSet}(X)$ is non empty.
- (23) G is connected if and only if for all v_1, v_2 such that $v_1 \neq v_2$ there exist c, v_3 such that c is non empty and v_3 is vertex sequence of c and $v_3(1) = v_1$ and $v_3(\text{len } v_3) = v_2$.
- (24) Let G be a connected graph, X be a set, and v be a vertex of G . Suppose X meets the edges of G and $v \notin G\text{-VSet}(X)$. Then there exists a vertex v' of G and there exists an element e of the edges of G such that $v' \in G\text{-VSet}(X)$ but $e \notin X$ but $v' = (\text{the target of } G)(e)$ or $v' = (\text{the source of } G)(e)$.

2. DEGREE OF A VERTEX

Let G be a graph, let v be a vertex of G , and let X be a set. The functor $\text{EdgesIn}(v, X)$ yields a subset of the edges of G and is defined as follows:

(Def. 1) For every set e holds $e \in \text{EdgesIn}(v, X)$ iff $e \in$ the edges of G and $e \in X$ and (the target of G)(e) = v .

The functor $\text{EdgesOut}(v, X)$ yields a subset of the edges of G and is defined as follows:

(Def. 2) For every set e holds $e \in \text{EdgesOut}(v, X)$ iff $e \in$ the edges of G and $e \in X$ and (the source of G)(e) = v .

Let G be a graph, let v be a vertex of G , and let X be a set. The functor $\text{EdgesAt}(v, X)$ yields a subset of the edges of G and is defined as follows:

(Def. 3) $\text{EdgesAt}(v, X) = \text{EdgesIn}(v, X) \cup \text{EdgesOut}(v, X)$.

Let G be a finite graph, let v be a vertex of G , and let X be a set. One can check the following observations:

- * $\text{EdgesIn}(v, X)$ is finite,
- * $\text{EdgesOut}(v, X)$ is finite, and
- * $\text{EdgesAt}(v, X)$ is finite.

Let G be a graph, let v be a vertex of G , and let X be an empty set. One can check the following observations:

- * $\text{EdgesIn}(v, X)$ is empty,
- * $\text{EdgesOut}(v, X)$ is empty, and
- * $\text{EdgesAt}(v, X)$ is empty.

Let G be a graph and let v be a vertex of G . The functor $\text{EdgesIn } v$ yields a subset of the edges of G and is defined as follows:

(Def. 4) $\text{EdgesIn } v = \text{EdgesIn}(v, \text{the edges of } G)$.

The functor $\text{EdgesOut } v$ yields a subset of the edges of G and is defined by:

(Def. 5) $\text{EdgesOut } v = \text{EdgesOut}(v, \text{the edges of } G)$.

We now state two propositions:

$$(25) \quad \text{EdgesIn}(v, X) \subseteq \text{EdgesIn } v.$$

$$(26) \quad \text{EdgesOut}(v, X) \subseteq \text{EdgesOut } v.$$

Let G be a finite graph and let v be a vertex of G . Observe that $\text{EdgesIn } v$ is finite and $\text{EdgesOut } v$ is finite.

For simplicity, we adopt the following convention: G is a finite graph, v is a vertex of G , c is a chain of G , v_3 is a finite sequence of elements of the vertices of G , and X_1, X_2 are sets.

One can prove the following propositions:

$$(27) \quad \text{cardEdgesIn } v = \text{EdgIn}(v).$$

$$(28) \quad \text{cardEdgesOut } v = \text{EdgOut}(v).$$

Let G be a finite graph, let v be a vertex of G , and let X be a set. The functor $\text{Degree}(v, X)$ yields a natural number and is defined as follows:

(Def. 6) $\text{Degree}(v, X) = \text{cardEdgesIn}(v, X) + \text{cardEdgesOut}(v, X)$.

The following propositions are true:

- (29) The degree of $v = \text{Degree}(v, \text{the edges of } G)$.
- (30) If $\text{Degree}(v, X) \neq 0$, then $\text{EdgesAt}(v, X)$ is non empty.
- (31) Suppose $e \in \text{the edges of } G$ but $e \notin X$ but $v = (\text{the target of } G)(e)$ or $v = (\text{the source of } G)(e)$. Then the degree of $v \neq \text{Degree}(v, X)$.
- (32) If $X_2 \subseteq X_1$, then $\text{cardEdgesIn}(v, X_1 \setminus X_2) = \text{cardEdgesIn}(v, X_1) - \text{cardEdgesIn}(v, X_2)$.
- (33) If $X_2 \subseteq X_1$, then $\text{cardEdgesOut}(v, X_1 \setminus X_2) = \text{cardEdgesOut}(v, X_1) - \text{cardEdgesOut}(v, X_2)$.
- (34) If $X_2 \subseteq X_1$, then $\text{Degree}(v, X_1 \setminus X_2) = \text{Degree}(v, X_1) - \text{Degree}(v, X_2)$.
- (35) $\text{EdgesIn}(v, X) = \text{EdgesIn}(v, X \cap \text{the edges of } G)$ and $\text{EdgesOut}(v, X) = \text{EdgesOut}(v, X \cap \text{the edges of } G)$.
- (36) $\text{Degree}(v, X) = \text{Degree}(v, X \cap \text{the edges of } G)$.
- (37) If c is non empty and v_3 is vertex sequence of c , then $v \in \text{rng } v_3$ iff $\text{Degree}(v, \text{rng } c) \neq 0$.
- (38) For every non empty finite connected graph G and for every vertex v of G holds the degree of $v \neq 0$.

3. ADDING AN EDGE TO A GRAPH

Let G be a graph and let v_1, v_2 be vertices of G . The functor $\text{AddNewEdge}(v_1, v_2)$ yields a strict graph and is defined by the conditions (Def. 7).

- (Def. 7)(i) The vertices of $\text{AddNewEdge}(v_1, v_2) = \text{the vertices of } G$,
- (ii) the edges of $\text{AddNewEdge}(v_1, v_2) = (\text{the edges of } G) \cup \{\text{the edges of } G\}$,
- (iii) the source of $\text{AddNewEdge}(v_1, v_2) = (\text{the source of } G) + \cdot ((\text{the edges of } G) \mapsto (v_1))$, and
- (iv) the target of $\text{AddNewEdge}(v_1, v_2) = (\text{the target of } G) + \cdot ((\text{the edges of } G) \mapsto (v_2))$.

Let G be a finite graph and let v_1, v_2 be vertices of G . Observe that $\text{AddNewEdge}(v_1, v_2)$ is finite.

For simplicity, we adopt the following convention: G denotes a graph, v, v_1, v_2 denote vertices of G , c denotes a chain of G , p denotes a path of G , v_3 denotes a finite sequence of elements of the vertices of G , v' denotes a vertex of $\text{AddNewEdge}(v_1, v_2)$, p' denotes a path of $\text{AddNewEdge}(v_1, v_2)$, and v'_1 denotes a finite sequence of elements of the vertices of $\text{AddNewEdge}(v_1, v_2)$.

The following propositions are true:

- (39)(i) The edges of $G \in \text{the edges of } \text{AddNewEdge}(v_1, v_2)$,
- (ii) the edges of $G = (\text{the edges of } \text{AddNewEdge}(v_1, v_2)) \setminus \{\text{the edges of } G\}$,
- (iii) $(\text{the source of } \text{AddNewEdge}(v_1, v_2))(\text{the edges of } G) = v_1$, and
- (iv) $(\text{the target of } \text{AddNewEdge}(v_1, v_2))(\text{the edges of } G) = v_2$.
- (40) Suppose $e \in \text{the edges of } G$. Then $(\text{the source of } \text{AddNewEdge}(v_1, v_2))(e) = (\text{the source of } G)(e)$ and $(\text{the target of } \text{AddNewEdge}(v_1, v_2))(e) = (\text{the target of } G)(e)$.
- (41) If $v'_1 = v_3$ and v_3 is vertex sequence of c , then v'_1 is vertex sequence of c .
- (42) c is a chain of $\text{AddNewEdge}(v_1, v_2)$.
- (43) p is a path of $\text{AddNewEdge}(v_1, v_2)$.
- (44) If $v' = v_1$ and $v_1 \neq v_2$, then $\text{EdgesIn}(v', X) = \text{EdgesIn}(v_1, X)$.
- (45) If $v' = v_2$ and $v_1 \neq v_2$, then $\text{EdgesOut}(v', X) = \text{EdgesOut}(v_2, X)$.

- (46) If $v' = v_1$ and $v_1 \neq v_2$ and the edges of $G \in X$, then $\text{EdgesOut}(v', X) = \text{EdgesOut}(v_1, X) \cup \{\text{the edges of } G\}$ and $\text{EdgesOut}(v_1, X)$ misses $\{\text{the edges of } G\}$.
- (47) If $v' = v_2$ and $v_1 \neq v_2$ and the edges of $G \in X$, then $\text{EdgesIn}(v', X) = \text{EdgesIn}(v_2, X) \cup \{\text{the edges of } G\}$ and $\text{EdgesIn}(v_2, X)$ misses $\{\text{the edges of } G\}$.
- (48) If $v' = v$ and $v \neq v_1$ and $v \neq v_2$, then $\text{EdgesIn}(v', X) = \text{EdgesIn}(v, X)$.
- (49) If $v' = v$ and $v \neq v_1$ and $v \neq v_2$, then $\text{EdgesOut}(v', X) = \text{EdgesOut}(v, X)$.
- (50) If the edges of $G \notin \text{rng } p'$, then p' is a path of G .
- (51) If the edges of $G \notin \text{rng } p'$ and $v_3 = v'_1$ and v'_1 is vertex sequence of p' , then v_3 is vertex sequence of p' .

Let G be a connected graph and let v_1, v_2 be vertices of G . Observe that $\text{AddNewEdge}(v_1, v_2)$ is connected.

For simplicity, we use the following convention: G is a finite graph, v, v_1, v_2 are vertices of G , v_3 is a finite sequence of elements of the vertices of G , and v' is a vertex of $\text{AddNewEdge}(v_1, v_2)$.

The following two propositions are true:

- (52) If $v' = v$ and $v_1 \neq v_2$ and $v = v_1$ or $v = v_2$ and the edges of $G \in X$, then $\text{Degree}(v', X) = \text{Degree}(v, X) + 1$.
- (53) If $v' = v$ and $v \neq v_1$ and $v \neq v_2$, then $\text{Degree}(v', X) = \text{Degree}(v, X)$.

4. SOME PROPERTIES OF AND OPERATIONS ON CYCLES

We now state two propositions:

- (54) For every cyclic path c of G holds $\text{Degree}(v, \text{rng } c)$ is even.
- (55) Let c be a path of G . Suppose c is non cyclic and v_3 is vertex sequence of c . Then $\text{Degree}(v, \text{rng } c)$ is even if and only if $v \neq v_3(1)$ and $v \neq v_3(\text{len } v_3)$.

In the sequel G is a graph, v is a vertex of G , and v_3 is a finite sequence of elements of the vertices of G .

Let G be a graph. The functor $G\text{-CycleSet}$ yields a non empty set of finite sequences of the edges of G and is defined by:

(Def. 8) For every set x holds $x \in G\text{-CycleSet}$ iff x is a cyclic path of G .

One can prove the following two propositions:

- (56) \emptyset is an element of $G\text{-CycleSet}$.
- (57) Let c be an element of $G\text{-CycleSet}$. Suppose $v \in G\text{-VSet}(\text{rng } c)$. Then $\{c'; c' \text{ ranges over elements of } G\text{-CycleSet: } \text{rng } c' = \text{rng } c \wedge \bigvee_{v_3} (v_3 \text{ is vertex sequence of } c' \wedge v_3(1) = v)\}$ is a non empty subset of $G\text{-CycleSet}$.

Let us consider G, v and let c be an element of $G\text{-CycleSet}$. Let us assume that $v \in G\text{-VSet}(\text{rng } c)$. The functor $c \circ v$ yielding an element of $G\text{-CycleSet}$ is defined as follows:

(Def. 9) $c \circ v = \text{choose}(\{c'; c' \text{ ranges over elements of } G\text{-CycleSet: } \text{rng } c' = \text{rng } c \wedge \bigvee_{v_3} (v_3 \text{ is vertex sequence of } c' \wedge v_3(1) = v)\})$.

Let G be a graph and let c_1, c_2 be elements of $G\text{-CycleSet}$. Let us assume that $G\text{-VSet}(\text{rng } c_1)$ meets $G\text{-VSet}(\text{rng } c_2)$ and $\text{rng } c_1$ misses $\text{rng } c_2$. The functor $\text{CatCycles}(c_1, c_2)$ yields an element of $G\text{-CycleSet}$ and is defined as follows:

(Def. 10) There exists a vertex v of G such that $v = \text{choose}((G\text{-VSet}(\text{rng } c_1)) \cap (G\text{-VSet}(\text{rng } c_2)))$ and $\text{CatCycles}(c_1, c_2) = (c_1 \circ v) \wedge (c_2 \circ v)$.

The following proposition is true

- (58) Let G be a graph and c_1, c_2 be elements of $G\text{-CycleSet}$. Suppose $G\text{-VSet}(\text{rng } c_1)$ meets $G\text{-VSet}(\text{rng } c_2)$ but $\text{rng } c_1$ misses $\text{rng } c_2$ but $c_1 \neq \emptyset$ or $c_2 \neq \emptyset$. Then $\text{CatCycles}(c_1, c_2)$ is non empty.

In the sequel G is a finite graph, v is a vertex of G , and v_3 is a finite sequence of elements of the vertices of G .

Let us consider G, v and let X be a set. Let us assume that $\text{Degree}(v, X) \neq 0$. The functor $X\text{-PathSet}(v)$ yields a non empty set of finite sequences of the edges of G and is defined as follows:

- (Def. 11) $X\text{-PathSet}(v) = \{c; c \text{ ranges over elements of } X^*: c \text{ is a path of } G \wedge c \text{ is non empty} \wedge \bigvee_{v_3} (v_3 \text{ is vertex sequence of } c \wedge v_3(1) = v)\}$.

We now state the proposition

- (59) For every element p of $X\text{-PathSet}(v)$ and for every finite set Y such that $Y = \text{the edges of } G$ and $\text{Degree}(v, X) \neq 0$ holds $\text{len } p \leq \text{card } Y$.

Let us consider G, v and let X be a set. Let us assume that for every vertex v of G holds $\text{Degree}(v, X)$ is even and $\text{Degree}(v, X) \neq 0$. The functor $X\text{-CycleSet}v$ yields a non empty subset of $G\text{-CycleSet}$ and is defined as follows:

- (Def. 12) $X\text{-CycleSet}v = \{c; c \text{ ranges over elements of } G\text{-CycleSet}: \text{rng } c \subseteq X \wedge c \text{ is non empty} \wedge \bigvee_{v_3} (v_3 \text{ is vertex sequence of } c \wedge v_3(1) = v)\}$.

The following two propositions are true:

- (60) If $\text{Degree}(v, X) \neq 0$ and for every v holds $\text{Degree}(v, X)$ is even, then for every element c of $X\text{-CycleSet}v$ holds c is non empty and $\text{rng } c \subseteq X$ and $v \in G\text{-VSet}(\text{rng } c)$.
- (61) Let G be a finite connected graph and c be an element of $G\text{-CycleSet}$. Suppose $\text{rng } c \neq \text{the edges of } G$ and c is non empty. Then $\{v'; v' \text{ ranges over vertices of } G: v' \in G\text{-VSet}(\text{rng } c) \wedge \text{the degree of } v' \neq \text{Degree}(v', \text{rng } c)\}$ is a non empty subset of the vertices of G .

Let G be a finite connected graph and let c be an element of $G\text{-CycleSet}$. Let us assume that $\text{rng } c \neq \text{the edges of } G$ and c is non empty. The functor $\text{ExtendCycle } c$ yields an element of $G\text{-CycleSet}$ and is defined by the condition (Def. 13).

- (Def. 13) There exists an element c' of $G\text{-CycleSet}$ and there exists a vertex v of G such that $v = \text{choose}(\{v'; v' \text{ ranges over vertices of } G: v' \in G\text{-VSet}(\text{rng } c) \wedge \text{the degree of } v' \neq \text{Degree}(v', \text{rng } c)\})$ and $c' = \text{choose}(\text{((the edges of } G) \setminus \text{rng } c)\text{-CycleSet}v)$ and $\text{ExtendCycle } c = \text{CatCycles}(c, c')$.

Next we state the proposition

- (62) Let G be a finite connected graph and c be an element of $G\text{-CycleSet}$. Suppose $\text{rng } c \neq \text{the edges of } G$ and c is non empty and for every vertex v of G holds the degree of v is even. Then $\text{ExtendCycle } c$ is non empty and $\text{card } \text{rng } c < \text{card } \text{rng } \text{ExtendCycle } c$.

5. EULER CIRCUITS AND PATHS

Let G be a graph and let p be a path of G . We say that p is Eulerian if and only if:

- (Def. 14) $\text{rng } p = \text{the edges of } G$.

One can prove the following three propositions:

- (63) Let G be a connected graph, p be a path of G , and v_3 be a finite sequence of elements of the vertices of G . Suppose p is Eulerian and v_3 is vertex sequence of p . Then $\text{rng } v_3 = \text{the vertices of } G$.

- (64) Let G be a finite connected graph. Then there exists a cyclic path of G which is Eulerian if and only if for every vertex v of G holds the degree of v is even.
- (65) Let G be a finite connected graph. Then there exists a path of G which is non cyclic and Eulerian if and only if there exist vertices v_1, v_2 of G such that $v_1 \neq v_2$ and for every vertex v of G holds the degree of v is even iff $v \neq v_1$ and $v \neq v_2$.

ACKNOWLEDGMENTS

We would like to thank Cz. Byliński for many helpful comments.

REFERENCES

- [1] Grzegorz Bancerek. Cardinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/card_1.html.
- [2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/nat_1.html.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.
- [4] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [5] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [6] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/partfun1.html>.
- [7] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [8] Czesław Byliński. A classical first order language. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/cqc_lang.html.
- [9] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_2.html.
- [10] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/funct_4.html.
- [11] Czesław Byliński and Piotr Rudnicki. The correspondence between monotonic many sorted signatures and well-founded graphs. Part I. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/msscyc_1.html.
- [12] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.
- [13] Krzysztof Hryniewiecki. Graphs. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/graph_1.html.
- [14] Yatsuka Nakamura and Piotr Rudnicki. Vertex sequences induced by chains. *Journal of Formalized Mathematics*, 7, 1995. http://mizar.org/JFM/Vol7/graph_2.html.
- [15] Piotr Rudnicki and Andrzej Trybulec. Abian's fixed point theorem. *Journal of Formalized Mathematics*, 9, 1997. <http://mizar.org/JFM/Vol9/abian.html>.
- [16] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [17] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [18] Michał J. Trybulec. Integers. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/int_1.html.
- [19] Wojciech A. Trybulec. Pigeon hole principle. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_4.html.
- [20] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

- [21] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received July 29, 1997

Published January 2, 2004
