

Hilbert Positive Propositional Calculus

Adam Grabowski
University of Białystok

MML Identifier: HILBERT1.

WWW: <http://mizar.org/JFM/Vol11/hilbert1.html>

The articles [4], [6], [5], [7], [3], [1], and [2] provide the notation and terminology for this paper.

1. DEFINITION OF THE LANGUAGE

Let D be a set. We say that D has VERUM if and only if:

(Def. 1) $\langle 0 \rangle \in D$.

Let D be a set. We say that D has implication if and only if:

(Def. 2) For all finite sequences p, q such that $p \in D$ and $q \in D$ holds $\langle 1 \rangle \wedge p \wedge q \in D$.

Let D be a set. We say that D has conjunction if and only if:

(Def. 3) For all finite sequences p, q such that $p \in D$ and $q \in D$ holds $\langle 2 \rangle \wedge p \wedge q \in D$.

Let D be a set. We say that D has propositional variables if and only if:

(Def. 4) For every natural number n holds $\langle 3 + n \rangle \in D$.

Let D be a set. We say that D is HP-closed if and only if:

(Def. 5) $D \subseteq \mathbb{N}^*$ and D has VERUM, implication, conjunction, and propositional variables.

Let us observe that every set which is HP-closed is also non empty and has VERUM, implication, conjunction, and propositional variables and every subset of \mathbb{N}^* which has VERUM, implication, conjunction, and propositional variables is also HP-closed.

The set HP-WFF is defined by:

(Def. 6) HP-WFF is HP-closed and for every set D such that D is HP-closed holds $\text{HP-WFF} \subseteq D$.

Let us mention that HP-WFF is HP-closed.

Let us note that there exists a set which is HP-closed and non empty.

One can check that every element of HP-WFF is relation-like and function-like.

Let us observe that every element of HP-WFF is finite sequence-like.

A HP-formula is an element of HP-WFF.

The HP-formula VERUM is defined by:

(Def. 7) $\text{VERUM} = \langle 0 \rangle$.

Let p, q be elements of HP-WFF. The functor $p \Rightarrow q$ yielding a HP-formula is defined as follows:

(Def. 8) $p \Rightarrow q = \langle 1 \rangle \wedge p \wedge q$.

The functor $p \wedge q$ yields a HP-formula and is defined as follows:

(Def. 9) $p \wedge q = \langle 2 \rangle \wedge p \wedge q$.

We follow the rules: T, X, Y are subsets of HP-WFF and p, q, r, s are elements of HP-WFF.

Let T be a subset of HP-WFF. We say that T is Hilbert theory if and only if the conditions (Def. 10) are satisfied.

(Def. 10)(i) VERUM $\in T$, and

(ii) for all elements p, q, r of HP-WFF holds $p \Rightarrow (q \Rightarrow p) \in T$ and $(p \Rightarrow (q \Rightarrow r)) \Rightarrow ((p \Rightarrow q) \Rightarrow (p \Rightarrow r)) \in T$ and $p \wedge q \Rightarrow p \in T$ and $p \wedge q \Rightarrow q \in T$ and $p \Rightarrow (q \Rightarrow p \wedge q) \in T$ and if $p \in T$ and $p \Rightarrow q \in T$, then $q \in T$.

Let us consider X . The functor CnPos X yielding a subset of HP-WFF is defined as follows:

(Def. 11) $r \in \text{CnPos}X$ iff for every T such that T is Hilbert theory and $X \subseteq T$ holds $r \in T$.

The subset HP_TAUT of HP-WFF is defined as follows:

(Def. 12) HP_TAUT = CnPos $\emptyset_{\text{HP-WFF}}$.

The following propositions are true:

- (1) VERUM $\in \text{CnPos}X$.
- (2) $p \Rightarrow (q \Rightarrow p \wedge q) \in \text{CnPos}X$.
- (3) $(p \Rightarrow (q \Rightarrow r)) \Rightarrow ((p \Rightarrow q) \Rightarrow (p \Rightarrow r)) \in \text{CnPos}X$.
- (4) $p \Rightarrow (q \Rightarrow p) \in \text{CnPos}X$.
- (5) $p \wedge q \Rightarrow p \in \text{CnPos}X$.
- (6) $p \wedge q \Rightarrow q \in \text{CnPos}X$.
- (7) If $p \in \text{CnPos}X$ and $p \Rightarrow q \in \text{CnPos}X$, then $q \in \text{CnPos}X$.
- (8) If T is Hilbert theory and $X \subseteq T$, then $\text{CnPos}X \subseteq T$.
- (9) $X \subseteq \text{CnPos}X$.
- (10) If $X \subseteq Y$, then $\text{CnPos}X \subseteq \text{CnPos}Y$.
- (11) $\text{CnPos} \text{CnPos}X = \text{CnPos}X$.

Let X be a subset of HP-WFF. Observe that CnPos X is Hilbert theory.

The following propositions are true:

- (12) T is Hilbert theory iff $\text{CnPos}T = T$.
- (13) If T is Hilbert theory, then HP_TAUT $\subseteq T$.

Let us note that HP_TAUT is Hilbert theory.

2. THE TAUTOLOGIES OF THE HILBERT CALCULUS - IMPLICATIONAL PART

The following propositions are true:

- (14) $p \Rightarrow p \in \text{HP_TAUT}$.
- (15) If $q \in \text{HP_TAUT}$, then $p \Rightarrow q \in \text{HP_TAUT}$.
- (16) $p \Rightarrow \text{VERUM} \in \text{HP_TAUT}$.
- (17) $(p \Rightarrow q) \Rightarrow (p \Rightarrow p) \in \text{HP_TAUT}$.
- (18) $(q \Rightarrow p) \Rightarrow (p \Rightarrow p) \in \text{HP_TAUT}$.
- (19) $(q \Rightarrow r) \Rightarrow ((p \Rightarrow q) \Rightarrow (p \Rightarrow r)) \in \text{HP_TAUT}$.
- (20) If $p \Rightarrow (q \Rightarrow r) \in \text{HP_TAUT}$, then $q \Rightarrow (p \Rightarrow r) \in \text{HP_TAUT}$.
- (21) $(p \Rightarrow q) \Rightarrow ((q \Rightarrow r) \Rightarrow (p \Rightarrow r)) \in \text{HP_TAUT}$.
- (22) If $p \Rightarrow q \in \text{HP_TAUT}$, then $(q \Rightarrow r) \Rightarrow (p \Rightarrow r) \in \text{HP_TAUT}$.
- (23) If $p \Rightarrow q \in \text{HP_TAUT}$ and $q \Rightarrow r \in \text{HP_TAUT}$, then $p \Rightarrow r \in \text{HP_TAUT}$.
- (24) $(p \Rightarrow (q \Rightarrow r)) \Rightarrow ((s \Rightarrow q) \Rightarrow (p \Rightarrow (s \Rightarrow r))) \in \text{HP_TAUT}$.
- (25) $((p \Rightarrow q) \Rightarrow r) \Rightarrow (q \Rightarrow r) \in \text{HP_TAUT}$.
- (26) $(p \Rightarrow (q \Rightarrow r)) \Rightarrow (q \Rightarrow (p \Rightarrow r)) \in \text{HP_TAUT}$.
- (27) $(p \Rightarrow (p \Rightarrow q)) \Rightarrow (p \Rightarrow q) \in \text{HP_TAUT}$.
- (28) $q \Rightarrow ((q \Rightarrow p) \Rightarrow p) \in \text{HP_TAUT}$.
- (29) If $s \Rightarrow (q \Rightarrow p) \in \text{HP_TAUT}$ and $q \in \text{HP_TAUT}$, then $s \Rightarrow p \in \text{HP_TAUT}$.

3. CONJUNCTIONAL PART OF THE CALCULUS

The following propositions are true:

- (30) $p \Rightarrow p \wedge p \in \text{HP_TAUT}$.
- (31) $p \wedge q \in \text{HP_TAUT}$ iff $p \in \text{HP_TAUT}$ and $q \in \text{HP_TAUT}$.
- (32) $p \wedge q \in \text{HP_TAUT}$ iff $q \wedge p \in \text{HP_TAUT}$.
- (33) $(p \wedge q \Rightarrow r) \Rightarrow (p \Rightarrow (q \Rightarrow r)) \in \text{HP_TAUT}$.
- (34) $(p \Rightarrow (q \Rightarrow r)) \Rightarrow (p \wedge q \Rightarrow r) \in \text{HP_TAUT}$.
- (35) $(r \Rightarrow p) \Rightarrow ((r \Rightarrow q) \Rightarrow (r \Rightarrow p \wedge q)) \in \text{HP_TAUT}$.
- (36) $(p \Rightarrow q) \wedge p \Rightarrow q \in \text{HP_TAUT}$.
- (37) $(p \Rightarrow q) \wedge p \wedge s \Rightarrow q \in \text{HP_TAUT}$.
- (38) $(q \Rightarrow s) \Rightarrow (p \wedge q \Rightarrow s) \in \text{HP_TAUT}$.
- (39) $(q \Rightarrow s) \Rightarrow (q \wedge p \Rightarrow s) \in \text{HP_TAUT}$.
- (40) $(p \wedge s \Rightarrow q) \Rightarrow (p \wedge s \Rightarrow q \wedge s) \in \text{HP_TAUT}$.
- (41) $(p \Rightarrow q) \Rightarrow (p \wedge s \Rightarrow q \wedge s) \in \text{HP_TAUT}$.
- (42) $(p \Rightarrow q) \wedge (p \wedge s) \Rightarrow q \wedge s \in \text{HP_TAUT}$.

- (43) $p \wedge q \Rightarrow q \wedge p \in \text{HP_TAUT}$.
- (44) $(p \Rightarrow q) \wedge (p \wedge s) \Rightarrow s \wedge q \in \text{HP_TAUT}$.
- (45) $(p \Rightarrow q) \Rightarrow (p \wedge s \Rightarrow s \wedge q) \in \text{HP_TAUT}$.
- (46) $(p \Rightarrow q) \Rightarrow (s \wedge p \Rightarrow s \wedge q) \in \text{HP_TAUT}$.
- (47) $p \wedge (s \wedge q) \Rightarrow p \wedge (q \wedge s) \in \text{HP_TAUT}$.
- (48) $(p \Rightarrow q) \wedge (p \Rightarrow s) \Rightarrow (p \Rightarrow q \wedge s) \in \text{HP_TAUT}$.
- (49) $p \wedge q \wedge s \Rightarrow p \wedge (q \wedge s) \in \text{HP_TAUT}$.
- (50) $p \wedge (q \wedge s) \Rightarrow p \wedge q \wedge s \in \text{HP_TAUT}$.

REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/nat_1.html.
- [2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/finseq_1.html.
- [3] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/func_1.html.
- [4] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [5] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [6] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/subset_1.html.
- [7] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/relat_1.html.

Received February 20, 1999

Published January 2, 2004
