

The Canonical Formulae

Andrzej Trybulec
University of Białystok

MML Identifier: HILBERT3.

WWW: <http://mizar.org/JFM/Vol12/hilbert3.html>

The articles [18], [9], [25], [23], [24], [26], [6], [4], [2], [17], [8], [7], [5], [19], [10], [1], [3], [13], [14], [20], [21], [15], [12], [16], [11], and [22] provide the notation and terminology for this paper.

1. PRELIMINARIES

One can prove the following three propositions:

- (1) For every integer i holds i is even iff $i - 1$ is odd.
- (2) For every integer i holds i is odd iff $i - 1$ is even.
- (3) Let X be a trivial set and x be a set. Suppose $x \in X$. Let f be a function from X into X . Then x is a fixpoint of f .

Let A, B, C be sets. Observe that every function from A into C^B is function yielding.

The following three propositions are true:

- (4) For every function yielding function f holds $\text{Sub}_f \text{rng } f = \text{rng } f$.
- (5) For all sets A, B, x and for every function f such that $x \in A$ and $f \in B^A$ holds $f(x) \in B$.
- (6) For all sets A, B, C such that if $C = \emptyset$, then $B = \emptyset$ or $A = \emptyset$ and for every function f from A into C^B holds $\text{dom}_\kappa f(\kappa) = A \mapsto B$.

Let us mention that \emptyset is function yielding.

In the sequel n denotes a natural number and p, q, r denote elements of HP-WFF.

The following proposition is true

- (7) For every set x holds $\emptyset(x) = \emptyset$.

Let A be a set and let B be a functional set. One can verify that every function from A into B is function yielding.

One can prove the following propositions:

- (8) For every set X and for every subset A of X holds $[0 \mapsto 1, 1 \mapsto 0] \cdot \mathcal{X}_{A,X} = \mathcal{X}_{A^c,X}$.
- (9) For every set X and for every subset A of X holds $[0 \mapsto 1, 1 \mapsto 0] \cdot \mathcal{X}_{A^c,X} = \mathcal{X}_{A,X}$.
- (10) For all sets a, b, x, y, x', y' such that $a \neq b$ and $[a \mapsto x, b \mapsto y] = [a \mapsto x', b \mapsto y']$ holds $x = x'$ and $y = y'$.

- (11) For all sets a, b, x, y, X, Y such that $a \neq b$ and $x \in X$ and $y \in Y$ holds $[a \mapsto x, b \mapsto y] \in \prod[a \mapsto X, b \mapsto Y]$.
- (12) For every non empty set D and for every function f from 2 into D there exist elements d_1, d_2 of D such that $f = [0 \mapsto d_1, 1 \mapsto d_2]$.
- (13) For all sets a, b, c, d such that $a \neq b$ holds $[a \mapsto c, b \mapsto d] \cdot [a \mapsto b, b \mapsto a] = [a \mapsto d, b \mapsto c]$.
- (14) For all sets a, b, c, d and for every function f such that $a \neq b$ and $c \in \text{dom } f$ and $d \in \text{dom } f$ holds $f \cdot [a \mapsto c, b \mapsto d] = [a \mapsto f(c), b \mapsto f(d)]$.

2. THE CARTESIAN PRODUCT OF FUNCTIONS AND THE FREGE FUNCTION

Let f, g be one-to-one functions. Note that $[:f, g:]$ is one-to-one.

We now state several propositions:

- (15) Let A, B be non empty sets, C, D be sets, f be a function from C into A , and g be a function from D into B . Then $\pi_1(A \times B) \cdot [:f, g:] = f \cdot \pi_1(C \times D)$.
- (16) Let A, B be non empty sets, C, D be sets, f be a function from C into A , and g be a function from D into B . Then $\pi_2(A \times B) \cdot [:f, g:] = g \cdot \pi_2(C \times D)$.
- (17) For every function g holds $\emptyset \leftrightarrow g = \emptyset$.
- (18) For every function yielding function f and for all functions g, h holds $f \leftrightarrow g \cdot h = (f \cdot h) \leftrightarrow (g \cdot h)$.
- (19) Let C be a set, A be a non empty set, f be a function from A into $C^{(\emptyset \text{ qua set})}$, and g be a function from A into \emptyset . Then $\text{rng}(f \leftrightarrow g) = \{\emptyset\}$.
- (20) Let A, B, C be sets such that if $B = \emptyset$, then $A = \emptyset$. Let f be a function from A into C^B and g be a function from A into B . Then $\text{rng}(f \leftrightarrow g) \subseteq C$.
- (21) For all sets A, B, C such that if $C = \emptyset$, then $B = \emptyset$ or $A = \emptyset$ and for every function f from A into C^B holds $\text{dom Frege}(f) = B^A$.
- (23)¹ For all sets A, B, C such that if $C = \emptyset$, then $B = \emptyset$ or $A = \emptyset$ and for every function f from A into C^B holds $\text{rng Frege}(f) \subseteq C^A$.
- (24) Let A, B, C be sets such that if $C = \emptyset$, then $B = \emptyset$ or $A = \emptyset$. Let f be a function from A into C^B . Then $\text{Frege}(f)$ is a function from B^A into C^A .

3. ABOUT PERMUTATIONS

The following proposition is true

- (25) For all sets A, B and for every permutation P of A and for every permutation Q of B holds $[:P, Q:]$ is bijective.

Let A, B be non empty sets, let P be a permutation of A , and let Q be a function from B into B . The functor $P \Rightarrow Q$ yields a function from B^A into B^A and is defined by:

(Def. 1) For every function f from A into B holds $(P \Rightarrow Q)(f) = Q \cdot f \cdot P^{-1}$.

Let A, B be non empty sets, let P be a permutation of A , and let Q be a permutation of B . One can verify that $P \Rightarrow Q$ is bijective.

We now state three propositions:

¹ The proposition (22) has been removed.

- (26) Let A, B be non empty sets, P be a permutation of A , Q be a permutation of B , and f be a function from A into B . Then $(P \Rightarrow Q)^{-1}(f) = Q^{-1} \cdot f \cdot P$.
- (27) For all non empty sets A, B and for every permutation P of A and for every permutation Q of B holds $(P \Rightarrow Q)^{-1} = P^{-1} \Rightarrow Q^{-1}$.
- (28) Let A, B, C be non empty sets, f be a function from A into C^B , g be a function from A into B , P be a permutation of B , and Q be a permutation of C . Then $((P \Rightarrow Q) \cdot f) \leftarrow P (P \cdot g) = Q \cdot f \leftarrow P g$.

4. SET VALUATIONS

A SetValuation is a non-empty many sorted set indexed by \mathbb{N} .

In the sequel V is a SetValuation.

Let us consider V . The functor $\text{SetVal}V$ yields a many sorted set indexed by HP-WFF and is defined by the conditions (Def. 2).

- (Def. 2)(i) $(\text{SetVal}V)(\text{VERUM}) = 1$,
- (ii) for every n holds $(\text{SetVal}V)(\text{prop}n) = V(n)$, and
- (iii) for all p, q holds $(\text{SetVal}V)(p \wedge q) = [:(\text{SetVal}V)(p), (\text{SetVal}V)(q):]$ and $(\text{SetVal}V)(p \Rightarrow q) = (\text{SetVal}V)(q)^{(\text{SetVal}V)(p)}$.

Let us consider V, p . The functor $\text{SetVal}(V, p)$ is defined by:

- (Def. 3) $\text{SetVal}(V, p) = (\text{SetVal}V)(p)$.

Let us consider V, p . One can check that $\text{SetVal}(V, p)$ is non empty.

One can prove the following propositions:

- (29) $\text{SetVal}(V, \text{VERUM}) = 1$.
- (30) $\text{SetVal}(V, \text{prop}n) = V(n)$.
- (31) $\text{SetVal}(V, p \wedge q) = [:\text{SetVal}(V, p), \text{SetVal}(V, q):]$.
- (32) $\text{SetVal}(V, p \Rightarrow q) = (\text{SetVal}(V, q))^{\text{SetVal}(V, p)}$.

Let us consider V, p, q . Observe that $\text{SetVal}(V, p \Rightarrow q)$ is functional.

Let us consider V, p, q, r . One can check that every element of $\text{SetVal}(V, p \Rightarrow (q \Rightarrow r))$ is function yielding.

Let us consider V, p, q, r . One can check that there exists a function from $\text{SetVal}(V, p \Rightarrow q)$ into $\text{SetVal}(V, p \Rightarrow r)$ which is function yielding and there exists an element of $\text{SetVal}(V, p \Rightarrow (q \Rightarrow r))$ which is function yielding.

5. PERMUTING SET VALUATIONS

Let us consider V . A function is called a permutation of V if:

- (Def. 4) $\text{dom}it = \mathbb{N}$ and for every n holds $it(n)$ is a permutation of $V(n)$.

In the sequel P is a permutation of V .

Let us consider V, P . The functor $\text{Perm}P$ yields a many sorted function from $\text{SetVal}V$ into $\text{SetVal}V$ and is defined by the conditions (Def. 5).

- (Def. 5)(i) $(\text{Perm}P)(\text{VERUM}) = \text{id}_1$,
- (ii) for every n holds $(\text{Perm}P)(\text{prop}n) = P(n)$, and
- (iii) for all p, q there exists a permutation p' of $\text{SetVal}(V, p)$ and there exists a permutation q' of $\text{SetVal}(V, q)$ such that $p' = (\text{Perm}P)(p)$ and $q' = (\text{Perm}P)(q)$ and $(\text{Perm}P)(p \wedge q) = [':p', q':]$ and $(\text{Perm}P)(p \Rightarrow q) = p' \Rightarrow q'$.

Let us consider V, P, p . The functor $\text{Perm}(P, p)$ yields a function from $\text{SetVal}(V, p)$ into $\text{SetVal}(V, p)$ and is defined as follows:

(Def. 6) $\text{Perm}(P, p) = (\text{Perm}P)(p)$.

Next we state four propositions:

(33) $\text{Perm}(P, \text{VERUM}) = \text{id}_{\text{SetVal}(V, \text{VERUM})}$.

(34) $\text{Perm}(P, \text{prop } n) = P(n)$.

(35) $\text{Perm}(P, p \wedge q) = [\text{Perm}(P, p), \text{Perm}(P, q)]$.

(36) For every permutation p' of $\text{SetVal}(V, p)$ and for every permutation q' of $\text{SetVal}(V, q)$ such that $p' = \text{Perm}(P, p)$ and $q' = \text{Perm}(P, q)$ holds $\text{Perm}(P, p \Rightarrow q) = p' \Rightarrow q'$.

Let us consider V, P, p . Observe that $\text{Perm}(P, p)$ is bijective.

One can prove the following propositions:

(37) For every function g from $\text{SetVal}(V, p)$ into $\text{SetVal}(V, q)$ holds $(\text{Perm}(P, p \Rightarrow q))(g) = \text{Perm}(P, q) \cdot g \cdot (\text{Perm}(P, p))^{-1}$.

(38) For every function g from $\text{SetVal}(V, p)$ into $\text{SetVal}(V, q)$ holds $(\text{Perm}(P, p \Rightarrow q))^{-1}(g) = (\text{Perm}(P, q))^{-1} \cdot g \cdot \text{Perm}(P, p)$.

(39) For all functions f, g from $\text{SetVal}(V, p)$ into $\text{SetVal}(V, q)$ such that $f = (\text{Perm}(P, p \Rightarrow q))(g)$ holds $\text{Perm}(P, q) \cdot g = f \cdot \text{Perm}(P, p)$.

(40) Let given V, P be a permutation of V , and x be a set. Suppose x is a fixpoint of $\text{Perm}(P, p)$. Let f be a function. If f is a fixpoint of $\text{Perm}(P, p \Rightarrow q)$, then $f(x)$ is a fixpoint of $\text{Perm}(P, q)$.

6. CANONICAL FORMULAE

Let us consider p . We say that p is canonical if and only if:

(Def. 7) For every V there exists a set x such that for every permutation P of V holds x is a fixpoint of $\text{Perm}(P, p)$.

Let us note that VERUM is canonical.

Next we state several propositions:

(41) $p \Rightarrow (q \Rightarrow p)$ is canonical.

(42) $(p \Rightarrow (q \Rightarrow r)) \Rightarrow ((p \Rightarrow q) \Rightarrow (p \Rightarrow r))$ is canonical.

(43) $p \wedge q \Rightarrow p$ is canonical.

(44) $p \wedge q \Rightarrow q$ is canonical.

(45) $p \Rightarrow (q \Rightarrow p \wedge q)$ is canonical.

(46) If p is canonical and $p \Rightarrow q$ is canonical, then q is canonical.

(47) If $p \in \text{HP_TAUT}$, then p is canonical.

Let us mention that there exists an element of HP-WFF which is canonical.

7. PSEUDO-CANONICAL FORMULAE

Let us consider p . We say that p is pseudo-canonical if and only if:

(Def. 8) For every V and for every permutation P of V holds there exists a set which is a fixpoint of $\text{Perm}(P, p)$.

Let us observe that every element of HP-WFF which is canonical is also pseudo-canonical. One can prove the following propositions:

- (48) $p \Rightarrow (q \Rightarrow p)$ is pseudo-canonical.
- (49) $(p \Rightarrow (q \Rightarrow r)) \Rightarrow ((p \Rightarrow q) \Rightarrow (p \Rightarrow r))$ is pseudo-canonical.
- (50) $p \wedge q \Rightarrow p$ is pseudo-canonical.
- (51) $p \wedge q \Rightarrow q$ is pseudo-canonical.
- (52) $p \Rightarrow (q \Rightarrow p \wedge q)$ is pseudo-canonical.
- (53) If p is pseudo-canonical and $p \Rightarrow q$ is pseudo-canonical, then q is pseudo-canonical.
- (54) Let given p, q , given V , and P be a permutation of V . Suppose there exists a set which is a fixpoint of $\text{Perm}(P, p)$ and there exists no set which is a fixpoint of $\text{Perm}(P, q)$. Then $p \Rightarrow q$ is not pseudo-canonical.
- (55) $((\text{prop}0 \Rightarrow \text{prop}1) \Rightarrow \text{prop}0) \Rightarrow \text{prop}0$ is not pseudo-canonical.

REFERENCES

- [1] Grzegorz Bancerek. Curried and uncurried functions. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/funct_5.html.
- [2] Grzegorz Bancerek. König's theorem. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/card_3.html.
- [3] Grzegorz Bancerek. Cartesian product of functions. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/funct_6.html.
- [4] Józef Białas. Group and field definitions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/realset1.html>.
- [5] Czesław Byliński. Basic functions and operations on functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_3.html.
- [6] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [7] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [8] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/partfun1.html>.
- [9] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [10] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/funct_4.html.
- [11] Adam Grabowski. Hilbert positive propositional calculus. *Journal of Formalized Mathematics*, 11, 1999. <http://mizar.org/JFM/Vol11/hilbert1.html>.
- [12] Małgorzata Korolkiewicz. Homomorphisms of many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg_3.html.
- [13] Beata Madras. Product of family of universal algebras. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/pralg_1.html.
- [14] Beata Madras. Products of many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/pralg_2.html.
- [15] Yatsuka Nakamura, Piotr Rudnicki, Andrzej Trybulec, and Pauline N. Kawamoto. Preliminaries to circuits, I. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/pre_circ.html.

- [16] Piotr Rudnicki and Andrzej Trybulec. Fixpoints in complete lattices. *Journal of Formalized Mathematics*, 8, 1996. <http://mizar.org/JFM/Vol8/knaster.html>.
- [17] Piotr Rudnicki and Andrzej Trybulec. Abian's fixed point theorem. *Journal of Formalized Mathematics*, 9, 1997. <http://mizar.org/JFM/Vol9/abian.html>.
- [18] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [19] Andrzej Trybulec. Function domains and Fränkel operator. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol2/fraenkel.html>.
- [20] Andrzej Trybulec. Many-sorted sets. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/pboole.html>.
- [21] Andrzej Trybulec. Many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg_1.html.
- [22] Andrzej Trybulec. Defining by structural induction in the positive propositional language. *Journal of Formalized Mathematics*, 11, 1999. <http://mizar.org/JFM/Vol11/hilbert2.html>.
- [23] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [24] Michał J. Trybulec. Integers. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/int_1.html.
- [25] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [26] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received July 4, 2000

Published January 2, 2004
