

Irrationality of e

Freek Wiedijk
University of Nijmegen

Summary. We prove the irrationality of square roots of prime numbers and of the number e . In order to be able to prove the last, a proof is given that `number_e = exp(1)` as defined in the Mizar library, that is that

$$\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = \sum_{k=0}^{\infty} \frac{1}{k!}$$

MML Identifier: IRRAT_1.

WWW: http://mizar.org/JFM/Vol11/irrat_1.html

The articles [21], [2], [18], [20], [1], [13], [4], [11], [9], [16], [15], [10], [3], [6], [17], [5], [12], [22], [8], [19], [7], and [14] provide the notation and terminology for this paper.

1. SQUARE ROOTS OF PRIMES ARE IRRATIONAL

For simplicity, we adopt the following convention: k, n, p, K, N are natural numbers, x, y, e_1 are real numbers, s_1, s_2, s_3 are sequences of real numbers, and s_4 is a finite sequence of elements of \mathbb{R} .

Let us consider x . We introduce x is irrational as an antonym of x is rational.

Let us consider x, y . We introduce x^y as a synonym of x^y .

One can prove the following propositions:

- (1) If p is prime, then \sqrt{p} is irrational.
- (2) There exist x, y such that x is irrational and y is irrational and x^y is rational.

2. A PROOF THAT $e = e$

The scheme *LambdaRealSeq* deals with a unary functor \mathcal{F} yielding a real number, and states that:

There exists s_1 such that for every n holds $s_1(n) = \mathcal{F}(n)$ and for all s_2, s_3 such that for every n holds $s_2(n) = \mathcal{F}(n)$ and for every n holds $s_3(n) = \mathcal{F}(n)$ holds $s_2 = s_3$

for all values of the parameter.

Let us consider k . The functor \mathbf{a}_k is a sequence of real numbers and is defined by:

(Def. 1) For every n holds $\mathbf{a}_k(n) = \frac{n-k}{n}$.

Let us consider k . The functor \mathbf{b}_k is a sequence of real numbers and is defined by:

(Def. 2) For every n holds $\mathbf{b}_k(n) = \binom{n}{k} \cdot n^{-k}$.

Let us consider n . The functor \mathbf{c}_n is a sequence of real numbers and is defined by:

(Def. 3) For every k holds $\mathbf{c}_n(k) = \binom{n}{k} \cdot n^{-k}$.

One can prove the following proposition

$$(3) \quad \mathbf{c}_n(k) = \mathbf{b}_k(n).$$

The sequence \mathbf{d} of real numbers is defined by:

$$(\text{Def. 4}) \quad \text{For every } n \text{ holds } \mathbf{d}(n) = \left(1 + \frac{1}{n}\right)^n.$$

The sequence \mathbf{e} of real numbers is defined by:

$$(\text{Def. 5}) \quad \text{For every } k \text{ holds } \mathbf{e}(k) = \frac{1}{k!}.$$

We now state a number of propositions:

$$(4) \quad \text{If } n > 0, \text{ then } n^{-(k+1)} = \frac{n^{-k}}{n}.$$

$$(5) \quad \text{For all real numbers } x, y, z, v, w \text{ holds } \frac{x}{y \cdot z \cdot \frac{v}{w}} = \frac{w}{z} \cdot \frac{x}{y \cdot v}.$$

$$(6) \quad \binom{n}{k+1} = \frac{n-k}{k+1} \cdot \binom{n}{k}.$$

$$(7) \quad \text{If } n > 0, \text{ then } \mathbf{b}_{k+1}(n) = \frac{1}{k+1} \cdot \mathbf{b}_k(n) \cdot \mathbf{a}_k(n).$$

$$(8) \quad \text{If } n > 0, \text{ then } \mathbf{a}_k(n) = 1 - \frac{k}{n}.$$

$$(9) \quad \mathbf{a}_k \text{ is convergent and } \lim(\mathbf{a}_k) = 1.$$

$$(10) \quad \text{For every } s_1 \text{ such that for every } n \text{ holds } s_1(n) = x \text{ holds } s_1 \text{ is convergent and } \lim s_1 = x.$$

$$(11) \quad \text{For every } n \text{ holds } \mathbf{b}_0(n) = 1.$$

$$(12) \quad \frac{1}{k+1} \cdot \frac{1}{k!} = \frac{1}{(k+1)!}.$$

$$(13) \quad \mathbf{b}_k \text{ is convergent and } \lim(\mathbf{b}_k) = \frac{1}{k!} \text{ and } \lim(\mathbf{b}_k) = \mathbf{e}(k).$$

$$(14) \quad \text{If } k < n, \text{ then } 0 < \mathbf{a}_k(n) \text{ and } \mathbf{a}_k(n) \leq 1.$$

$$(15) \quad \text{If } n > 0, \text{ then } 0 \leq \mathbf{b}_k(n) \text{ and } \mathbf{b}_k(n) \leq \frac{1}{k!} \text{ and } \mathbf{b}_k(n) \leq \mathbf{e}(k) \text{ and } 0 \leq \mathbf{c}_n(k) \text{ and } \mathbf{c}_n(k) \leq \frac{1}{k!} \text{ and } \mathbf{c}_n(k) \leq \mathbf{e}(k).$$

$$(16) \quad \text{For every } s_1 \text{ such that } s_1 \uparrow 1 \text{ is summable holds } s_1 \text{ is summable and } \sum s_1 = s_1(0) + \sum(s_1 \uparrow 1).$$

$$(17) \quad \text{Let } D \text{ be a non empty set and } s_4 \text{ be a finite sequence of elements of } D. \text{ If } 1 \leq k \text{ and } k < \text{len } s_4, \text{ then } (s_4)_{\uparrow 1}(k) = s_4(k+1).$$

$$(18) \quad \text{For every } s_4 \text{ such that } \text{len } s_4 > 0 \text{ holds } \sum s_4 = s_4(1) + \sum((s_4)_{\uparrow 1}).$$

$$(19) \quad \text{Let given } n \text{ and given } s_1, s_4. \text{ Suppose } \text{len } s_4 = n \text{ and for every } k \text{ such that } k < n \text{ holds } s_1(k) = s_4(k+1) \text{ and for every } k \text{ such that } k \geq n \text{ holds } s_1(k) = 0. \text{ Then } s_1 \text{ is summable and } \sum s_1 = \sum s_4.$$

$$(20) \quad \text{If } x \neq 0 \text{ and } y \neq 0 \text{ and } k \leq n, \text{ then } \langle \binom{n}{0} x^0 y^n, \dots, \binom{n}{n} x^n y^0 \rangle (k+1) = \binom{n}{k} \cdot x^{n-k} \cdot y^k.$$

$$(21) \quad \text{If } n > 0 \text{ and } k \leq n, \text{ then } \mathbf{c}_n(k) = \langle \binom{n}{0} 1^0 \left(\frac{1}{n}\right)^n, \dots, \binom{n}{n} 1^n \left(\frac{1}{n}\right)^0 \rangle (k+1).$$

$$(22) \quad \text{If } n > 0, \text{ then } \mathbf{c}_n \text{ is summable and } \sum(\mathbf{c}_n) = \left(1 + \frac{1}{n}\right)^n \text{ and } \sum(\mathbf{c}_n) = \mathbf{d}(n).$$

$$(23) \quad \mathbf{d} \text{ is convergent and } \lim \mathbf{d} = e.$$

$$(24) \quad \mathbf{e} \text{ is summable and } \sum \mathbf{e} = \exp 1.$$

$$(25) \quad \text{Let given } K \text{ and } d_1 \text{ be a sequence of real numbers. If for every } n \text{ holds } d_1(n) = \left(\sum_{\alpha=0}^k (\mathbf{c}_n)(\alpha)\right)_{\kappa \in \mathbb{N}(K)}, \text{ then } d_1 \text{ is convergent and } \lim d_1 = \left(\sum_{\alpha=0}^k \mathbf{e}(\alpha)\right)_{\kappa \in \mathbb{N}(K)}.$$

(26) If s_1 is convergent and $\lim s_1 = x$, then for every ϵ_1 such that $\epsilon_1 > 0$ there exists N such that for every n such that $n \geq N$ holds $s_1(n) > x - \epsilon_1$.

(27) Suppose that

(i) for every ϵ_1 such that $\epsilon_1 > 0$ there exists N such that for every n such that $n \geq N$ holds $s_1(n) > x - \epsilon_1$, and

(ii) there exists N such that for every n such that $n \geq N$ holds $s_1(n) \leq x$.

Then s_1 is convergent and $\lim s_1 = x$.

(28) If s_1 is summable, then for every ϵ_1 such that $\epsilon_1 > 0$ there exists K such that $(\sum_{\alpha=0}^k (s_1)(\alpha))_{k \in \mathbb{N}}(K) > \sum s_1 - \epsilon_1$.

(29) If $n \geq 1$, then $\mathbf{d}(n) \leq \sum \mathbf{e}$.

(30) If s_1 is summable and for every k holds $s_1(k) \geq 0$, then $\sum s_1 \geq (\sum_{\alpha=0}^k (s_1)(\alpha))_{k \in \mathbb{N}}(K)$.

(31) \mathbf{d} is convergent and $\lim \mathbf{d} = \sum \mathbf{e}$.

e can be characterized by the condition:

(Def. 6) $e = \sum \mathbf{e}$.

e can be characterized by the condition:

(Def. 7) $e = \exp 1$.

3. THE NUMBER e IS IRRATIONAL

The following propositions are true:

(32) If x is rational, then there exists n such that $n \geq 2$ and $n! \cdot x$ is integer.

(33) $n! \cdot \mathbf{e}(k) = \frac{n!}{k!}$.

(34) $\frac{n!}{k!} > 0$.

(35) If s_1 is summable and for every n holds $s_1(n) > 0$, then $\sum s_1 > 0$.

(36) $n! \cdot \sum(\mathbf{e} \uparrow (n+1)) > 0$.

(37) If $k \leq n$, then $\frac{n!}{k!}$ is integer.

(38) $n! \cdot (\sum_{\alpha=0}^k \mathbf{e}(\alpha))_{k \in \mathbb{N}}(n)$ is integer.

(39) If $x = \frac{1}{n+1}$, then $\frac{n!}{(n+k+1)!} \leq x^{k+1}$.

(40) If $n > 0$ and $x = \frac{1}{n+1}$, then $n! \cdot \sum(\mathbf{e} \uparrow (n+1)) \leq \frac{x}{1-x}$.

(41) For every real number n such that $n \geq 2$ and $x = \frac{1}{n+1}$ holds $\frac{x}{1-x} < 1$.

(42) e is irrational.

REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/nat_1.html.
- [2] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinal1.html>.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.
- [4] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [5] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_2.html.
- [6] Czesław Byliński. The sum and product of finite sequences of real numbers. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/rvsum_1.html.
- [7] Marek Chmur. The lattice of natural numbers and the sublattice of it. The set of prime numbers. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/nat_lat.html.
- [8] Andrzej Kondracki. Basic properties of rational numbers. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/rat_1.html.
- [9] Jarosław Kotowicz. Convergent sequences and the limit of sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/seq_2.html.
- [10] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/seqm_3.html.
- [11] Jarosław Kotowicz. Real sequences and basic operations on them. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/seq_1.html.
- [12] Jarosław Kotowicz. Functions and finite sequences of real numbers. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/rfinseq.html>.
- [13] Rafał Kwiatek. Factorial and Newton coefficients. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol2/newton.html>.
- [14] Rafał Kwiatek and Grzegorz Żwara. The divisibility of integers and integer relatively primes. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/int_2.html.
- [15] Jan Popiołek. Some properties of functions modul and signum. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/absvalue.html>.
- [16] Konrad Raczkowski and Andrzej Nędzusiak. Real exponents and logarithms. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol2/power.html>.
- [17] Konrad Raczkowski and Andrzej Nędzusiak. Series. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/series_1.html.
- [18] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [19] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers operations: min, max, square, and square root. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/square_1.html.
- [20] Michał J. Trybulec. Integers. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/int_1.html.
- [21] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [22] Yuguang Yang and Yasunari Shidama. Trigonometric functions and existence of circle ratio. *Journal of Formalized Mathematics*, 10, 1998. http://mizar.org/JFM/Vol10/sin_cos.html.

Received July 2, 1999

Published January 2, 2004
