

Banach Space of Bounded Linear Operators

Yasunari Shidama
Shinshu University
Nagano

Summary. On this article, the basic properties of linear spaces which are defined by the set of all linear operators from one linear space to another are described. Especially, the Banach space is introduced. This is defined by the set of all bounded linear operators.

MML Identifier: LOPBAN_1.

WWW: http://mizar.org/JFM/Vol15/lopban_1.html

The articles [24], [6], [23], [31], [25], [33], [32], [4], [5], [16], [22], [21], [3], [1], [2], [28], [27], [9], [7], [30], [14], [26], [17], [29], [18], [19], [8], [20], [13], [11], [12], [10], and [15] provide the notation and terminology for this paper.

1. REAL VECTOR SPACE OF OPERATORS

Let X be a set, let Y be a non empty set, let F be a function from $[\mathbb{R}, Y]$ into Y , let a be a real number, and let f be a function from X into Y . Then $F^\circ(a, f)$ is an element of Y^X .

One can prove the following two propositions:

- (1) Let X be a non empty set and Y be a non empty loop structure. Then there exists a binary operation A_1 on (the carrier of Y^X) such that for all elements f, g of (the carrier of Y^X) holds $A_1(f, g) = (\text{the addition of } Y)^\circ(f, g)$.
- (2) Let X be a non empty set and Y be a real linear space. Then there exists a function M_1 from $[\mathbb{R}, (\text{the carrier of } Y^X)]$ into (the carrier of Y^X) such that for every real number r and for every element f of (the carrier of Y^X) and for every element s of X holds $M_1(\langle r, f \rangle)(s) = r \cdot f(s)$.

Let X be a non empty set and let Y be a non empty loop structure. The functor $\text{FuncAdd}(X, Y)$ yielding a binary operation on (the carrier of Y^X) is defined as follows:

(Def. 1) For all elements f, g of (the carrier of Y^X) holds $(\text{FuncAdd}(X, Y))(f, g) = (\text{the addition of } Y)^\circ(f, g)$.

Let X be a non empty set and let Y be a real linear space. The functor $\text{FuncExtMult}(X, Y)$ yields a function from $[\mathbb{R}, (\text{the carrier of } Y^X)]$ into (the carrier of Y^X) and is defined by the condition (Def. 2).

(Def. 2) Let a be a real number, f be an element of (the carrier of Y^X), and x be an element of X . Then $(\text{FuncExtMult}(X, Y))(\langle a, f \rangle)(x) = a \cdot f(x)$.

Let X be a set and let Y be a non empty zero structure. The functor $\text{FuncZero}(X, Y)$ yields an element of (the carrier of Y^X) and is defined as follows:

(Def. 3) $\text{FuncZero}(X, Y) = X \mapsto 0_Y$.

We adopt the following rules: X denotes a non empty set, Y denotes a real linear space, and f, g, h denote elements of $(\text{the carrier of } Y)^X$.

Next we state two propositions:

- (3) Let Y be a non empty loop structure and f, g, h be elements of $(\text{the carrier of } Y)^X$. Then $h = (\text{FuncAdd}(X, Y))(f, g)$ if and only if for every element x of X holds $h(x) = f(x) + g(x)$.
- (4) For every element x of X holds $(\text{FuncZero}(X, Y))(x) = 0_Y$.

In the sequel a, b are real numbers.

We now state several propositions:

- (5) $h = (\text{FuncExtMult}(X, Y))(\langle a, f \rangle)$ iff for every element x of X holds $h(x) = a \cdot f(x)$.
- (6) $(\text{FuncAdd}(X, Y))(f, g) = (\text{FuncAdd}(X, Y))(g, f)$.
- (7) $(\text{FuncAdd}(X, Y))(f, (\text{FuncAdd}(X, Y))(g, h)) = (\text{FuncAdd}(X, Y))((\text{FuncAdd}(X, Y))(f, g), h)$.
- (8) $(\text{FuncAdd}(X, Y))(\text{FuncZero}(X, Y), f) = f$.
- (9) $(\text{FuncAdd}(X, Y))(f, (\text{FuncExtMult}(X, Y))(\langle -1, f \rangle)) = \text{FuncZero}(X, Y)$.
- (10) $(\text{FuncExtMult}(X, Y))(\langle 1, f \rangle) = f$.
- (11) $(\text{FuncExtMult}(X, Y))(\langle a, (\text{FuncExtMult}(X, Y))(\langle b, f \rangle) \rangle) = (\text{FuncExtMult}(X, Y))(\langle a \cdot b, f \rangle)$.
- (12) $(\text{FuncAdd}(X, Y))((\text{FuncExtMult}(X, Y))(\langle a, f \rangle), (\text{FuncExtMult}(X, Y))(\langle b, f \rangle)) = (\text{FuncExtMult}(X, Y))(\langle a + b, f \rangle)$.
- (13) $\langle (\text{the carrier of } Y)^X, \text{FuncZero}(X, Y), \text{FuncAdd}(X, Y), \text{FuncExtMult}(X, Y) \rangle$ is a real linear space.

Let X be a non empty set and let Y be a real linear space. The functor $\text{RealVectSpace}(X, Y)$ yielding a real linear space is defined by:

(Def. 4) $\text{RealVectSpace}(X, Y) = \langle (\text{the carrier of } Y)^X, \text{FuncZero}(X, Y), \text{FuncAdd}(X, Y), \text{FuncExtMult}(X, Y) \rangle$.

Let X be a non empty set and let Y be a real linear space. Note that $\text{RealVectSpace}(X, Y)$ is strict.

Next we state three propositions:

- (14) Let X be a non empty set, Y be a real linear space, f, g, h be vectors of $\text{RealVectSpace}(X, Y)$, and f', g', h' be elements of $(\text{the carrier of } Y)^X$. Suppose $f' = f$ and $g' = g$ and $h' = h$. Then $h = f + g$ if and only if for every element x of X holds $h'(x) = f'(x) + g'(x)$.
- (15) Let X be a non empty set, Y be a real linear space, f, h be vectors of $\text{RealVectSpace}(X, Y)$, and f', h' be elements of $(\text{the carrier of } Y)^X$. Suppose $f' = f$ and $h' = h$. Let a be a real number. Then $h = a \cdot f$ if and only if for every element x of X holds $h'(x) = a \cdot f'(x)$.
- (16) For every non empty set X and for every real linear space Y holds $0_{\text{RealVectSpace}(X, Y)} = X \mapsto 0_Y$.

2. REAL VECTOR SPACE OF LINEAR OPERATORS

Let X be a non empty RLS structure, let Y be a non empty loop structure, and let I_1 be a function from X into Y . We say that I_1 is additive if and only if:

(Def. 5) For all vectors x, y of X holds $I_1(x + y) = I_1(x) + I_1(y)$.

Let X, Y be non empty RLS structures and let I_1 be a function from X into Y . We say that I_1 is homogeneous if and only if:

(Def. 6) For every vector x of X and for every real number r holds $I_1(r \cdot x) = r \cdot I_1(x)$.

Let X be a non empty RLS structure and let Y be a real linear space. Note that there exists a function from X into Y which is additive and homogeneous.

Let X, Y be real linear spaces. A linear operator from X into Y is an additive homogeneous function from X into Y .

Let X, Y be real linear spaces. The functor $\text{LinearOperators}(X, Y)$ yields a subset of $\text{RealVectSpace}(\text{the carrier of } X, Y)$ and is defined by:

(Def. 7) For every set x holds $x \in \text{LinearOperators}(X, Y)$ iff x is a linear operator from X into Y .

Let X, Y be real linear spaces. One can check that $\text{LinearOperators}(X, Y)$ is non empty.

We now state two propositions:

(17) For all real linear spaces X, Y holds $\text{LinearOperators}(X, Y)$ is linearly closed.

(18) Let X, Y be real linear spaces. Then $\langle \text{LinearOperators}(X, Y), \text{Zero}_-(\text{LinearOperators}(X, Y), \text{RealVectSpace}(\text{the carrier of } X, Y)), \text{Add}_-(\text{LinearOperators}(X, Y), \text{RealVectSpace}(\text{the carrier of } X, Y)), \text{Mult}_-(\text{LinearOperators}(X, Y), \text{RealVectSpace}(\text{the carrier of } X, Y)) \rangle$ is a subspace of $\text{RealVectSpace}(\text{the carrier of } X, Y)$.

Let X, Y be real linear spaces. Note that $\langle \text{LinearOperators}(X, Y), \text{Zero}_-(\text{LinearOperators}(X, Y), \text{RealVectSpace}(\text{the carrier of } X, Y)), \text{Add}_-(\text{LinearOperators}(X, Y), \text{RealVectSpace}(\text{the carrier of } X, Y)), \text{Mult}_-(\text{LinearOperators}(X, Y), \text{RealVectSpace}(\text{the carrier of } X, Y)) \rangle$ is Abelian, add-associative, right zeroed, right complementable, and real linear space-like.

The following proposition is true

(19) Let X, Y be real linear spaces. Then $\langle \text{LinearOperators}(X, Y), \text{Zero}_-(\text{LinearOperators}(X, Y), \text{RealVectSpace}(\text{the carrier of } X, Y)), \text{Add}_-(\text{LinearOperators}(X, Y), \text{RealVectSpace}(\text{the carrier of } X, Y)), \text{Mult}_-(\text{LinearOperators}(X, Y), \text{RealVectSpace}(\text{the carrier of } X, Y)) \rangle$ is a real linear space.

Let X, Y be real linear spaces. The functor $\text{RVectorSpaceOfLinearOperators}(X, Y)$ yields a real linear space and is defined by:

(Def. 8) $\text{RVectorSpaceOfLinearOperators}(X, Y) = \langle \text{LinearOperators}(X, Y), \text{Zero}_-(\text{LinearOperators}(X, Y), \text{RealVectSpace}(\text{the carrier of } X, Y)), \text{Add}_-(\text{LinearOperators}(X, Y), \text{RealVectSpace}(\text{the carrier of } X, Y)), \text{Mult}_-(\text{LinearOperators}(X, Y), \text{RealVectSpace}(\text{the carrier of } X, Y)) \rangle$.

Let X, Y be real linear spaces. Note that $\text{RVectorSpaceOfLinearOperators}(X, Y)$ is strict.

One can prove the following four propositions:

(20) Let X, Y be real linear spaces, f, g, h be vectors of $\text{RVectorSpaceOfLinearOperators}(X, Y)$, and f', g', h' be linear operators from X into Y . Suppose $f' = f$ and $g' = g$ and $h' = h$. Then $h = f + g$ if and only if for every vector x of X holds $h'(x) = f'(x) + g'(x)$.

(21) Let X, Y be real linear spaces, f, h be vectors of $\text{RVectorSpaceOfLinearOperators}(X, Y)$, and f', h' be linear operators from X into Y . Suppose $f' = f$ and $h' = h$. Let a be a real number. Then $h = a \cdot f$ if and only if for every vector x of X holds $h'(x) = a \cdot f'(x)$.

(22) For all real linear spaces X, Y holds $0_{\text{RVectorSpaceOfLinearOperators}(X, Y)} = (\text{the carrier of } X) \longmapsto 0_Y$.

(23) For all real linear spaces X, Y holds $(\text{the carrier of } X) \longmapsto 0_Y$ is a linear operator from X into Y .

3. REAL NORMED LINEAR SPACE OF BOUNDED LINEAR OPERATORS

We now state the proposition

- (24) Let X be a real normed space, s_1 be a sequence of X , and g be a point of X . If s_1 is convergent and $\lim s_1 = g$, then $\|s_1\|$ is convergent and $\lim \|s_1\| = \|g\|$.

Let X, Y be real normed spaces and let I_1 be a linear operator from X into Y . We say that I_1 is bounded if and only if:

- (Def. 9) There exists a real number K such that $0 \leq K$ and for every vector x of X holds $\|I_1(x)\| \leq K \cdot \|x\|$.

We now state the proposition

- (25) Let X, Y be real normed spaces and f be a linear operator from X into Y . If for every vector x of X holds $f(x) = 0_Y$, then f is bounded.

Let X, Y be real normed spaces. Note that there exists a linear operator from X into Y which is bounded.

Let X, Y be real normed spaces. The functor $\text{BoundedLinearOperators}(X, Y)$ yields a subset of $\text{RVectorSpaceOfLinearOperators}(X, Y)$ and is defined as follows:

- (Def. 10) For every set x holds $x \in \text{BoundedLinearOperators}(X, Y)$ iff x is a bounded linear operator from X into Y .

Let X, Y be real normed spaces. Note that $\text{BoundedLinearOperators}(X, Y)$ is non empty. One can prove the following propositions:

- (26) For all real normed spaces X, Y holds $\text{BoundedLinearOperators}(X, Y)$ is linearly closed.

- (27) For all real normed spaces X, Y holds $\langle \text{BoundedLinearOperators}(X, Y), \text{Zero}(\text{BoundedLinearOperators}(X, Y)), \text{RVectorSpaceOfLinearOperators}(X, Y) \rangle$ is a subspace of $\text{RVectorSpaceOfLinearOperators}(X, Y)$.

Let X, Y be real normed spaces. Note that $\langle \text{BoundedLinearOperators}(X, Y), \text{Zero}(\text{BoundedLinearOperators}(X, Y)), \text{RVectorSpaceOfLinearOperators}(X, Y) \rangle$ is Abelian, add-associative, right zeroed, right complementable, and real linear space-like.

Next we state the proposition

- (28) For all real normed spaces X, Y holds $\langle \text{BoundedLinearOperators}(X, Y), \text{Zero}(\text{BoundedLinearOperators}(X, Y)), \text{RVectorSpaceOfLinearOperators}(X, Y) \rangle$ is a real linear space.

Let X, Y be real normed spaces. The functor $\text{RVectorSpaceOfBoundedLinearOperators}(X, Y)$ yielding a real linear space is defined as follows:

- (Def. 11) $\text{RVectorSpaceOfBoundedLinearOperators}(X, Y) = \langle \text{BoundedLinearOperators}(X, Y), \text{Zero}(\text{BoundedLinearOperators}(X, Y)), \text{RVectorSpaceOfLinearOperators}(X, Y) \rangle$

Let X, Y be real normed spaces. Observe that $\text{RVectorSpaceOfBoundedLinearOperators}(X, Y)$ is strict.

One can prove the following three propositions:

- (29) Let X, Y be real normed spaces, f, g, h be vectors of $\text{RVectorSpaceOfBoundedLinearOperators}(X, Y)$, and f', g', h' be bounded linear operators from X into Y . Suppose $f' = f$ and $g' = g$ and $h' = h$. Then $h = f + g$ if and only if for every vector x of X holds $h'(x) = f'(x) + g'(x)$.
- (30) Let X, Y be real normed spaces, f, h be vectors of $\text{RVectorSpaceOfBoundedLinearOperators}(X, Y)$, and f', h' be bounded linear operators from X into Y . Suppose $f' = f$ and $h' = h$. Let a be a real number. Then $h = a \cdot f$ if and only if for every vector x of X holds $h'(x) = a \cdot f'(x)$.
- (31) For all real normed spaces X, Y holds $0_{\text{RVectorSpaceOfBoundedLinearOperators}(X, Y)} = (\text{the carrier of } X) \mapsto 0_Y$.

Let X, Y be real normed spaces and let f be a set. Let us assume that $f \in \text{BoundedLinearOperators}(X, Y)$. The functor $\text{modetrans}(f, X, Y)$ yields a bounded linear operator from X into Y and is defined as follows:

(Def. 12) $\text{modetrans}(f, X, Y) = f$.

Let X, Y be real normed spaces and let u be a linear operator from X into Y . The functor $\text{PreNorms}(u)$ yields a non empty subset of \mathbb{R} and is defined by:

(Def. 13) $\text{PreNorms}(u) = \{\|u(t)\|; t \text{ ranges over vectors of } X: \|t\| \leq 1\}$.

Next we state three propositions:

(32) Let X, Y be real normed spaces and g be a bounded linear operator from X into Y . Then $\text{PreNorms}(g)$ is non empty and upper bounded.

(33) Let X, Y be real normed spaces and g be a linear operator from X into Y . Then g is bounded if and only if $\text{PreNorms}(g)$ is upper bounded.

(34) Let X, Y be real normed spaces. Then there exists a function N_1 from $\text{BoundedLinearOperators}(X, Y)$ into \mathbb{R} such that for every set f if $f \in \text{BoundedLinearOperators}(X, Y)$, then $N_1(f) = \sup \text{PreNorms}(\text{modetrans}(f, X, Y))$.

Let X, Y be real normed spaces. The functor $\text{BoundedLinearOperatorsNorm}(X, Y)$ yields a function from $\text{BoundedLinearOperators}(X, Y)$ into \mathbb{R} and is defined by:

(Def. 14) For every set x such that $x \in \text{BoundedLinearOperators}(X, Y)$ holds $(\text{BoundedLinearOperatorsNorm}(X, Y))(x) = \sup \text{PreNorms}(\text{modetrans}(x, X, Y))$.

One can prove the following propositions:

(35) For all real normed spaces X, Y and for every bounded linear operator f from X into Y holds $\text{modetrans}(f, X, Y) = f$.

(36) For all real normed spaces X, Y and for every bounded linear operator f from X into Y holds $(\text{BoundedLinearOperatorsNorm}(X, Y))(f) = \sup \text{PreNorms}(f)$.

Let X, Y be real normed spaces. The functor $\text{RNormSpaceOfBoundedLinearOperators}(X, Y)$ yielding a non empty normed structure is defined as follows:

(Def. 15) $\text{RNormSpaceOfBoundedLinearOperators}(X, Y) = \langle \text{BoundedLinearOperators}(X, Y), \text{Zero}(\text{BoundedLinearOperators}(X, Y)) \rangle$

We now state several propositions:

(37) For all real normed spaces X, Y holds $(\text{the carrier of } X) \mapsto 0_Y = 0_{\text{RNormSpaceOfBoundedLinearOperators}(X, Y)}$.

(38) Let X, Y be real normed spaces, f be a point of $\text{RNormSpaceOfBoundedLinearOperators}(X, Y)$, and g be a bounded linear operator from X into Y . If $g = f$, then for every vector t of X holds $\|g(t)\| \leq \|f\| \cdot \|t\|$.

(39) For all real normed spaces X, Y and for every point f of $\text{RNormSpaceOfBoundedLinearOperators}(X, Y)$ holds $0 \leq \|f\|$.

(40) For all real normed spaces X, Y and for every point f of $\text{RNormSpaceOfBoundedLinearOperators}(X, Y)$ such that $f = 0_{\text{RNormSpaceOfBoundedLinearOperators}(X, Y)}$ holds $0 = \|f\|$.

(41) Let X, Y be real normed spaces, f, g, h be points of $\text{RNormSpaceOfBoundedLinearOperators}(X, Y)$, and f', g', h' be bounded linear operators from X into Y . Suppose $f' = f$ and $g' = g$ and $h' = h$. Then $h = f + g$ if and only if for every vector x of X holds $h'(x) = f'(x) + g'(x)$.

(42) Let X, Y be real normed spaces, f, h be points of $\text{RNormSpaceOfBoundedLinearOperators}(X, Y)$, and f', h' be bounded linear operators from X into Y . Suppose $f' = f$ and $h' = h$. Let a be a real number. Then $h = a \cdot f$ if and only if for every vector x of X holds $h'(x) = a \cdot f'(x)$.

- (43) Let X be a real normed space, Y be a real normed space, f, g be points of $\text{RNormSpaceOfBoundedLinearOperators}(X, Y)$, and a be a real number. Then $\|f\| = 0$ iff $f = 0_{\text{RNormSpaceOfBoundedLinearOperators}(X, Y)}$ and $\|a \cdot f\| = |a| \cdot \|f\|$ and $\|f + g\| \leq \|f\| + \|g\|$.
- (44) For all real normed spaces X, Y holds $\text{RNormSpaceOfBoundedLinearOperators}(X, Y)$ is real normed space-like.
- (45) For all real normed spaces X, Y holds $\text{RNormSpaceOfBoundedLinearOperators}(X, Y)$ is a real normed space.

Let X, Y be real normed spaces. Observe that $\text{RNormSpaceOfBoundedLinearOperators}(X, Y)$ is real normed space-like, real linear space-like, Abelian, add-associative, right zeroed, and right complementable.

The following proposition is true

- (46) Let X, Y be real normed spaces, f, g, h be points of $\text{RNormSpaceOfBoundedLinearOperators}(X, Y)$, and f', g', h' be bounded linear operators from X into Y . Suppose $f' = f$ and $g' = g$ and $h' = h$. Then $h = f - g$ if and only if for every vector x of X holds $h'(x) = f'(x) - g'(x)$.

4. REAL BANACH SPACE OF BOUNDED LINEAR OPERATORS

Let X be a real normed space. We say that X is complete if and only if:

- (Def. 16) For every sequence s_1 of X such that s_1 is Cauchy sequence by norm holds s_1 is convergent.

One can verify that there exists a real normed space which is complete.

A real Banach space is a complete real normed space.

Next we state three propositions:

- (47) Let X be a real normed space and s_1 be a sequence of X . If s_1 is convergent, then $\|s_1\|$ is convergent and $\lim \|s_1\| = \|\lim s_1\|$.
- (48) Let X, Y be real normed spaces. Suppose Y is complete. Let s_1 be a sequence of $\text{RNormSpaceOfBoundedLinearOperators}(X, Y)$. If s_1 is Cauchy sequence by norm, then s_1 is convergent.
- (49) For every real normed space X and for every real Banach space Y holds $\text{RNormSpaceOfBoundedLinearOperators}(X, Y)$ is a real Banach space.

Let X be a real normed space and let Y be a real Banach space. Observe that $\text{RNormSpaceOfBoundedLinearOperators}(X, Y)$ is complete.

REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinal1.html>.
- [2] Grzegorz Bancerek. Sequences of ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinal2.html>.
- [3] Czesław Byliński. Binary operations. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/binop_1.html.
- [4] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [5] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [6] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [7] Czesław Byliński and Piotr Rudnicki. Bounding boxes for compact sets in E^2 . *Journal of Formalized Mathematics*, 9, 1997. http://mizar.org/JFM/Vol9/pscomp_1.html.
- [8] Noboru Endou, Yasumasa Suzuki, and Yasunari Shidama. Real linear space of real sequences. *Journal of Formalized Mathematics*, 15, 2003. <http://mizar.org/JFM/Vol15/rsspace.html>.

- [9] Krzysztof Hryniewiecki. Basic properties of real numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/real_1.html.
- [10] Jarosław Kotowicz. Convergent real sequences. Upper and lower bound of sets of real numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/seq_4.html.
- [11] Jarosław Kotowicz. Convergent sequences and the limit of sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/seq_2.html.
- [12] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/seqm_3.html.
- [13] Jarosław Kotowicz. Real sequences and basic operations on them. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/seq_1.html.
- [14] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/vectsp_1.html.
- [15] Henryk Orszyszczyn and Krzysztof Prażmowski. Real functions spaces. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol2/funcsdom.html>.
- [16] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/pre_topc.html.
- [17] Jan Popiołek. Some properties of functions modul and signum. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/absvalue.html>.
- [18] Jan Popiołek. Real normed space. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/normsp_1.html.
- [19] Jan Popiołek. Introduction to Banach and Hilbert spaces — part I. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/bhsp_1.html.
- [20] Yasunasa Suzuki, Noboru Endou, and Yasunari Shidama. Banach space of absolute summable real sequences. *Journal of Formalized Mathematics*, 15, 2003. <http://mizar.org/JFM/Vol15/rsspace3.html>.
- [21] Andrzej Trybulec. Binary operations applied to functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funcop_1.html.
- [22] Andrzej Trybulec. Domains and their Cartesian products. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/domain_1.html.
- [23] Andrzej Trybulec. Enumerated sets. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/enumset1.html>.
- [24] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [25] Andrzej Trybulec. Tuples, projections and Cartesian products. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/mcart_1.html.
- [26] Andrzej Trybulec. Function domains and Fränkel operator. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol2/raenkel.html>.
- [27] Andrzej Trybulec. On the sets inhabited by numbers. *Journal of Formalized Mathematics*, 15, 2003. <http://mizar.org/JFM/Vol16/membered.html>.
- [28] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [29] Wojciech A. Trybulec. Subspaces and cosets of subspaces in real linear space. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/rbsub_1.html.
- [30] Wojciech A. Trybulec. Vectors in real linear space. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/rlvect_1.html.
- [31] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [32] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.
- [33] Edmund Woronowicz. Relations defined on sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relset_1.html.

Received December 22, 2003

Published December 22, 2003
