

# The Product and the Determinant of Matrices with Entries in a Field

Katarzyna Zawadzka  
Warsaw University  
Białystok

**Summary.** Concerned with a generalization of concepts introduced in [14], i.e. there are introduced the sum and the product of matrices of any dimension of elements of any field.

MML Identifier: MATRIX\_3.

WWW: [http://mizar.org/JFM/Vol5/matrix\\_3.html](http://mizar.org/JFM/Vol5/matrix_3.html)

The articles [19], [8], [24], [20], [13], [25], [6], [7], [3], [5], [4], [17], [23], [16], [18], [11], [10], [9], [14], [22], [15], [1], [21], [26], [2], and [12] provide the notation and terminology for this paper.

## 1. AUXILIARY THEOREMS

We adopt the following rules:  $i, j, k, l, n, m$  are natural numbers,  $D$  is a non empty set, and  $K$  is a field.

The following proposition is true

- (1) If  $n = n + k$ , then  $k = 0$ .

Let us consider  $K, n, m$ . The functor  $\left( \begin{array}{ccc} 0 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & 0 \end{array} \right)_{n \times m}^K$  yields a matrix over  $K$  of dimension  $n \times m$  and is defined by:

$$\text{(Def. 1)} \quad \left( \begin{array}{ccc} 0 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & 0 \end{array} \right)_{n \times m}^K = n \mapsto (m \mapsto 0_K).$$

Let us consider  $K$  and let  $A$  be a matrix over  $K$ . The functor  $-A$  yields a matrix over  $K$  and is defined by:

$$\text{(Def. 2)} \quad \text{len}(-A) = \text{len}A \text{ and } \text{width}(-A) = \text{width}A \text{ and for all } i, j \text{ such that } \langle i, j \rangle \in \text{the indices of } A \text{ holds } (-A) \circ (i, j) = -(A \circ (i, j)).$$

Let us consider  $K$  and let  $A, B$  be matrices over  $K$ . The functor  $A + B$  yields a matrix over  $K$  and is defined by:

$$\text{(Def. 3)} \quad \text{len}(A + B) = \text{len}A \text{ and } \text{width}(A + B) = \text{width}A \text{ and for all } i, j \text{ such that } \langle i, j \rangle \in \text{the indices of } A \text{ holds } (A + B) \circ (i, j) = (A \circ (i, j)) + (B \circ (i, j)).$$

The following propositions are true:

(3)<sup>1</sup> For all  $i, j$  such that  $\langle i, j \rangle \in$  the indices of  $\begin{pmatrix} 0 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & 0 \end{pmatrix}_K^{n \times m}$  holds  $\begin{pmatrix} 0 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & 0 \end{pmatrix}_K^{n \times m} \circ (i, j) = 0_K$ .

(4) For all matrices  $A, B$  over  $K$  such that  $\text{len}A = \text{len}B$  and  $\text{width}A = \text{width}B$  holds  $A + B = B + A$ .

(5) For all matrices  $A, B, C$  over  $K$  such that  $\text{len}A = \text{len}B$  and  $\text{len}A = \text{len}C$  and  $\text{width}A = \text{width}B$  and  $\text{width}A = \text{width}C$  holds  $(A + B) + C = A + (B + C)$ .

(6) For every matrix  $A$  over  $K$  of dimension  $n \times m$  holds  $A + \begin{pmatrix} 0 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & 0 \end{pmatrix}_K^{n \times m} = A$ .

(7) For every matrix  $A$  over  $K$  of dimension  $n \times m$  holds  $A + -A = \begin{pmatrix} 0 & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & 0 \end{pmatrix}_K^{n \times m}$ .

Let us consider  $K$  and let  $A, B$  be matrices over  $K$ . Let us assume that  $\text{width}A = \text{len}B$ . The functor  $A \cdot B$  yielding a matrix over  $K$  is defined by:

(Def. 4)  $\text{len}(A \cdot B) = \text{len}A$  and  $\text{width}(A \cdot B) = \text{width}B$  and for all  $i, j$  such that  $\langle i, j \rangle \in$  the indices of  $A \cdot B$  holds  $(A \cdot B) \circ (i, j) = \text{Line}(A, i) \cdot B_{\square, j}$ .

Let us consider  $n, k, m$ , let us consider  $K$ , let  $A$  be a matrix over  $K$  of dimension  $n \times k$ , and let  $B$  be a matrix over  $K$  of dimension  $\text{width}A \times m$ . Then  $A \cdot B$  is a matrix over  $K$  of dimension  $\text{len}A \times \text{width}B$ .

Let us consider  $K$ , let  $M$  be a matrix over  $K$ , and let  $a$  be an element of  $K$ . The functor  $a \cdot M$  yields a matrix over  $K$  and is defined as follows:

(Def. 5)  $\text{len}(a \cdot M) = \text{len}M$  and  $\text{width}(a \cdot M) = \text{width}M$  and for all  $i, j$  such that  $\langle i, j \rangle \in$  the indices of  $M$  holds  $(a \cdot M) \circ (i, j) = a \cdot (M \circ (i, j))$ .

Let us consider  $K$ , let  $M$  be a matrix over  $K$ , and let  $a$  be an element of  $K$ . The functor  $M \cdot a$  yielding a matrix over  $K$  is defined as follows:

(Def. 6)  $M \cdot a = a \cdot M$ .

One can prove the following propositions:

(8) For all finite sequences  $p, q$  of elements of the carrier of  $K$  such that  $\text{len}p = \text{len}q$  holds  $\text{len}(p \bullet q) = \text{len}p$  and  $\text{len}(p \bullet q) = \text{len}q$ .

(9) For all  $i, l$  such that  $\langle i, l \rangle \in$  the indices of  $\begin{pmatrix} 1 & & 0 \\ & \ddots & \\ 0 & & 1 \end{pmatrix}_K^{n \times n}$  and  $l = i$  holds

$$\text{Line}\left(\begin{pmatrix} 1 & & 0 \\ & \ddots & \\ 0 & & 1 \end{pmatrix}_K^{n \times n}, i\right)(l) = \mathbf{1}_K.$$

(10) For all  $i, l$  such that  $\langle i, l \rangle \in$  the indices of  $\begin{pmatrix} 1 & & 0 \\ & \ddots & \\ 0 & & 1 \end{pmatrix}_K^{n \times n}$  and  $l \neq i$  holds

$$\text{Line}\left(\begin{pmatrix} 1 & & 0 \\ & \ddots & \\ 0 & & 1 \end{pmatrix}_K^{n \times n}, i\right)(l) = 0_K.$$

<sup>1</sup> The proposition (2) has been removed.

(11) For all  $l, j$  such that  $\langle l, j \rangle \in$  the indices of  $\begin{pmatrix} 1 & 0 \\ & \ddots \\ 0 & 1 \end{pmatrix}_K^{n \times n}$  and  $l = j$  holds

$$\left( \begin{pmatrix} 1 & 0 \\ & \ddots \\ 0 & 1 \end{pmatrix}_K^{n \times n} \right)_{\square, j}(l) = \mathbf{1}_K.$$

(12) For all  $l, j$  such that  $\langle l, j \rangle \in$  the indices of  $\begin{pmatrix} 1 & 0 \\ & \ddots \\ 0 & 1 \end{pmatrix}_K^{n \times n}$  and  $l \neq j$  holds

$$\left( \begin{pmatrix} 1 & 0 \\ & \ddots \\ 0 & 1 \end{pmatrix}_K^{n \times n} \right)_{\square, j}(l) = 0_K.$$

(13) For every add-associative right zeroed right complementable non empty loop structure  $K$  holds  $\Sigma(n \mapsto 0_K) = 0_K$ .

(14) Let  $K$  be an add-associative right zeroed right complementable non empty loop structure,  $p$  be a finite sequence of elements of the carrier of  $K$ , and given  $i$ . If  $i \in \text{dom } p$  and for every  $k$  such that  $k \in \text{dom } p$  and  $k \neq i$  holds  $p(k) = 0_K$ , then  $\Sigma p = p(i)$ .

(15) For all finite sequences  $p, q$  of elements of the carrier of  $K$  holds  $\text{len}(p \bullet q) = \min(\text{len } p, \text{len } q)$ .

(16) Let  $p, q$  be finite sequences of elements of the carrier of  $K$  and given  $i$ . Suppose  $i \in \text{dom } p$  and  $p(i) = \mathbf{1}_K$  and for every  $k$  such that  $k \in \text{dom } p$  and  $k \neq i$  holds  $p(k) = 0_K$ . Let given  $j$  such that  $j \in \text{dom}(p \bullet q)$ . Then

- (i) if  $i = j$ , then  $(p \bullet q)(j) = q(i)$ , and
- (ii) if  $i \neq j$ , then  $(p \bullet q)(j) = 0_K$ .

(17) For all  $i, j$  such that  $\langle i, j \rangle \in$  the indices of  $\begin{pmatrix} 1 & 0 \\ & \ddots \\ 0 & 1 \end{pmatrix}_K^{n \times n}$  holds if  $i = j$ , then

$$\text{Line} \left( \begin{pmatrix} 1 & 0 \\ & \ddots \\ 0 & 1 \end{pmatrix}_K^{n \times n}, i \right)(j) = \mathbf{1}_K \text{ and if } i \neq j, \text{ then } \text{Line} \left( \begin{pmatrix} 1 & 0 \\ & \ddots \\ 0 & 1 \end{pmatrix}_K^{n \times n}, i \right)(j) = 0_K.$$

(18) For all  $i, j$  such that  $\langle i, j \rangle \in$  the indices of  $\begin{pmatrix} 1 & 0 \\ & \ddots \\ 0 & 1 \end{pmatrix}_K^{n \times n}$  holds if  $i = j$ , then

$$\left( \begin{pmatrix} 1 & 0 \\ & \ddots \\ 0 & 1 \end{pmatrix}_K^{n \times n} \right)_{\square, j}(i) = \mathbf{1}_K \text{ and if } i \neq j, \text{ then } \left( \begin{pmatrix} 1 & 0 \\ & \ddots \\ 0 & 1 \end{pmatrix}_K^{n \times n} \right)_{\square, j}(i) = 0_K.$$

(19) Let  $p, q$  be finite sequences of elements of the carrier of  $K$  and given  $i$ . Suppose  $i \in \text{dom } p$  and  $i \in \text{dom } q$  and  $p(i) = \mathbf{1}_K$  and for every  $k$  such that  $k \in \text{dom } p$  and  $k \neq i$  holds  $p(k) = 0_K$ . Then  $\Sigma(p \bullet q) = q(i)$ .

(20) For every matrix  $A$  over  $K$  of dimension  $n$  holds  $\begin{pmatrix} 1 & 0 \\ & \ddots \\ 0 & 1 \end{pmatrix}_K^{n \times n} \cdot A = A$ .

- (21) For every matrix  $A$  over  $K$  of dimension  $n$  holds  $A \cdot \begin{pmatrix} 1 & & 0 \\ & \ddots & \\ 0 & & 1 \end{pmatrix}_K^{n \times n} = A$ .
- (22) For all elements  $a, b$  of  $K$  holds  $\langle\langle a \rangle\rangle \cdot \langle\langle b \rangle\rangle = \langle\langle a \cdot b \rangle\rangle$ .
- (23) For all elements  $a_1, a_2, b_1, b_2, c_1, c_2, d_1, d_2$  of  $K$  holds  $\begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix} \cdot \begin{pmatrix} a_2 & b_2 \\ c_2 & d_2 \end{pmatrix} = \begin{pmatrix} a_1 \cdot a_2 + b_1 \cdot c_2 & a_1 \cdot b_2 + b_1 \cdot d_2 \\ c_1 \cdot a_2 + d_1 \cdot c_2 & c_1 \cdot b_2 + d_1 \cdot d_2 \end{pmatrix}$ .
- (24) For all matrices  $A, B$  over  $K$  such that  $\text{width}A = \text{len}B$  and  $\text{width}B \neq 0$  holds  $(A \cdot B)^T = B^T \cdot A^T$ .

## 2. THE PRODUCT OF MATRICES

Let  $I, J$  be non empty sets, let  $X$  be an element of  $\text{Fin}I$ , and let  $Y$  be an element of  $\text{Fin}J$ . Then  $[X, Y]$  is an element of  $\text{Fin}[I, J]$ .

Let  $I, J, D$  be non empty sets, let  $G$  be a binary operation on  $D$ , let  $f$  be a function from  $I$  into  $D$ , and let  $g$  be a function from  $J$  into  $D$ . Then  $G \circ (f, g)$  is a function from  $[I, J]$  into  $D$ .

We now state a number of propositions:

- (25) Let  $I, J, D$  be non empty sets,  $F, G$  be binary operations on  $D$ ,  $f$  be a function from  $I$  into  $D$ ,  $g$  be a function from  $J$  into  $D$ ,  $X$  be an element of  $\text{Fin}I$ , and  $Y$  be an element of  $\text{Fin}J$ . Suppose  $F$  is commutative and associative but  $[Y, X] \neq \emptyset$  or  $F$  has a unity but  $G$  is commutative. Then  $F\text{-}\sum_{[X, Y]}(G \circ (f, g)) = F\text{-}\sum_{[Y, X]}(G \circ (g, f))$ .
- (26) Let  $I, J$  be non empty sets,  $F, G$  be binary operations on  $D$ ,  $f$  be a function from  $I$  into  $D$ , and  $g$  be a function from  $J$  into  $D$ . Suppose  $F$  is commutative and associative and has a unity. Let  $x$  be an element of  $I$  and  $y$  be an element of  $J$ . Then  $F\text{-}\sum_{[\{x\}, \{y\}]}(G \circ (f, g)) = F\text{-}\sum_{\{x\}} G^\circ(f, F\text{-}\sum_{\{y\}} g)$ .
- (27) Let  $I, J$  be non empty sets,  $F, G$  be binary operations on  $D$ ,  $f$  be a function from  $I$  into  $D$ ,  $g$  be a function from  $J$  into  $D$ ,  $X$  be an element of  $\text{Fin}I$ , and  $Y$  be an element of  $\text{Fin}J$ . Suppose  $F$  is commutative and associative and has a unity and an inverse operation and  $G$  is distributive w.r.t.  $F$ . Let  $x$  be an element of  $I$ . Then  $F\text{-}\sum_{[\{x\}, Y]}(G \circ (f, g)) = F\text{-}\sum_{\{x\}} G^\circ(f, F\text{-}\sum_Y g)$ .
- (28) Let  $I, J$  be non empty sets,  $F, G$  be binary operations on  $D$ ,  $f$  be a function from  $I$  into  $D$ ,  $g$  be a function from  $J$  into  $D$ ,  $X$  be an element of  $\text{Fin}I$ , and  $Y$  be an element of  $\text{Fin}J$ . Suppose  $F$  is commutative and associative and has a unity and an inverse operation and  $G$  is distributive w.r.t.  $F$ . Then  $F\text{-}\sum_{[X, Y]}(G \circ (f, g)) = F\text{-}\sum_X G^\circ(f, F\text{-}\sum_Y g)$ .
- (29) Let  $I, J$  be non empty sets,  $F, G$  be binary operations on  $D$ ,  $f$  be a function from  $I$  into  $D$ , and  $g$  be a function from  $J$  into  $D$ . Suppose  $F$  is commutative and associative and has a unity and  $G$  is commutative. Let  $x$  be an element of  $I$  and  $y$  be an element of  $J$ . Then  $F\text{-}\sum_{[\{x\}, \{y\}]}(G \circ (f, g)) = F\text{-}\sum_{\{y\}} G^\circ(F\text{-}\sum_{\{x\}} f, g)$ .
- (30) Let  $I, J$  be non empty sets,  $F, G$  be binary operations on  $D$ ,  $f$  be a function from  $I$  into  $D$ ,  $g$  be a function from  $J$  into  $D$ ,  $X$  be an element of  $\text{Fin}I$ , and  $Y$  be an element of  $\text{Fin}J$ . Suppose that
- (i)  $F$  is commutative and associative and has a unity and an inverse operation, and
  - (ii)  $G$  is distributive w.r.t.  $F$  and commutative.
- Then  $F\text{-}\sum_{[X, Y]}(G \circ (f, g)) = F\text{-}\sum_Y G^\circ(F\text{-}\sum_X f, g)$ .
- (31) Let  $I, J$  be non empty sets,  $F$  be a binary operation on  $D$ ,  $f$  be a function from  $[I, J]$  into  $D$ ,  $g$  be a function from  $I$  into  $D$ , and  $Y$  be an element of  $\text{Fin}J$ . Suppose  $F$  is commutative and associative and has a unity and an inverse operation. Let  $x$  be an element of  $I$ . If for every element  $i$  of  $I$  holds  $g(i) = F\text{-}\sum_Y(\text{curry } f)(i)$ , then  $F\text{-}\sum_{[\{x\}, Y]} f = F\text{-}\sum_{\{x\}} g$ .

- (32) Let  $I, J$  be non empty sets,  $F$  be a binary operation on  $D$ ,  $f$  be a function from  $[I, J]$  into  $D$ ,  $g$  be a function from  $I$  into  $D$ ,  $X$  be an element of  $\text{Fin}I$ , and  $Y$  be an element of  $\text{Fin}J$ . Suppose for every element  $i$  of  $I$  holds  $g(i) = F\text{-}\sum_Y(\text{curry } f)(i)$  and  $F$  is commutative and associative and has a unity and an inverse operation. Then  $F\text{-}\sum_{[X, Y]} f = F\text{-}\sum_X g$ .
- (33) Let  $I, J$  be non empty sets,  $F$  be a binary operation on  $D$ ,  $f$  be a function from  $[I, J]$  into  $D$ ,  $g$  be a function from  $J$  into  $D$ , and  $X$  be an element of  $\text{Fin}I$ . Suppose  $F$  is commutative and associative and has a unity and an inverse operation. Let  $y$  be an element of  $J$ . If for every element  $j$  of  $J$  holds  $g(j) = F\text{-}\sum_X(\text{curry}' f)(j)$ , then  $F\text{-}\sum_{[X, \{y\}]} f = F\text{-}\sum_{\{y\}} g$ .
- (34) Let  $I, J$  be non empty sets,  $F$  be a binary operation on  $D$ ,  $f$  be a function from  $[I, J]$  into  $D$ ,  $g$  be a function from  $J$  into  $D$ ,  $X$  be an element of  $\text{Fin}I$ , and  $Y$  be an element of  $\text{Fin}J$ . Suppose for every element  $j$  of  $J$  holds  $g(j) = F\text{-}\sum_X(\text{curry}' f)(j)$  and  $F$  is commutative and associative and has a unity and an inverse operation. Then  $F\text{-}\sum_{[X, Y]} f = F\text{-}\sum_Y g$ .
- (35) For all matrices  $A, B, C$  over  $K$  such that  $\text{width}A = \text{len}B$  and  $\text{width}B = \text{len}C$  holds  $(A \cdot B) \cdot C = A \cdot (B \cdot C)$ .

### 3. DETERMINANT

Let us consider  $n, K$ , let  $M$  be a matrix over  $K$  of dimension  $n$ , and let  $p$  be an element of the permutations of  $n$ -element set. The functor  $p\text{-Path}M$  yielding a finite sequence of elements of the carrier of  $K$  is defined by:

- (Def. 7)  $\text{len}(p\text{-Path}M) = n$  and for all  $i, j$  such that  $i \in \text{dom}(p\text{-Path}M)$  and  $j = p(i)$  holds  $(p\text{-Path}M)(i) = M \circ (i, j)$ .

Let us consider  $n, K$  and let  $M$  be a matrix over  $K$  of dimension  $n$ . The product on paths of  $M$  yields a function from the permutations of  $n$ -element set into the carrier of  $K$  and is defined by the condition (Def. 8).

- (Def. 8) Let  $p$  be an element of the permutations of  $n$ -element set. Then (the product on paths of  $M$ )( $p$ ) =  $(-1)^{\text{sgn}(p)}$ (the multiplication of  $K \otimes (p\text{-Path}M)$ ).

Let us consider  $n$ , let us consider  $K$ , and let  $M$  be a matrix over  $K$  of dimension  $n$ . The functor  $\text{Det}M$  yielding an element of the carrier of  $K$  is defined as follows:

- (Def. 9)  $\text{Det}M = (\text{the addition of } K)\text{-}\sum_{\Omega_{\text{the permutations of } n\text{-element set}}^f}$  (the product on paths of  $M$ ).

In the sequel  $a$  denotes an element of  $K$ .

Next we state the proposition

- (36)  $\text{Det}\langle\langle a \rangle\rangle = a$ .

Let us consider  $n$ , let us consider  $K$ , and let  $M$  be a matrix over  $K$  of dimension  $n$ . The diagonal of  $M$  yields a finite sequence of elements of the carrier of  $K$  and is defined by:

- (Def. 10)  $\text{len}(\text{the diagonal of } M) = n$  and for every  $i$  such that  $i \in \text{Seg } n$  holds  $(\text{the diagonal of } M)(i) = M \circ (i, i)$ .

### REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/nat\\_1.html](http://mizar.org/JFM/Vol1/nat_1.html).
- [2] Grzegorz Bancerek. Curried and uncurried functions. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/funct\\_5.html](http://mizar.org/JFM/Vol2/funct_5.html).
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/finseq\\_1.html](http://mizar.org/JFM/Vol1/finseq_1.html).
- [4] Czesław Byliński. Basic functions and operations on functions. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/funct\\_3.html](http://mizar.org/JFM/Vol1/funct_3.html).

- [5] Czesław Byliński. Binary operations. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/binop\\_1.html](http://mizar.org/JFM/Vol1/binop_1.html).
- [6] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/funct\\_1.html](http://mizar.org/JFM/Vol1/funct_1.html).
- [7] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/funct\\_2.html](http://mizar.org/JFM/Vol1/funct_2.html).
- [8] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/zfmisc\\_1.html](http://mizar.org/JFM/Vol1/zfmisc_1.html).
- [9] Czesław Byliński. Binary operations applied to finite sequences. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol2/finseqop.html>.
- [10] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/finseq\\_2.html](http://mizar.org/JFM/Vol2/finseq_2.html).
- [11] Czesław Byliński. Semigroup operations on finite subsets. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/setwop\\_2.html](http://mizar.org/JFM/Vol2/setwop_2.html).
- [12] Czesław Byliński. Subcategories and products of categories. *Journal of Formalized Mathematics*, 2, 1990. [http://mizar.org/JFM/Vol2/cat\\_2.html](http://mizar.org/JFM/Vol2/cat_2.html).
- [13] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/finset\\_1.html](http://mizar.org/JFM/Vol1/finset_1.html).
- [14] Katarzyna Jankowska. Matrices. Abelian group of matrices. *Journal of Formalized Mathematics*, 3, 1991. [http://mizar.org/JFM/Vol3/matrix\\_1.html](http://mizar.org/JFM/Vol3/matrix_1.html).
- [15] Katarzyna Jankowska. Transpose matrices and groups of permutations. *Journal of Formalized Mathematics*, 4, 1992. [http://mizar.org/JFM/Vol4/matrix\\_2.html](http://mizar.org/JFM/Vol4/matrix_2.html).
- [16] Eugeniusz Kusak, Wojciech Leoniczuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/vectsp\\_1.html](http://mizar.org/JFM/Vol1/vectsp_1.html).
- [17] Andrzej Trybulec. Binary operations applied to functions. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/funcop\\_1.html](http://mizar.org/JFM/Vol1/funcop_1.html).
- [18] Andrzej Trybulec. Semilattice operations on finite subsets. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/setwiseo.html>.
- [19] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [20] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [21] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers operations: min, max, square, and square root. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/square\\_1.html](http://mizar.org/JFM/Vol1/square_1.html).
- [22] Andrzej Trybulec and Agata Darmochwał. Boolean domains. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/finsub\\_1.html](http://mizar.org/JFM/Vol1/finsub_1.html).
- [23] Wojciech A. Trybulec. Vectors in real linear space. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/rlvect\\_1.html](http://mizar.org/JFM/Vol1/rlvect_1.html).
- [24] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/subset\\_1.html](http://mizar.org/JFM/Vol1/subset_1.html).
- [25] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/relat\\_1.html](http://mizar.org/JFM/Vol1/relat_1.html).
- [26] Katarzyna Zawadzka. Sum and product of finite sequences of elements of a field. *Journal of Formalized Mathematics*, 4, 1992. [http://mizar.org/JFM/Vol4/fvsum\\_1.html](http://mizar.org/JFM/Vol4/fvsum_1.html).

*Received May 17, 1993*

*Published January 2, 2004*

---