

Reper Algebras

Michał Muzalewski
Warsaw University
Białystok

Summary. We shall describe n -dimensional spaces with the reper operation [7, pages 72–79]. An inspiration to such approach comes from the monograph [9] and so-called Leibniz program. Let us recall that the Leibniz program is a program of algebraization of geometry using purely geometric notions. Leibniz formulated his program in opposition to algebraization method developed by Descartes. The Euclidean geometry in Szmelew's approach [9] is a theory of structures $\langle S; \parallel, \oplus, O \rangle$, where $\langle S; \parallel, \oplus, O \rangle$ is Desarguesian midpoint plane and $O \subseteq S \times S \times S$ is the relation of equi-orthogonal basis. Points o, p, q are in relation O if they form an isosceles triangle with the right angle in vertex a . If we fix vertices a, p , then there exist exactly two points q, q' such that $O(apq), O(apq')$. Moreover $q \oplus q' = a$. In accordance with the Leibniz program we replace the relation of equi-orthogonal basis by a binary operation $*$: $S \times S \rightarrow S$, called the reper operation. A standard model for the Euclidean geometry in the above sense is the oriented plane over the field of real numbers with the reper operations $*$ defined by the condition: $a * b = q$ iff the point q is the result of rotating of p about right angle around the center a .

MML Identifier: MIDSP_3.

WWW: http://mizar.org/JFM/Vol4/midsp_3.html

The articles [10], [12], [3], [4], [2], [5], [1], [11], [6], and [8] provide the notation and terminology for this paper.

1. SUBSTITUTIONS IN TUPLES

For simplicity, we adopt the following rules: n, i, j, k, l are natural numbers, D is a non empty set, c, d are elements of D , and p, q, r are finite sequences of elements of D .

The following propositions are true:

- (1) If $\text{len } p = j + 1 + k$, then there exist q, r, c such that $\text{len } q = j$ and $\text{len } r = k$ and $p = q \hat{\ } \langle c \rangle \hat{\ } r$.
- (2) If $i \in \text{Seg } n$, then there exist j, k such that $n = j + 1 + k$ and $i = j + 1$.
- (3) Suppose $p = q \hat{\ } \langle c \rangle \hat{\ } r$ and $i = \text{len } q + 1$. Then for every l such that $1 \leq l$ and $l \leq \text{len } q$ holds $p(l) = q(l)$ and $p(i) = c$ and for every l such that $i + 1 \leq l$ and $l \leq \text{len } p$ holds $p(l) = r(l - i)$.
- (4) $l \leq j$ or $l = j + 1$ or $j + 2 \leq l$.
- (5) If $l \in \text{Seg } n \setminus \{i\}$ and $i = j + 1$, then $1 \leq l$ and $l \leq j$ or $i + 1 \leq l$ and $l \leq n$.

Let us consider n, i, D, d and let p be an element of D^{n+1} . Let us assume that $i \in \text{Seg}(n + 1)$. The functor $p(i/d)$ yielding an element of D^{n+1} is defined by:

(Def. 1) $p(i/d)(i) = d$ and for every l such that $l \in \text{dom } p \setminus \{i\}$ holds $p(i/d)(l) = p(l)$.

2. REPER ALGEBRA STRUCTURE AND ITS PROPERTIES

Let us consider n . We consider structures of reper algebra over n as extensions of midpoint algebra structure as systems

\langle a carrier, a midpoint operation, a reper \rangle ,

where the carrier is a set, the midpoint operation is a binary operation on the carrier, and the reper is a function from the carrier ^{n} into the carrier.

Let us consider n , let A be a non empty set, let m be a binary operation on A , and let r be a function from A^n into A . Note that $\langle A, m, r \rangle$ is non empty.

Let us consider n . Observe that there exists a structure of reper algebra over n which is non empty.

Let us consider n . Note that there exists a non empty structure of reper algebra over $n + 2$ which is midpoint algebra-like.

We follow the rules: R_1 is a midpoint algebra-like non empty structure of reper algebra over $n + 2$ and a, b, d, p_1, p'_1 are points of R_1 .

Let us consider i, D . A i -tuple of D is an element of D^i .

Let us consider n, R_1, i . A i -tuple of R_1 is a i -tuple of the carrier of R_1 .

In the sequel p, q are $n + 1$ -tuples of R_1 .

Let us consider n, R_1, a . Then $\langle a \rangle$ is a 1-tuple of R_1 .

Let us consider n, R_1, i, j , let p be a i -tuple of R_1 , and let q be a j -tuple of R_1 . Then $p \wedge q$ is a $i + j$ -tuple of R_1 .

Next we state the proposition

(6) $\langle a \rangle \wedge p$ is a $n + 2$ -tuple of R_1 .

Let us consider n, R_1, a, p . The functor $*(a, p)$ yields a point of R_1 and is defined as follows:

(Def. 2) $*(a, p) = (\text{the reper of } R_1)(\langle a \rangle \wedge p)$.

Let us consider n, i, R_1, d, p . The functor $p_{\uparrow i \rightarrow d}$ yielding a $n + 1$ -tuple of R_1 is defined by the condition (Def. 3).

(Def. 3) Let given D, p' be an element of D^{n+1} , and d' be an element of D . If $D =$ the carrier of R_1 and $p' = p$ and $d' = d$, then $p_{\uparrow i \rightarrow d} = p'(i/d')$.

One can prove the following proposition

(7) If $i \in \text{Seg}(n + 1)$, then $p_{\uparrow i \rightarrow d}(i) = d$ and for every l such that $l \in \text{dom } p \setminus \{i\}$ holds $p_{\uparrow i \rightarrow d}(l) = p(l)$.

Let us consider n . A natural number is called a natural number of n if:

(Def. 4) $1 \leq i$ and $i \leq n + 1$.

In the sequel m denotes a natural number of n .

The following four propositions are true:

(8) i is a natural number of n iff $i \in \text{Seg}(n + 1)$.

(10)¹ If $i \leq n$, then $i + 1$ is a natural number of n .

(11) If for every m holds $p(m) = q(m)$, then $p = q$.

(12)(i) For every natural number l of n such that $l = i$ holds $p_{\uparrow i \rightarrow d}(l) = d$, and

(ii) for all natural numbers l, i of n such that $l \neq i$ holds $p_{\uparrow i \rightarrow d}(l) = p(l)$.

Let us consider n, D , let p be an element of D^{n+1} , and let us consider m . Then $p(m)$ is an element of D .

Let us consider n, R_1 . We say that R_1 is invariance if and only if:

¹ The proposition (9) has been removed.

(Def. 5) For all a, b, p, q such that for every m holds $a^{\otimes} q(m) = b^{\otimes} p(m)$ holds $a^{\otimes} *(b, q) = b^{\otimes} *(a, p)$.

We introduce R_1 is invariance as a synonym of R_1 is invariance.

Let us consider n, i, R_1 . We say that R_1 has property of zero in i if and only if:

(Def. 6) For all a, p holds $*(a, p_{\uparrow i \rightarrow a}) = a$.

Let us consider n, i, R_1 . We say that R_1 is semi additive in i if and only if:

(Def. 7) For all a, p_1, p such that $p(i) = p_1$ holds $*(a, p_{\uparrow i \rightarrow a^{\otimes} p_1}) = a^{\otimes} *(a, p)$.

The following proposition is true

(13) If R_1 is semi additive in m , then for all a, d, p, q such that $q = p_{\uparrow m \rightarrow d}$ holds $*(a, p_{\uparrow m \rightarrow a^{\otimes} d}) = a^{\otimes} *(a, q)$.

Let us consider n, i, R_1 . We say that R_1 is additive in i if and only if:

(Def. 8) For all a, p_1, p'_1, p such that $p(i) = p_1$ holds $*(a, p_{\uparrow i \rightarrow p_1^{\otimes} p'_1}) = *(a, p)^{\otimes} *(a, p_{\uparrow i \rightarrow p'_1})$.

Let us consider n, i, R_1 . We say that R_1 is alternative in i if and only if:

(Def. 9) For all a, p, p_1 such that $p(i) = p_1$ holds $*(a, p_{\uparrow i+1 \rightarrow p_1}) = a$.

In the sequel W is an atlas of R_1 and v is a vector of W .

Let us consider n, R_1, W, i . A i -tuple of W is a i -tuple of the carrier of the algebra of W .

In the sequel x, y denote $n+1$ -tuples of W .

Let us consider n, R_1, W, x, i, v . The functor $x_{\uparrow i \rightarrow v}$ yields a $n+1$ -tuple of W and is defined by the condition (Def. 10).

(Def. 10) Let given D, x' be an element of D^{n+1} , and v' be an element of D . Suppose $D =$ the carrier of the algebra of W and $x' = x$ and $v' = v$. Then $x_{\uparrow i \rightarrow v} = x'(i/v')$.

Next we state three propositions:

(14) If $i \in \text{Seg}(n+1)$, then $x_{\uparrow i \rightarrow v}(i) = v$ and for every l such that $l \in \text{dom } x \setminus \{i\}$ holds $x_{\uparrow i \rightarrow v}(l) = x(l)$.

(15)(i) For every natural number l of n such that $l = i$ holds $x_{\uparrow i \rightarrow v}(l) = v$, and

(ii) for all natural numbers l, i of n such that $l \neq i$ holds $x_{\uparrow i \rightarrow v}(l) = x(l)$.

(16) If for every m holds $x(m) = y(m)$, then $x = y$.

The scheme *SeqLambdaD'* deals with a natural number \mathcal{A} , a non empty set \mathcal{B} , and a unary functor \mathcal{F} yielding an element of \mathcal{B} , and states that:

There exists a finite sequence z of elements of \mathcal{B} such that $\text{len } z = \mathcal{A} + 1$ and for every natural number j of \mathcal{A} holds $z(j) = \mathcal{F}(j)$

for all values of the parameters.

Let us consider n, R_1, W, a, x . The functor $(a, x).W$ yields a $n+1$ -tuple of R_1 and is defined as follows:

(Def. 11) $((a, x).W)(m) = (a, x(m)).W$.

Let us consider n, R_1, W, a, p . The functor $W(a, p)$ yielding a $n+1$ -tuple of W is defined by:

(Def. 12) $W(a, p)(m) = W(a, p(m))$.

Next we state three propositions:

(17) $W(a, p) = x$ iff $(a, x).W = p$.

(18) $W(a, (a, x).W) = x$.

$$(19) \quad (a, W(a, p)).W = p.$$

Let us consider n, R_1, W, a, x . The functor $\Phi(a, x)$ yields a vector of W and is defined by:

$$(Def. 13) \quad \Phi(a, x) = W(a, *(a, (a, x).W)).$$

We now state four propositions:

$$(20) \quad \text{If } W(a, p) = x \text{ and } W(a, b) = v, \text{ then } *(a, p) = b \text{ iff } \Phi(a, x) = v.$$

$$(21) \quad R_1 \text{ is invariance iff for all } a, b, x \text{ holds } \Phi(a, x) = \Phi(b, x).$$

$$(22) \quad 1 \in \text{Seg}(n+1).$$

$$(24)^2 \quad 1 \text{ is a natural number of } n.$$

3. REPER ALGEBRA AND ITS ATLAS

Let us consider n . A midpoint algebra-like non empty structure of reper algebra over $n+2$ is said to be a reper algebra of n if:

(Def. 14) It is invariance.

For simplicity, we follow the rules: R_1 is a reper algebra of n , a, b are points of R_1 , p is a $n+1$ -tuple of R_1 , W is an atlas of R_1 , v is a vector of W , and x is a $n+1$ -tuple of W .

One can prove the following proposition

$$(25) \quad \Phi(a, x) = \Phi(b, x).$$

Let us consider n, R_1, W, x . The functor $\Phi(x)$ yielding a vector of W is defined as follows:

(Def. 15) For every a holds $\Phi(x) = \Phi(a, x)$.

Next we state a number of propositions:

$$(26) \quad \text{If } W(a, p) = x \text{ and } W(a, b) = v \text{ and } \Phi(x) = v, \text{ then } *(a, p) = b.$$

$$(27) \quad \text{If } (a, x).W = p \text{ and } (a, v).W = b \text{ and } *(a, p) = b, \text{ then } \Phi(x) = v.$$

$$(28) \quad \text{If } W(a, p) = x \text{ and } W(a, b) = v, \text{ then } W(a, p \upharpoonright_{m \rightarrow b}) = x \upharpoonright_{m \rightarrow v}.$$

$$(29) \quad \text{If } (a, x).W = p \text{ and } (a, v).W = b, \text{ then } (a, x \upharpoonright_{m \rightarrow v}).W = p \upharpoonright_{m \rightarrow b}.$$

$$(30) \quad R_1 \text{ has property of zero in } m \text{ iff for every } x \text{ holds } \Phi((x \upharpoonright_{m \rightarrow 0_W})) = 0_W.$$

$$(31) \quad R_1 \text{ is semi additive in } m \text{ iff for every } x \text{ holds } \Phi((x \upharpoonright_{m \rightarrow 2x(m)})) = 2\Phi(x).$$

$$(32) \quad \text{If } R_1 \text{ is additive in } m \text{ and has property of zero in } m, \text{ then } R_1 \text{ is semi additive in } m.$$

$$(33) \quad \text{If } R_1 \text{ has property of zero in } m, \text{ then } R_1 \text{ is additive in } m \text{ iff for all } x, v \text{ holds } \Phi((x \upharpoonright_{m \rightarrow x(m)+v})) = \Phi(x) + \Phi((x \upharpoonright_{m \rightarrow v})).$$

$$(34) \quad \text{If } W(a, p) = x \text{ and } m \leq n, \text{ then } W(a, p \upharpoonright_{m+1 \rightarrow p(m)}) = x \upharpoonright_{m+1 \rightarrow x(m)}.$$

$$(35) \quad \text{If } (a, x).W = p \text{ and } m \leq n, \text{ then } (a, x \upharpoonright_{m+1 \rightarrow x(m)}).W = p \upharpoonright_{m+1 \rightarrow p(m)}.$$

$$(36) \quad \text{If } m \leq n, \text{ then } R_1 \text{ is alternative in } m \text{ iff for every } x \text{ holds } \Phi((x \upharpoonright_{m+1 \rightarrow x(m)})) = 0_W.$$

² The proposition (23) has been removed.

REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/nat_1.html.
- [2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.
- [3] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [4] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [5] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_2.html.
- [6] Michał Muzalewski. Midpoint algebras. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/midsp_1.html.
- [7] Michał Muzalewski. *Foundations of Metric-Affine Geometry*. Dział Wydawnictw Filii UW w Białymstoku, Filia UW w Białymstoku, 1990.
- [8] Michał Muzalewski. Atlas of midpoint algebra. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/midsp_2.html.
- [9] Wanda Szmielew. *From Affine to Euclidean Geometry*, volume 27. PWN – D.Reidel Publ. Co., Warszawa – Dordrecht, 1983.
- [10] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [11] Wojciech A. Trybulec. Vectors in real linear space. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/rlvect_1.html.
- [12] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

Received May 28, 1992

Published January 2, 2004
