

Terms Over Many Sorted Universal Algebra¹

Grzegorz Bancerek
Institute of Mathematics
Polish Academy of Sciences

Summary. Pure terms (without constants) over a signature of many sorted universal algebra and terms with constants from algebra are introduced. Facts on evaluation of a term in some valuation are proved.

MML Identifier: MSATERM.

WWW: <http://mizar.org/JFM/Vol6/msaterm.html>

The articles [18], [12], [23], [21], [24], [10], [8], [11], [14], [1], [4], [16], [2], [22], [3], [13], [5], [6], [7], [19], [20], [9], [15], and [17] provide the notation and terminology for this paper.

1. TERMS OVER A SIGNATURE AND OVER AN ALGEBRA

Let I be a non empty set, let X be a non-empty many sorted set indexed by I , and let i be an element of I . Observe that $X(i)$ is non empty.

In the sequel S denotes a non void non empty many sorted signature and V denotes a non-empty many sorted set indexed by the carrier of S .

Let us consider S and let V be a many sorted set indexed by the carrier of S . The functor S -Terms(V) yielding a subset of FinTrees(the carrier of DTConMSA(V)) is defined by:

(Def. 1) S -Terms(V) = TS(DTConMSA(V)).

Let us consider S, V . Note that S -Terms(V) is non empty.

Let us consider S, V . A term of S over V is an element of S -Terms(V).

In the sequel A is an algebra over S and t is a term of S over V .

Let us consider S, V and let o be an operation symbol of S . Then Sym(o, V) is a nonterminal of DTConMSA(V).

Let us consider S, V and let s_1 be a nonterminal of DTConMSA(V). A finite sequence of elements of S -Terms(V) is said to be an argument sequence of s_1 if:

(Def. 2) It is a subtree sequence joinable by s_1 .

We now state the proposition

- (1) Let o be an operation symbol of S and a be a finite sequence. Then $\langle o, \text{the carrier of } S \rangle$ -tree(a) $\in S$ -Terms(V) and a is decorated tree yielding if and only if a is an argument sequence of Sym(o, V).

The scheme *TermInd* deals with a non void non empty many sorted signature \mathcal{A} , a non-empty many sorted set \mathcal{B} indexed by the carrier of \mathcal{A} , and a unary predicate \mathcal{P} , and states that:

¹This article has been prepared during the visit of the author in Nagano in Summer 1994.

For every term t of \mathcal{A} over \mathcal{B} holds $\mathcal{P}[t]$
 provided the following conditions are satisfied:

- For every sort symbol s of \mathcal{A} and for every element v of $\mathcal{B}(s)$ holds \mathcal{P} [the root tree of $\langle v, s \rangle$], and
- Let o be an operation symbol of \mathcal{A} and p be an argument sequence of $\text{Sym}(o, \mathcal{B})$. Suppose that for every term t of \mathcal{A} over \mathcal{B} such that $t \in \text{rng } p$ holds $\mathcal{P}[t]$. Then $\mathcal{P}[\langle o, \text{the carrier of } \mathcal{A} \rangle\text{-tree}(p)]$.

Let us consider S, A, V . A term of A over V is a term of S over $(\text{the sorts of } A) \cup V$.

Let us consider S, A, V and let o be an operation symbol of S . An argument sequence of o, A , and V is an argument sequence of $\text{Sym}(o, (\text{the sorts of } A) \cup V)$.

The scheme *CTermInd* deals with a non void non empty many sorted signature \mathcal{A} , a non-empty algebra \mathcal{B} over \mathcal{A} , a non-empty many sorted set \mathcal{C} indexed by the carrier of \mathcal{A} , and a unary predicate \mathcal{P} , and states that:

For every term t of \mathcal{B} over \mathcal{C} holds $\mathcal{P}[t]$
 provided the parameters meet the following requirements:

- For every sort symbol s of \mathcal{A} and for every element x of $(\text{the sorts of } \mathcal{B})(s)$ holds \mathcal{P} [the root tree of $\langle x, s \rangle$],
- For every sort symbol s of \mathcal{A} and for every element v of $\mathcal{C}(s)$ holds \mathcal{P} [the root tree of $\langle v, s \rangle$], and
- Let o be an operation symbol of \mathcal{A} and p be an argument sequence of o, \mathcal{B} , and \mathcal{C} . Suppose that for every term t of \mathcal{B} over \mathcal{C} such that $t \in \text{rng } p$ holds $\mathcal{P}[t]$. Then $\mathcal{P}[\text{Sym}(o, (\text{the sorts of } \mathcal{B}) \cup \mathcal{C})\text{-tree}(p)]$.

Let us consider S, V, t and let p be a node of t . Then $t(p)$ is a symbol of $\text{DTConMSA}(V)$.

Let us consider S, V . Note that every term of S over V is finite.

The following propositions are true:

- (2)(i) There exists a sort symbol s of S and there exists an element v of $V(s)$ such that $t(\emptyset) = \langle v, s \rangle$, or
- (ii) $t(\emptyset) \in [\text{the operation symbols of } S, \{ \text{the carrier of } S \}]$.
- (3) Let t be a term of A over V . Then
 - (i) there exists a sort symbol s of S and there exists a set x such that $x \in (\text{the sorts of } A)(s)$ and $t(\emptyset) = \langle x, s \rangle$, or
 - (ii) there exists a sort symbol s of S and there exists an element v of $V(s)$ such that $t(\emptyset) = \langle v, s \rangle$, or
 - (iii) $t(\emptyset) \in [\text{the operation symbols of } S, \{ \text{the carrier of } S \}]$.
- (4) For every sort symbol s of S and for every element v of $V(s)$ holds the root tree of $\langle v, s \rangle$ is a term of S over V .
- (5) For every sort symbol s of S and for every element v of $V(s)$ such that $t(\emptyset) = \langle v, s \rangle$ holds $t = \text{the root tree of } \langle v, s \rangle$.
- (6) Let s be a sort symbol of S and x be a set. Suppose $x \in (\text{the sorts of } A)(s)$. Then the root tree of $\langle x, s \rangle$ is a term of A over V .
- (7) Let t be a term of A over V , s be a sort symbol of S , and x be a set. If $x \in (\text{the sorts of } A)(s)$ and $t(\emptyset) = \langle x, s \rangle$, then $t = \text{the root tree of } \langle x, s \rangle$.
- (8) For every sort symbol s of S and for every element v of $V(s)$ holds the root tree of $\langle v, s \rangle$ is a term of A over V .
- (9) Let t be a term of A over V , s be a sort symbol of S , and v be an element of $V(s)$. If $t(\emptyset) = \langle v, s \rangle$, then $t = \text{the root tree of } \langle v, s \rangle$.
- (10) Let o be an operation symbol of S . Suppose $t(\emptyset) = \langle o, \text{the carrier of } S \rangle$. Then there exists an argument sequence a of $\text{Sym}(o, V)$ such that $t = \langle o, \text{the carrier of } S \rangle\text{-tree}(a)$.

Let us consider S , let A be a non-empty algebra over S , let us consider V , let s be a sort symbol of S , and let x be an element of $(\text{the sorts of } A)(s)$. The functor $x_{A,V}$ yields a term of A over V and is defined as follows:

(Def. 3) $x_{A,V}$ = the root tree of $\langle x, s \rangle$.

Let us consider S, A, V , let s be a sort symbol of S , and let v be an element of $V(s)$. The functor v_A yields a term of A over V and is defined by:

(Def. 4) v_A = the root tree of $\langle v, s \rangle$.

Let us consider S, V , let s_1 be a nonterminal of $\text{DTConMSA}(V)$, and let p be an argument sequence of s_1 . Then $s_1\text{-tree}(p)$ is a term of S over V .

The scheme *TermInd2* deals with a non void non empty many sorted signature \mathcal{A} , a non-empty algebra \mathcal{B} over \mathcal{A} , a non-empty many sorted set \mathcal{C} indexed by the carrier of \mathcal{A} , and a unary predicate \mathcal{P} , and states that:

For every term t of \mathcal{B} over \mathcal{C} holds $\mathcal{P}[t]$

provided the parameters meet the following conditions:

- For every sort symbol s of \mathcal{A} and for every element x of $(\text{the sorts of } \mathcal{B})(s)$ holds $\mathcal{P}[x_{\mathcal{B},\mathcal{C}}]$,
- For every sort symbol s of \mathcal{A} and for every element v of $\mathcal{C}(s)$ holds $\mathcal{P}[v_{\mathcal{B}}]$, and
- Let o be an operation symbol of \mathcal{A} and p be an argument sequence of $\text{Sym}(o, (\text{the sorts of } \mathcal{B}) \cup \mathcal{C})$. Suppose that for every term t of \mathcal{B} over \mathcal{C} such that $t \in \text{rng } p$ holds $\mathcal{P}[t]$. Then $\mathcal{P}[\text{Sym}(o, (\text{the sorts of } \mathcal{B}) \cup \mathcal{C})\text{-tree}(p)]$.

2. SORT OF A TERM

We now state three propositions:

- (11) For every term t of S over V there exists a sort symbol s of S such that $t \in \text{FreeSort}(V, s)$.
- (12) For every sort symbol s of S holds $\text{FreeSort}(V, s) \subseteq S\text{-Terms}(V)$.
- (13) $S\text{-Terms}(V) = \bigcup \text{FreeSorts}(V)$.

Let us consider S, V, t . The sort of t yields a sort symbol of S and is defined by:

(Def. 5) $t \in \text{FreeSort}(V, \text{the sort of } t)$.

The following propositions are true:

- (14) Let s be a sort symbol of S and v be an element of $V(s)$. If t = the root tree of $\langle v, s \rangle$, then the sort of t = s .
- (15) Let t be a term of A over V , s be a sort symbol of S , and x be a set. Suppose $x \in (\text{the sorts of } A)(s)$ and t = the root tree of $\langle x, s \rangle$. Then the sort of t = s .
- (16) Let t be a term of A over V , s be a sort symbol of S , and v be an element of $V(s)$. If t = the root tree of $\langle v, s \rangle$, then the sort of t = s .
- (17) Let o be an operation symbol of S . Suppose $t(\emptyset) = \langle o, \text{the carrier of } S \rangle$. Then the sort of t = the result sort of o .
- (18) Let A be a non-empty algebra over S , s be a sort symbol of S , and x be an element of $(\text{the sorts of } A)(s)$. Then the sort of $x_{A,V}$ = s .
- (19) For every sort symbol s of S and for every element v of $V(s)$ holds the sort of v_A = s .
- (20) Let o be an operation symbol of S and p be an argument sequence of $\text{Sym}(o, V)$. Then the sort of $(\text{Sym}(o, V)\text{-tree}(p))$ **qua** term of S over V = the result sort of o .

3. ARGUMENT SEQUENCE

Next we state several propositions:

- (21) Let o be an operation symbol of S and a be a finite sequence of elements of $S\text{-Terms}(V)$. Then a is an argument sequence of $\text{Sym}(o, V)$ if and only if $\text{Sym}(o, V) \Rightarrow$ the roots of a .
- (22) Let o be an operation symbol of S and a be an argument sequence of $\text{Sym}(o, V)$. Then $\text{len } a = \text{len Arity}(o)$ and $\text{dom } a = \text{dom Arity}(o)$ and for every natural number i such that $i \in \text{dom } a$ holds $a(i)$ is a term of S over V .
- (23) Let o be an operation symbol of S , a be an argument sequence of $\text{Sym}(o, V)$, and i be a natural number. Suppose $i \in \text{dom } a$. Let t be a term of S over V . Suppose $t = a(i)$. Then
- (i) $t = (a \text{ qua finite sequence of elements of } S\text{-Terms}(V) \text{ qua non empty set})_i$,
 - (ii) the sort of $t = \text{Arity}(o)(i)$, and
 - (iii) the sort of $t = \text{Arity}(o)_i$.
- (24) Let o be an operation symbol of S and a be a finite sequence. Suppose that
- (i) $\text{len } a = \text{len Arity}(o)$ or $\text{dom } a = \text{dom Arity}(o)$, and
 - (ii) for every natural number i such that $i \in \text{dom } a$ there exists a term t of S over V such that $t = a(i)$ and the sort of $t = \text{Arity}(o)(i)$ or for every natural number i such that $i \in \text{dom } a$ there exists a term t of S over V such that $t = a(i)$ and the sort of $t = \text{Arity}(o)_i$.
- Then a is an argument sequence of $\text{Sym}(o, V)$.
- (25) Let o be an operation symbol of S and a be a finite sequence of elements of $S\text{-Terms}(V)$. Suppose that
- (i) $\text{len } a = \text{len Arity}(o)$ or $\text{dom } a = \text{dom Arity}(o)$, and
 - (ii) for every natural number i such that $i \in \text{dom } a$ and for every term t of S over V such that $t = a(i)$ holds the sort of $t = \text{Arity}(o)(i)$ or for every natural number i such that $i \in \text{dom } a$ and for every term t of S over V such that $t = a(i)$ holds the sort of $t = \text{Arity}(o)_i$.
- Then a is an argument sequence of $\text{Sym}(o, V)$.
- (26) Let S be a non void non empty many sorted signature and V_1, V_2 be non-empty many sorted sets indexed by the carrier of S . If $V_1 \subseteq V_2$, then every term of S over V_1 is a term of S over V_2 .
- (27) Let S be a non void non empty many sorted signature, A be an algebra over S , and V be a non-empty many sorted set indexed by the carrier of S . Then every term of S over V is a term of A over V .

4. COMPOUND TERMS

Let S be a non void non empty many sorted signature and let V be a non-empty many sorted set indexed by the carrier of S . A term of S over V is called a compound term of S over V if:

(Def. 6) $\text{It}(\emptyset) \in \{ \text{the operation symbols of } S, \{ \text{the carrier of } S \} \}$.

Let S be a non void non empty many sorted signature and let V be a non-empty many sorted set indexed by the carrier of S . A non empty subset of $S\text{-Terms}(V)$ is said to be a set with a compound term of S over V if:

(Def. 7) There exists a compound term t of S over V such that $t \in \text{it}$.

The following propositions are true:

- (28) If t is not root, then t is a compound term of S over V .
- (29) For every node p of t holds $t \upharpoonright p$ is a term of S over V .

Let S be a non void non empty many sorted signature, let V be a non-empty many sorted set indexed by the carrier of S , let t be a term of S over V , and let p be a node of t . Then $t \upharpoonright p$ is a term of S over V .

5. EVALUATION OF TERMS

Let S be a non void non empty many sorted signature and let A be an algebra over S . A non-empty many sorted set indexed by the carrier of S is said to be a variables family of A if:

(Def. 8) It misses the sorts of A .

We now state the proposition

(30) Let V be a variables family of A , s be a sort symbol of S , and x be a set. If $x \in$ (the sorts of A)(s), then for every element v of $V(s)$ holds $x \neq v$.

Let S be a non void non empty many sorted signature, let A be a non-empty algebra over S , let V be a non-empty many sorted set indexed by the carrier of S , let t be a term of A over V , let f be a many sorted function from V into the sorts of A , and let v_1 be a finite decorated tree. We say that v_1 is an evaluation of t w.r.t. f if and only if the conditions (Def. 9) are satisfied.

(Def. 9)(i) $\text{dom } v_1 = \text{dom } t$, and

(ii) for every node p of v_1 holds for every sort symbol s of S and for every element v of $V(s)$ such that $t(p) = \langle v, s \rangle$ holds $v_1(p) = f(s)(v)$ and for every sort symbol s of S and for every element x of (the sorts of A)(s) such that $t(p) = \langle x, s \rangle$ holds $v_1(p) = x$ and for every operation symbol o of S such that $t(p) = \langle o, \text{the carrier of } S \rangle$ holds $v_1(p) = (\text{Den}(o, A))(\text{succ}(v_1, p))$.

For simplicity, we follow the rules: S denotes a non void non empty many sorted signature, A denotes a non-empty algebra over S , V denotes a variables family of A , t denotes a term of A over V , and f denotes a many sorted function from V into the sorts of A .

The following propositions are true:

(31) Let s be a sort symbol of S and x be an element of (the sorts of A)(s). Suppose $t =$ the root tree of $\langle x, s \rangle$. Then the root tree of x is an evaluation of t w.r.t. f .

(32) Let s be a sort symbol of S and v be an element of $V(s)$. Suppose $t =$ the root tree of $\langle v, s \rangle$. Then the root tree of $f(s)(v)$ is an evaluation of t w.r.t. f .

(33) Let o be an operation symbol of S , p be an argument sequence of o , A , and V , and q be a decorated tree yielding finite sequence. Suppose that

(i) $\text{len } q = \text{len } p$, and

(ii) for every natural number i and for every term t of A over V such that $i \in \text{dom } p$ and $t = p(i)$ there exists a finite decorated tree v_1 such that $v_1 = q(i)$ and v_1 is an evaluation of t w.r.t. f .

Then there exists a finite decorated tree v_1 such that $v_1 = (\text{Den}(o, A))(\text{the roots of } q)\text{-tree}(q)$ and v_1 is an evaluation of $\text{Sym}(o, (\text{the sorts of } A) \cup V)\text{-tree}(p)$ **qua** term of A over V w.r.t. f .

(34) Let t be a term of A over V and e be a finite decorated tree. Suppose e is an evaluation of t w.r.t. f . Let p be a node of t and n be a node of e . If $n = p$, then $e \upharpoonright n$ is an evaluation of $t \upharpoonright p$ w.r.t. f .

(35) Let o be an operation symbol of S , p be an argument sequence of o , A , and V , and v_1 be a finite decorated tree. Suppose v_1 is an evaluation of $\text{Sym}(o, (\text{the sorts of } A) \cup V)\text{-tree}(p)$ **qua** term of A over V w.r.t. f . Then there exists a decorated tree yielding finite sequence q such that

(i) $\text{len } q = \text{len } p$,

(ii) $v_1 = (\text{Den}(o, A))(\text{the roots of } q)\text{-tree}(q)$, and

(iii) for every natural number i and for every term t of A over V such that $i \in \text{dom } p$ and $t = p(i)$ there exists a finite decorated tree v_1 such that $v_1 = q(i)$ and v_1 is an evaluation of t w.r.t. f .

(36) There exists a finite decorated tree which is an evaluation of t w.r.t. f .

- (37) Let e_1, e_2 be finite decorated trees. Suppose e_1 is an evaluation of t w.r.t. f and e_2 is an evaluation of t w.r.t. f . Then $e_1 = e_2$.
- (38) Let v_1 be a finite decorated tree. Suppose v_1 is an evaluation of t w.r.t. f . Then $v_1(\emptyset) \in (\text{the sorts of } A)(\text{the sort of } t)$.

Let S be a non void non empty many sorted signature, let A be a non-empty algebra over S , let V be a variables family of A , let t be a term of A over V , and let f be a many sorted function from V into the sorts of A . The functor $t^{\textcircled{A}} f$ yielding an element of $(\text{the sorts of } A)(\text{the sort of } t)$ is defined by:

(Def. 10) There exists a finite decorated tree v_1 such that v_1 is an evaluation of t w.r.t. f and $t^{\textcircled{A}} f = v_1(\emptyset)$.

In the sequel t is a term of A over V .

We now state several propositions:

- (39) For every finite decorated tree v_1 such that v_1 is an evaluation of t w.r.t. f holds $t^{\textcircled{A}} f = v_1(\emptyset)$.
- (40) Let v_1 be a finite decorated tree. Suppose v_1 is an evaluation of t w.r.t. f . Let p be a node of t . Then $v_1(p) = t|_p^{\textcircled{A}} f$.
- (41) For every sort symbol s of S and for every element x of $(\text{the sorts of } A)(s)$ holds $x_{A,V}^{\textcircled{A}} f = x$.
- (42) For every sort symbol s of S and for every element v of $V(s)$ holds $v_A^{\textcircled{A}} f = f(s)(v)$.
- (43) Let o be an operation symbol of S , p be an argument sequence of o , A , and V , and q be a finite sequence. Suppose that
- (i) $\text{len } q = \text{len } p$, and
 - (ii) for every natural number i such that $i \in \text{dom } p$ and for every term t of A over V such that $t = p(i)$ holds $q(i) = t^{\textcircled{A}} f$.
- Then $(\text{Sym}(o, (\text{the sorts of } A) \cup V)\text{-tree}(p) \text{ qua term of } A \text{ over } V)^{\textcircled{A}} f = (\text{Den}(o, A))(q)$.

REFERENCES

- [1] Grzegorz Bancerek. Introduction to trees. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/trees_1.html.
- [2] Grzegorz Bancerek. König's theorem. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/card_3.html.
- [3] Grzegorz Bancerek. Cartesian product of functions. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/funct_6.html.
- [4] Grzegorz Bancerek. König's Lemma. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/trees_2.html.
- [5] Grzegorz Bancerek. Sets and functions of trees and joining operations of trees. *Journal of Formalized Mathematics*, 4, 1992. http://mizar.org/JFM/Vol4/trees_3.html.
- [6] Grzegorz Bancerek. Joining of decorated trees. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/trees_4.html.
- [7] Grzegorz Bancerek. Subtrees. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/trees_9.html.
- [8] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.
- [9] Grzegorz Bancerek and Piotr Rudnicki. On defining functions on trees. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/dtconstr.html>.
- [10] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [11] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [12] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.

- [13] Patricia L. Carlson and Grzegorz Bancerek. Context-free grammar — part I. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/lang1.html>.
- [14] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.
- [15] Małgorzata Korolkiewicz. Homomorphisms of many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg_3.html.
- [16] Andrzej Nędzusiak. σ -fields and probability. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/prob_1.html.
- [17] Beata Perkowska. Free many sorted universal algebra. *Journal of Formalized Mathematics*, 6, 1994. <http://mizar.org/JFM/Vol6/msafree.html>.
- [18] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [19] Andrzej Trybulec. Many-sorted sets. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/pboole.html>.
- [20] Andrzej Trybulec. Many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg_1.html.
- [21] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [22] Wojciech A. Trybulec. Pigeon hole principle. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_4.html.
- [23] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [24] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received November 25, 1994

Published January 2, 2004
