## The Correspondence Between Lattices of Subalgebras of Universal Algebras and Many Sorted Algebras

Adam Naumowicz University of Białystok Agnieszka Julia Marasik Warsaw University of Technology

**Summary.** The main goal of this paper is to show some properties of subalgebras of universal algebras and many sorted algebras, and then the isomorphic correspondence between lattices of such subalgebras.

MML Identifier: MSSUBLAT.

WWW: http://mizar.org/JFM/Vol10/mssublat.html

The articles [11], [15], [8], [17], [16], [5], [7], [6], [12], [10], [3], [2], [14], [1], [9], [13], and [4] provide the notation and terminology for this paper.

## 1. Preliminaries

In this paper a is a set and i is a natural number.

The following propositions are true:

- $(1) \quad (\square \longmapsto a)(0) = \emptyset.$
- (2)  $(\square \longmapsto a)(1) = \langle a \rangle$ .
- (3)  $(\square \longmapsto a)(2) = \langle a, a \rangle$ .
- $(4) \quad (\Box \longmapsto a)(3) = \langle a, a, a \rangle.$
- (5) For every finite sequence f of elements of  $\{0\}$  holds  $f = i \mapsto 0$  iff len f = i.
- (6) For every finite sequence f such that  $f = (\square \longmapsto 0)(i)$  holds len f = i.
- 2. Some Properties of Subalgebras of Universal and Many Sorted Algebras

Next we state the proposition

(7) For all universal algebras  $U_1$ ,  $U_2$  such that  $U_1$  is a subalgebra of  $U_2$  holds  $MSSign(U_1) = MSSign(U_2)$ .

Let  $U_0$  be a universal algebra. One can check that the characteristic of  $U_0$  is function yielding. One can prove the following propositions:

- (8) Let  $U_1$ ,  $U_2$  be universal algebras. Suppose  $U_1$  is a subalgebra of  $U_2$ . Let B be a subset of  $MSAlg(U_2)$ . Suppose B = the sorts of  $MSAlg(U_1)$ . Let o be an operation symbol of  $MSSign(U_2)$  and a be an operation symbol of  $MSSign(U_1)$ . If a = o, then  $Den(a, MSAlg(U_1)) = Den(o, MSAlg(U_2)) \upharpoonright Args(a, MSAlg(U_1))$ .
- (9) For all universal algebras  $U_1$ ,  $U_2$  such that  $U_1$  is a subalgebra of  $U_2$  holds the sorts of  $MSAlg(U_1)$  are a subset of  $MSAlg(U_2)$ .
- (10) Let  $U_1$ ,  $U_2$  be universal algebras. Suppose  $U_1$  is a subalgebra of  $U_2$ . Let B be a subset of  $MSAlg(U_2)$ . If B = the sorts of  $MSAlg(U_1)$ , then B is operations closed.
- (11) Let  $U_1$ ,  $U_2$  be universal algebras. Suppose  $U_1$  is a subalgebra of  $U_2$ . Let B be a subset of  $MSAlg(U_2)$ . If B = the sorts of  $MSAlg(U_1)$ , then the characteristics of  $MSAlg(U_1)$  =  $Opers(MSAlg(U_2), B)$ .
- (12) For all universal algebras  $U_1$ ,  $U_2$  such that  $U_1$  is a subalgebra of  $U_2$  holds  $MSAlg(U_1)$  is a subalgebra of  $MSAlg(U_2)$ .
- (13) For all universal algebras  $U_1$ ,  $U_2$  such that  $MSAlg(U_1)$  is a subalgebra of  $MSAlg(U_2)$  holds the carrier of  $U_1$  is a subset of  $U_2$ .
- (14) Let  $U_1$ ,  $U_2$  be universal algebras. Suppose  $MSAlg(U_1)$  is a subalgebra of  $MSAlg(U_2)$ . Let B be a non empty subset of  $U_2$ . If B = the carrier of  $U_1$ , then B is operations closed.
- (15) Let  $U_1$ ,  $U_2$  be universal algebras. Suppose  $MSAlg(U_1)$  is a subalgebra of  $MSAlg(U_2)$ . Let B be a non empty subset of  $U_2$ . If B = the carrier of  $U_1$ , then the characteristic of  $U_1$  =  $Opers(U_2, B)$ .
- (16) For all universal algebras  $U_1$ ,  $U_2$  such that  $MSAlg(U_1)$  is a subalgebra of  $MSAlg(U_2)$  holds  $U_1$  is a subalgebra of  $U_2$ .

In the sequel  $M_1$  is a segmental trivial non void non empty many sorted signature and A is a non-empty algebra over  $M_1$ .

The following propositions are true:

- (17) For every non-empty subalgebra B of A holds the carrier of  $Alg_1(B)$  is a subset of  $Alg_1(A)$ .
- (18) Let B be a non-empty subalgebra of A and S be a non empty subset of  $Alg_1(A)$ . If S = the carrier of  $Alg_1(B)$ , then S is operations closed.
- (19) Let B be a non-empty subalgebra of A and S be a non empty subset of  $Alg_1(A)$ . If S = the carrier of  $Alg_1(B)$ , then the characteristic of  $Alg_1(B) = Opers(Alg_1(A), S)$ .
- (20) For every non-empty subalgebra B of A holds  $Alg_1(B)$  is a subalgebra of  $Alg_1(A)$ .
- (21) Let *S* be a non empty non void many sorted signature and *A*, *B* be algebras over *S*. Then *A* is a subalgebra of *B* if and only if *A* is a subalgebra of the algebra of *B*.
- (22) For all universal algebras A, B holds signature A = signature B iff MSSign(A) = MSSign(B).
- (23) Let A be a non-empty algebra over  $M_1$ . Suppose the carrier of  $M_1 = \{0\}$ . Then  $MSSign(Alg_1(A)) =$  the many sorted signature of  $M_1$ .
- (24) Let A, B be non-empty algebras over  $M_1$ . Suppose the carrier of  $M_1 = \{0\}$  and  $Alg_1(A) = Alg_1(B)$ . Then the algebra of A = the algebra of B.
- (25) Let A be a non-empty algebra over  $M_1$ . If the carrier of  $M_1 = \{0\}$ , then the sorts of A = the sorts of MSAlg(Alg<sub>1</sub>(A)).
- (26) For every non-empty algebra A over  $M_1$  such that the carrier of  $M_1 = \{0\}$  holds  $MSAlg(Alg_1(A)) =$ the algebra of A.
- (27) Let A be a universal algebra and B be a strict non-empty subalgebra of MSAlg(A). If the carrier of  $MSSign(A) = \{0\}$ , then  $Alg_1(B)$  is a subalgebra of A.

3. THE CORRESPONDENCE BETWEEN LATTICES OF SUBALGEBRAS OF UNIVERSAL AND
MANY SORTED ALGEBRAS

Next we state three propositions:

- (28) Let A be a universal algebra,  $a_1$ ,  $b_1$  be strict non-empty subalgebras of A, and  $a_2$ ,  $b_2$  be strict non-empty subalgebras of MSAlg(A). Suppose  $a_2 = \text{MSAlg}(a_1)$  and  $b_2 = \text{MSAlg}(b_1)$ . Then (the sorts of  $a_2$ )  $\cup$  (the sorts of  $b_2$ )  $= 0 \mapsto (\text{the carrier of } a_1) \cup (\text{the carrier of } b_1))$ .
- (29) Let A be a universal algebra,  $a_1$ ,  $b_1$  be strict non-empty subalgebras of A, and  $a_2$ ,  $b_2$  be strict non-empty subalgebras of MSAlg(A). Suppose  $a_2 = \text{MSAlg}(a_1)$  and  $b_2 = \text{MSAlg}(b_1)$ . Then (the sorts of  $a_2$ )  $\cap$  (the sorts of  $b_2$ )  $= 0 \mapsto$  (the carrier of  $a_1$ )  $\cap$  (the carrier of  $b_1$ ).
- (30) Let A be a strict universal algebra,  $a_1$ ,  $b_1$  be strict non-empty subalgebras of A, and  $a_2$ ,  $b_2$  be strict non-empty subalgebras of MSAlg(A). If  $a_2 = \text{MSAlg}(a_1)$  and  $b_2 = \text{MSAlg}(b_1)$ , then MSAlg( $a_1 \sqcup b_1$ ) =  $a_2 \sqcup b_2$ .

Let A be a universal algebra with constants. One can verify that MSSign(A) is non void, strict, segmental, and trivial and has constant operations.

The following proposition is true

(31) Let A be a universal algebra with constants,  $a_1$ ,  $b_1$  be strict non-empty subalgebras of A, and  $a_2$ ,  $b_2$  be strict non-empty subalgebras of MSAlg(A). If  $a_2 = MSAlg(a_1)$  and  $b_2 = MSAlg(b_1)$ , then  $MSAlg(a_1 \cap b_1) = a_2 \cap b_2$ .

Let *A* be a quasi total universal algebra structure. Note that the universal algebra structure of *A* is quasi total.

Let A be a partial universal algebra structure. Note that the universal algebra structure of A is partial.

Let *A* be a non-empty universal algebra structure. Observe that the universal algebra structure of *A* is non-empty.

Let A be a universal algebra with constants. Note that the universal algebra structure of A has constants.

We now state the proposition

(32) Let *A* be a universal algebra with constants. Then the lattice of subalgebras of the universal algebra structure of *A* and the lattice of subalgebras of MSAlg(the universal algebra structure of *A*) are isomorphic.

## REFERENCES

- [1] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq\_1.html.
- [2] Józef Białas. Group and field definitions. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/realsetl.
- [3] Ewa Burakowska. Subalgebras of the universal algebra. Lattices of subalgebras. Journal of Formalized Mathematics, 5, 1993. http://mizar.org/JFM/Vol5/unialg\_2.html.
- [4] Ewa Burakowska. Subalgebras of many sorted algebra. Lattice of subalgebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg 2.html.
- [5] Czesław Byliński. Functions and their basic properties. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/funct\_1.html.
- [6] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct\_2 html
- [7] Czesław Byliński. Partial functions. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/partfun1.html.
- [8] Czesław Byliński. A classical first order language. Journal of Formalized Mathematics, 2, 1990. http://mizar.org/JFM/Vol2/cqc\_lang.html.
- [9] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. Journal of Formalized Mathematics, 2, 1990. http://mizar.org/JFM/Vol2/finseq\_2.html.

- [10] Jarosław Kotowicz, Beata Madras, and Małgorzata Korolkiewicz. Basic notation of universal algebra. Journal of Formalized Mathematics, 4, 1992. http://mizar.org/JFM/Vol4/unialg\_1.html.
- [11] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.
- $[12] \ \ And rzej\ Trybulec.\ Many-sorted\ sets.\ \textit{Journal of Formalized Mathematics}, 5, 1993.\ \texttt{http://mizar.org/JFM/Vol5/pboole.html}.$
- [13] Andrzej Trybulec. Many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg\_l.
- [14] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. http://mizar.org/JFM/Addenda/numbers.html.
- [15] Zinaida Trybulec. Properties of subsets. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/subset\_1.html.
- [16] Edmund Woronowicz. Relations and their basic properties. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/relat 1.html.
- [17] Stanisław Żukowski. Introduction to lattice theory. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/lattices.html.

Received September 22, 1998

Published January 2, 2004