

Subalgebras of Many Sorted Algebra. Lattice of Subalgebras

Ewa Burakowska
Warsaw University
Białystok

MML Identifier: MSUALG_2.

WWW: http://mizar.org/JFM/Vol6/msualg_2.html

The articles [10], [6], [13], [14], [4], [5], [2], [9], [7], [15], [3], [8], [1], [11], and [12] provide the notation and terminology for this paper.

1. AUXILIARY FACTS ABOUT MANY SORTED SETS

In this paper x denotes a set.

The scheme *LambdaB* deals with a non empty set \mathcal{A} and a unary functor \mathcal{F} yielding a set, and states that:

There exists a function f such that $\text{dom } f = \mathcal{A}$ and for every element d of \mathcal{A} holds
 $f(d) = \mathcal{F}(d)$

for all values of the parameters.

Let I be a set, let X be a many sorted set indexed by I , and let Y be a non-empty many sorted set indexed by I . One can check that $X \cup Y$ is non-empty and $Y \cup X$ is non-empty.

The following proposition is true

(2)¹ Let I be a non empty set, X, Y be many sorted sets indexed by I , and i be an element of I^* .
Then $\prod((X \cap Y) \cdot i) = \prod(X \cdot i) \cap \prod(Y \cdot i)$.

Let I be a set and let M be a many sorted set indexed by I . A many sorted set indexed by I is said to be a many sorted subset indexed by M if:

(Def. 1) $It \subseteq M$.

Let I be a set and let M be a non-empty many sorted set indexed by I . One can verify that there exists a many sorted subset indexed by M which is non-empty.

2. CONSTANTS OF A MANY SORTED ALGEBRA

We follow the rules: S is a non void non empty many sorted signature, o is an operation symbol of S , and U_0, U_1, U_2 are algebras over S .

Let S be a non empty many sorted signature and let U_0 be an algebra over S . A subset of U_0 is a many sorted subset indexed by the sorts of U_0 .

Let S be a non empty many sorted signature and let I_1 be a sort symbol of S . We say that I_1 has constants if and only if:

¹ The proposition (1) has been removed.

(Def. 2) There exists an operation symbol o of S such that (the arity of S)(o) = \emptyset and (the result sort of S)(o) = I_1 .

Let I_1 be a non empty many sorted signature. We say that I_1 has constant operations if and only if:

(Def. 3) Every sort symbol of I_1 has constants.

Let A be a non empty set, let B be a set, let a be a function from B into A^* , and let r be a function from B into A . One can check that $\langle A, B, a, r \rangle$ is non empty.

Let us mention that there exists a non empty many sorted signature which is non void and strict and has constant operations.

Let S be a non void non empty many sorted signature, let U_0 be an algebra over S , and let s be a sort symbol of S . The functor $\text{Constants}(U_0, s)$ yielding a subset of (the sorts of U_0)(s) is defined by:

(Def. 4)(i) There exists a non empty set A such that $A = (\text{the sorts of } U_0)(s)$ and $\text{Constants}(U_0, s) = \{a; a \text{ ranges over elements of } A: \bigvee_{o: \text{operation symbol of } S} ((\text{the arity of } S)(o) = \emptyset \wedge (\text{the result sort of } S)(o) = s \wedge a \in \text{rngDen}(o, U_0))\}$ if (the sorts of U_0)(s) $\neq \emptyset$,

(ii) $\text{Constants}(U_0, s) = \emptyset$, otherwise.

Let S be a non void non empty many sorted signature and let U_0 be an algebra over S . The functor $\text{Constants}(U_0)$ yields a subset of U_0 and is defined by:

(Def. 5) For every sort symbol s of S holds $(\text{Constants}(U_0))(s) = \text{Constants}(U_0, s)$.

Let S be a non void non empty many sorted signature with constant operations, let U_0 be a non-empty algebra over S , and let s be a sort symbol of S . Observe that $\text{Constants}(U_0, s)$ is non empty.

Let S be a non void non empty many sorted signature with constant operations and let U_0 be a non-empty algebra over S . One can verify that $\text{Constants}(U_0)$ is non-empty.

3. SUBALGEBRAS OF A MANY SORTED ALGEBRA

Let S be a non void non empty many sorted signature, let U_0 be an algebra over S , let o be an operation symbol of S , and let A be a subset of U_0 . We say that A is closed on o if and only if:

(Def. 6) $\text{rng}(\text{Den}(o, U_0) \upharpoonright (A^\# \cdot \text{the arity of } S)(o)) \subseteq (A \cdot \text{the result sort of } S)(o)$.

Let S be a non void non empty many sorted signature, let U_0 be an algebra over S , and let A be a subset of U_0 . We say that A is operations closed if and only if:

(Def. 7) For every operation symbol o of S holds A is closed on o .

Next we state the proposition

(3) Let S be a non void non empty many sorted signature, o be an operation symbol of S , U_0 be an algebra over S , and B_0, B_1 be subsets of U_0 . If $B_0 \subseteq B_1$, then $(B_0^\# \cdot \text{the arity of } S)(o) \subseteq (B_1^\# \cdot \text{the arity of } S)(o)$.

Let S be a non void non empty many sorted signature, let U_0 be an algebra over S , let o be an operation symbol of S , and let A be a subset of U_0 . Let us assume that A is closed on o . The functor o_A yielding a function from $(A^\# \cdot \text{the arity of } S)(o)$ into $(A \cdot \text{the result sort of } S)(o)$ is defined by:

(Def. 8) $o_A = \text{Den}(o, U_0) \upharpoonright (A^\# \cdot \text{the arity of } S)(o)$.

Let S be a non void non empty many sorted signature, let U_0 be an algebra over S , and let A be a subset of U_0 . The functor $\text{Opers}(U_0, A)$ yields a many sorted function from $A^\# \cdot \text{the arity of } S$ into $A \cdot \text{the result sort of } S$ and is defined as follows:

(Def. 9) For every operation symbol o of S holds $(\text{Opers}(U_0, A))(o) = o_A$.

We now state two propositions:

- (4) Let U_0 be an algebra over S and B be a subset of U_0 . Suppose $B =$ the sorts of U_0 . Then B is operations closed and for every o holds $o_B = \text{Den}(o, U_0)$.
- (5) For every subset B of U_0 such that $B =$ the sorts of U_0 holds $\text{Opers}(U_0, B) =$ the characteristics of U_0 .

Let S be a non void non empty many sorted signature and let U_0 be an algebra over S . An algebra over S is called a subalgebra of U_0 if it satisfies the conditions (Def. 10).

- (Def. 10)(i) The sorts of it are a subset of U_0 , and
- (ii) for every subset B of U_0 such that $B =$ the sorts of it holds B is operations closed and the characteristics of it $= \text{Opers}(U_0, B)$.

Let S be a non void non empty many sorted signature and let U_0 be an algebra over S . Observe that there exists a subalgebra of U_0 which is strict.

Let S be a non void non empty many sorted signature and let U_0 be a non-empty algebra over S . Observe that \langle the sorts of U_0 , the characteristics of $U_0\rangle$ is non-empty.

Let S be a non void non empty many sorted signature and let U_0 be a non-empty algebra over S . Note that there exists a subalgebra of U_0 which is non-empty and strict.

One can prove the following propositions:

- (6) U_0 is a subalgebra of U_0 .
- (7) If U_0 is a subalgebra of U_1 and U_1 is a subalgebra of U_2 , then U_0 is a subalgebra of U_2 .
- (8) If U_1 is a strict subalgebra of U_2 and U_2 is a strict subalgebra of U_1 , then $U_1 = U_2$.
- (9) For all subalgebras U_1, U_2 of U_0 such that the sorts of $U_1 \subseteq$ the sorts of U_2 holds U_1 is a subalgebra of U_2 .
- (10) For all strict subalgebras U_1, U_2 of U_0 such that the sorts of $U_1 =$ the sorts of U_2 holds $U_1 = U_2$.
- (11) Let S be a non void non empty many sorted signature, U_0 be an algebra over S , and U_1 be a subalgebra of U_0 . Then $\text{Constants}(U_0)$ is a subset of U_1 .
- (12) Let S be a non void non empty many sorted signature with constant operations, U_0 be a non-empty algebra over S , and U_1 be a non-empty subalgebra of U_0 . Then $\text{Constants}(U_0)$ is a non-empty subset of U_1 .
- (13) Let S be a non void non empty many sorted signature with constant operations, U_0 be a non-empty algebra over S , and U_1, U_2 be non-empty subalgebras of U_0 . Then \langle the sorts of $U_1\rangle \cap \langle$ the sorts of $U_2\rangle$ is non-empty.

4. MANY SORTED SUBSETS OF MANY SORTED ALGEBRA

Let S be a non void non empty many sorted signature, let U_0 be an algebra over S , and let A be a subset of U_0 . The functor $\text{SubSorts}(A)$ yielding a set is defined by the condition (Def. 11).

- (Def. 11) Let x be a set. Then $x \in \text{SubSorts}(A)$ if and only if the following conditions are satisfied:
- (i) $x \in (2^{\cup(\text{the sorts of } U_0)})^{\text{the carrier of } S}$,
 - (ii) x is a subset of U_0 , and
 - (iii) for every subset B of U_0 such that $B = x$ holds B is operations closed and $\text{Constants}(U_0) \subseteq B$ and $A \subseteq B$.

Let S be a non void non empty many sorted signature, let U_0 be an algebra over S , and let A be a subset of U_0 . Note that $\text{SubSorts}(A)$ is non empty.

Let S be a non void non empty many sorted signature and let U_0 be an algebra over S . The functor $\text{SubSorts}(U_0)$ yielding a set is defined by the condition (Def. 12).

(Def. 12) Let x be a set. Then $x \in \text{SubSorts}(U_0)$ if and only if the following conditions are satisfied:

- (i) $x \in (2^{\cup(\text{the sorts of } U_0)})^{\text{the carrier of } S}$,
- (ii) x is a subset of U_0 , and
- (iii) for every subset B of U_0 such that $B = x$ holds B is operations closed.

Let S be a non void non empty many sorted signature and let U_0 be an algebra over S . One can verify that $\text{SubSorts}(U_0)$ is non empty.

Let S be a non void non empty many sorted signature, let U_0 be an algebra over S , and let e be an element of $\text{SubSorts}(U_0)$. The functor ${}^@e$ yields a subset of U_0 and is defined as follows:

(Def. 13) ${}^@e = e$.

The following two propositions are true:

- (14) For all subsets A, B of U_0 holds $B \in \text{SubSorts}(A)$ iff B is operations closed and $\text{Constants}(U_0) \subseteq B$ and $A \subseteq B$.
- (15) For every subset B of U_0 holds $B \in \text{SubSorts}(U_0)$ iff B is operations closed.

Let S be a non void non empty many sorted signature, let U_0 be an algebra over S , let A be a subset of U_0 , and let s be a sort symbol of S . The functor $\text{SubSort}(A, s)$ yielding a set is defined as follows:

(Def. 14) For every set x holds $x \in \text{SubSort}(A, s)$ iff there exists a subset B of U_0 such that $B \in \text{SubSorts}(A)$ and $x = B(s)$.

Let S be a non void non empty many sorted signature, let U_0 be an algebra over S , let A be a subset of U_0 , and let s be a sort symbol of S . Note that $\text{SubSort}(A, s)$ is non empty.

Let S be a non void non empty many sorted signature, let U_0 be an algebra over S , and let A be a subset of U_0 . The functor $\text{MSSubSort}(A)$ yielding a subset of U_0 is defined as follows:

(Def. 15) For every sort symbol s of S holds $(\text{MSSubSort}(A))(s) = \bigcap \text{SubSort}(A, s)$.

Next we state several propositions:

- (16) For every subset A of U_0 holds $\text{Constants}(U_0) \cup A \subseteq \text{MSSubSort}(A)$.
- (17) For every subset A of U_0 such that $\text{Constants}(U_0) \cup A$ is non-empty holds $\text{MSSubSort}(A)$ is non-empty.
- (18) Let A be a subset of U_0 and B be a subset of U_0 . If $B \in \text{SubSorts}(A)$, then $((\text{MSSubSort}(A))^\# \cdot \text{the arity of } S)(o) \subseteq (B^\# \cdot \text{the arity of } S)(o)$.
- (19) Let A be a subset of U_0 and B be a subset of U_0 . Suppose $B \in \text{SubSorts}(A)$. Then $\text{rng}(\text{Den}(o, U_0) \upharpoonright ((\text{MSSubSort}(A))^\# \cdot \text{the arity of } S)(o)) \subseteq (B \cdot \text{the result sort of } S)(o)$.
- (20) For every subset A of U_0 holds $\text{rng}(\text{Den}(o, U_0) \upharpoonright ((\text{MSSubSort}(A))^\# \cdot \text{the arity of } S)(o)) \subseteq (\text{MSSubSort}(A) \cdot \text{the result sort of } S)(o)$.
- (21) For every subset A of U_0 holds $\text{MSSubSort}(A)$ is operations closed and $A \subseteq \text{MSSubSort}(A)$.

5. OPERATIONS ON MANY SORTED ALGEBRA AND ITS SUBALGEBRAS

Let S be a non void non empty many sorted signature, let U_0 be an algebra over S , and let A be a subset of U_0 . Let us assume that A is operations closed. The functor $U_0|A$ yields a strict subalgebra of U_0 and is defined by:

(Def. 16) $U_0|A = \langle A, (\text{Opers}(U_0, A) \text{ qua many sorted function from } A^\# \cdot \text{the arity of } S \text{ into } A \cdot \text{the result sort of } S) \rangle$.

Let S be a non void non empty many sorted signature, let U_0 be an algebra over S , and let U_1, U_2 be subalgebras of U_0 . The functor $U_1 \cap U_2$ yields a strict subalgebra of U_0 and is defined by the conditions (Def. 17).

(Def. 17)(i) The sorts of $U_1 \cap U_2 = (\text{the sorts of } U_1) \cap (\text{the sorts of } U_2)$, and
(ii) for every subset B of U_0 such that $B = \text{the sorts of } U_1 \cap U_2$ holds B is operations closed and the characteristics of $U_1 \cap U_2 = \text{Opers}(U_0, B)$.

Let S be a non void non empty many sorted signature, let U_0 be an algebra over S , and let A be a subset of U_0 . The functor $\text{Gen}(A)$ yielding a strict subalgebra of U_0 is defined by the conditions (Def. 18).

(Def. 18)(i) A is a subset of $\text{Gen}(A)$, and
(ii) for every subalgebra U_1 of U_0 such that A is a subset of U_1 holds $\text{Gen}(A)$ is a subalgebra of U_1 .

Let S be a non void non empty many sorted signature, let U_0 be a non-empty algebra over S , and let A be a non-empty subset of U_0 . Observe that $\text{Gen}(A)$ is non-empty.

We now state three propositions:

(22) Let S be a non void non empty many sorted signature, U_0 be a strict algebra over S , and B be a subset of U_0 . If $B = \text{the sorts of } U_0$, then $\text{Gen}(B) = U_0$.

(23) Let S be a non void non empty many sorted signature, U_0 be an algebra over S , U_1 be a strict subalgebra of U_0 , and B be a subset of U_0 . If $B = \text{the sorts of } U_1$, then $\text{Gen}(B) = U_1$.

(24) Let S be a non void non empty many sorted signature, U_0 be a non-empty algebra over S , and U_1 be a subalgebra of U_0 . Then $\text{Gen}(\text{Constants}(U_0)) \cap U_1 = \text{Gen}(\text{Constants}(U_0))$.

Let S be a non void non empty many sorted signature, let U_0 be a non-empty algebra over S , and let U_1, U_2 be subalgebras of U_0 . The functor $U_1 \sqcup U_2$ yielding a strict subalgebra of U_0 is defined by:

(Def. 19) For every subset A of U_0 such that $A = (\text{the sorts of } U_1) \cup (\text{the sorts of } U_2)$ holds $U_1 \sqcup U_2 = \text{Gen}(A)$.

Next we state several propositions:

(25) Let S be a non void non empty many sorted signature, U_0 be a non-empty algebra over S , U_1 be a subalgebra of U_0 , and A, B be subsets of U_0 . If $B = A \cup \text{the sorts of } U_1$, then $\text{Gen}(A) \sqcup U_1 = \text{Gen}(B)$.

(26) Let S be a non void non empty many sorted signature, U_0 be a non-empty algebra over S , U_1 be a subalgebra of U_0 , and B be a subset of U_0 . If $B = \text{the sorts of } U_0$, then $\text{Gen}(B) \sqcup U_1 = \text{Gen}(B)$.

(27) Let S be a non void non empty many sorted signature, U_0 be a non-empty algebra over S , and U_1, U_2 be subalgebras of U_0 . Then $U_1 \sqcup U_2 = U_2 \sqcup U_1$.

(28) Let S be a non void non empty many sorted signature, U_0 be a non-empty algebra over S , and U_1, U_2 be strict subalgebras of U_0 . Then $U_1 \cap (U_1 \sqcup U_2) = U_1$.

(29) Let S be a non void non empty many sorted signature, U_0 be a non-empty algebra over S , and U_1, U_2 be strict subalgebras of U_0 . Then $U_1 \cap U_2 \sqcup U_2 = U_2$.

6. LATTICE OF SUBALGEBRAS OF MANY SORTED ALGEBRA

Let S be a non void non empty many sorted signature and let U_0 be an algebra over S . The functor $\text{Subalgebras}(U_0)$ yielding a set is defined as follows:

(Def. 20) For every x holds $x \in \text{Subalgebras}(U_0)$ iff x is a strict subalgebra of U_0 .

Let S be a non void non empty many sorted signature and let U_0 be an algebra over S . Observe that $\text{Subalgebras}(U_0)$ is non empty.

Let S be a non void non empty many sorted signature and let U_0 be a non-empty algebra over S . The functor $\text{MSAlgJoin}(U_0)$ yields a binary operation on $\text{Subalgebras}(U_0)$ and is defined as follows:

(Def. 21) For all elements x, y of $\text{Subalgebras}(U_0)$ and for all strict subalgebras U_1, U_2 of U_0 such that $x = U_1$ and $y = U_2$ holds $(\text{MSAlgJoin}(U_0))(x, y) = U_1 \sqcup U_2$.

Let S be a non void non empty many sorted signature and let U_0 be a non-empty algebra over S . The functor $\text{MSAlgMeet}(U_0)$ yields a binary operation on $\text{Subalgebras}(U_0)$ and is defined as follows:

(Def. 22) For all elements x, y of $\text{Subalgebras}(U_0)$ and for all strict subalgebras U_1, U_2 of U_0 such that $x = U_1$ and $y = U_2$ holds $(\text{MSAlgMeet}(U_0))(x, y) = U_1 \cap U_2$.

In the sequel U_0 denotes a non-empty algebra over S .

The following four propositions are true:

- (30) $\text{MSAlgJoin}(U_0)$ is commutative.
- (31) $\text{MSAlgJoin}(U_0)$ is associative.
- (32) For every non void non empty many sorted signature S and for every non-empty algebra U_0 over S holds $\text{MSAlgMeet}(U_0)$ is commutative.
- (33) For every non void non empty many sorted signature S and for every non-empty algebra U_0 over S holds $\text{MSAlgMeet}(U_0)$ is associative.

Let S be a non void non empty many sorted signature and let U_0 be a non-empty algebra over S . The lattice of subalgebras of U_0 yields a strict lattice and is defined by:

(Def. 23) The lattice of subalgebras of $U_0 = \langle \text{Subalgebras}(U_0), \text{MSAlgJoin}(U_0), \text{MSAlgMeet}(U_0) \rangle$.

The following proposition is true

- (34) Let S be a non void non empty many sorted signature and U_0 be a non-empty algebra over S . Then the lattice of subalgebras of U_0 is bounded.

Let S be a non void non empty many sorted signature and let U_0 be a non-empty algebra over S . Observe that the lattice of subalgebras of U_0 is bounded.

One can prove the following propositions:

- (35) Let S be a non void non empty many sorted signature and U_0 be a non-empty algebra over S . Then $\perp_{\text{the lattice of subalgebras of } U_0} = \text{Gen}(\text{Constants}(U_0))$.
- (36) Let S be a non void non empty many sorted signature, U_0 be a non-empty algebra over S , and B be a subset of U_0 . If $B = \text{the sorts of } U_0$, then $\top_{\text{the lattice of subalgebras of } U_0} = \text{Gen}(B)$.
- (37) Let S be a non void non empty many sorted signature and U_0 be a strict non-empty algebra over S . Then $\top_{\text{the lattice of subalgebras of } U_0} = U_0$.
- (38) Let S be a non void non empty many sorted signature and U_0 be an algebra over S . Then $\langle \text{the sorts of } U_0, \text{the characteristics of } U_0 \rangle$ is a subalgebra of U_0 .
- (39) Let S be a non void non empty many sorted signature and U_0 be a non-empty algebra over S . Then $\langle \text{the sorts of } U_0, \text{the characteristics of } U_0 \rangle$ is non-empty.

- (40) Let S be a non void non empty many sorted signature, U_0 be an algebra over S , and A be a subset of U_0 . Then the sorts of $U_0 \in \text{SubSorts}(A)$.
- (41) Let S be a non void non empty many sorted signature, U_0 be an algebra over S , and A be a subset of U_0 . Then $\text{SubSorts}(A) \subseteq \text{SubSorts}(U_0)$.

REFERENCES

- [1] Grzegorz Bancerek. König's theorem. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/card_3.html.
- [2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.
- [3] Czesław Byliński. Binary operations. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/binop_1.html.
- [4] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [5] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [6] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [7] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_2.html.
- [8] Andrzej Nędzusiak. σ -fields and probability. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/prob_1.html.
- [9] Beata Padlewska. Families of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/setfam_1.html.
- [10] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [11] Andrzej Trybulec. Many-sorted sets. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/pboole.html>.
- [12] Andrzej Trybulec. Many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/msualg_1.html.
- [13] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [14] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.
- [15] Stanisław Żukowski. Introduction to lattice theory. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/lattices.html>.

Received April 25, 1994

Published January 2, 2004
