

# Solving Roots of Polynomial Equation of Degree 4 with Real Coefficients

Xiquan Liang  
 Qingdao University of Science and Technology

**Summary.** In this paper, we describe the definition of the fourth degree algebraic equations and their properties. We clarify the relation between the four roots of this equation and its coefficient. Also, the form of these roots for various conditions is discussed. This solution is known as the Cardano solution.

MML Identifier: POLYEQ\_2.

WWW: [http://mizar.org/JFM/Vol15/polyeq\\_2.html](http://mizar.org/JFM/Vol15/polyeq_2.html)

The articles [1], [4], [2], and [3] provide the notation and terminology for this paper.

Let  $a, b, c, d, e, x$  be real numbers. The functor  $\text{Four}(a, b, c, d, e, x)$  is defined as follows:

(Def. 1)  $\text{Four}(a, b, c, d, e, x) = a \cdot x^4 + b \cdot x^3 + c \cdot x^2 + d \cdot x + e$ .

Let  $a, b, c, d, e, x$  be real numbers. Observe that  $\text{Four}(a, b, c, d, e, x)$  is real.

We now state several propositions:

- (1) Let  $a, c, e, x$  be real numbers. Suppose  $a \neq 0$  and  $e \neq 0$  and  $c^2 - 4 \cdot a \cdot e > 0$ . Suppose  $\text{Four}(a, 0, c, 0, e, x) = 0$ . Then  $x \neq 0$  but  $x = \sqrt{\frac{-c + \sqrt{\Delta(a, c, e)}}{2 \cdot a}}$  or  $x = \sqrt{\frac{-c - \sqrt{\Delta(a, c, e)}}{2 \cdot a}}$  or  $x = -\sqrt{\frac{-c + \sqrt{\Delta(a, c, e)}}{2 \cdot a}}$  or  $x = -\sqrt{\frac{-c - \sqrt{\Delta(a, c, e)}}{2 \cdot a}}$ .
- (2) Let  $a, b, c, x, y$  be real numbers. Suppose  $a \neq 0$  and  $y = x + \frac{1}{x}$ . If  $\text{Four}(a, b, c, b, a, x) = 0$ , then  $x \neq 0$  and  $(a \cdot y^2 + b \cdot y + c) - 2 \cdot a = 0$ .
- (3) Let  $a, b, c, x, y$  be real numbers. Suppose  $a \neq 0$  and  $(b^2 - 4 \cdot a \cdot c) + 8 \cdot a^2 > 0$  and  $y = x + \frac{1}{x}$ . Suppose  $\text{Four}(a, b, c, b, a, x) = 0$ . Let  $y_1, y_2$  be real numbers. Suppose  $y_1 = \frac{-b + \sqrt{(b^2 - 4 \cdot a \cdot c) + 8 \cdot a^2}}{2 \cdot a}$  and  $y_2 = \frac{-b - \sqrt{(b^2 - 4 \cdot a \cdot c) + 8 \cdot a^2}}{2 \cdot a}$ . Then  $x \neq 0$  but  $x = \frac{y_1 + \sqrt{\Delta(1, -y_1, 1)}}{2}$  or  $x = \frac{y_2 + \sqrt{\Delta(1, -y_2, 1)}}{2}$  or  $x = \frac{y_1 - \sqrt{\Delta(1, -y_1, 1)}}{2}$  or  $x = \frac{y_2 - \sqrt{\Delta(1, -y_2, 1)}}{2}$ .
- (4) For every real number  $x$  holds  $x^3 = x^2 \cdot x$  and  $x^3 \cdot x = x^4$  and  $x^2 \cdot x^2 = x^4$ .
- (5) For all real numbers  $x, y$  such that  $x + y \neq 0$  holds  $(x + y)^4 = (x^3 + (3 \cdot y \cdot x^2 + 3 \cdot y^2 \cdot x) + y^3) \cdot x + (x^3 + (3 \cdot y \cdot x^2 + 3 \cdot y^2 \cdot x) + y^3) \cdot y$ .
- (6) For all real numbers  $x, y$  such that  $x + y \neq 0$  holds  $(x + y)^4 = x^4 + (4 \cdot y \cdot x^3 + 6 \cdot y^2 \cdot x^2 + 4 \cdot y^3 \cdot x) + y^4$ .
- (7) Let  $a_1, a_2, a_3, a_4, a_5, b_1, b_2, b_3, b_4, b_5$  be real numbers. Suppose that for every real number  $x$  holds  $\text{Four}(a_1, a_2, a_3, a_4, a_5, x) = \text{Four}(b_1, b_2, b_3, b_4, b_5, x)$ . Then  $a_5 = b_5$  and  $((a_1 - a_2) + a_3) - a_4 = ((b_1 - b_2) + b_3) - b_4$  and  $a_1 + a_2 + a_3 + a_4 = b_1 + b_2 + b_3 + b_4$ .

- (8) Let  $a_1, a_2, a_3, a_4, a_5, b_1, b_2, b_3, b_4, b_5$  be real numbers. Suppose that for every real number  $x$  holds  $\text{Four}(a_1, a_2, a_3, a_4, a_5, x) = \text{Four}(b_1, b_2, b_3, b_4, b_5, x)$ . Then  $a_1 - b_1 = b_3 - a_3$  and  $a_2 - b_2 = b_4 - a_4$ .
- (9) Let  $a_1, a_2, a_3, a_4, a_5, b_1, b_2, b_3, b_4, b_5$  be real numbers. Suppose that for every real number  $x$  holds  $\text{Four}(a_1, a_2, a_3, a_4, a_5, x) = \text{Four}(b_1, b_2, b_3, b_4, b_5, x)$ . Then  $a_1 = b_1$  and  $a_2 = b_2$  and  $a_3 = b_3$  and  $a_4 = b_4$  and  $a_5 = b_5$ .

Let  $a_1, x_1, x_2, x_3, x_4, x$  be real numbers. The functor  $\text{Four0}(a_1, x_1, x_2, x_3, x_4, x)$  is defined as follows:

(Def. 2)  $\text{Four0}(a_1, x_1, x_2, x_3, x_4, x) = a_1 \cdot ((x - x_1) \cdot (x - x_2) \cdot (x - x_3) \cdot (x - x_4))$ .

Let  $a_1, x_1, x_2, x_3, x_4, x$  be real numbers. One can check that  $\text{Four0}(a_1, x_1, x_2, x_3, x_4, x)$  is real.

Next we state four propositions:

- (10) Let  $a_1, a_2, a_3, a_4, a_5, x, x_1, x_2, x_3, x_4$  be real numbers. Suppose  $a_1 \neq 0$ . Suppose that for every real number  $x$  holds  $\text{Four}(a_1, a_2, a_3, a_4, a_5, x) = \text{Four0}(a_1, x_1, x_2, x_3, x_4, x)$ . Then  $\frac{a_1 \cdot x^4 + a_2 \cdot x^3 + a_3 \cdot x^2 + a_4 \cdot x + a_5}{a_1} = ((x^2 \cdot x^2 - (x_1 + x_2 + x_3) \cdot (x^2 \cdot x)) + (x_1 \cdot x_3 + x_2 \cdot x_3 + x_1 \cdot x_2) \cdot x^2) - x_1 \cdot x_2 \cdot x_3 \cdot x - (x - x_1) \cdot (x - x_2) \cdot (x - x_3) \cdot x_4$ .
- (11) Let  $a_1, a_2, a_3, a_4, a_5, x, x_1, x_2, x_3, x_4$  be real numbers. Suppose  $a_1 \neq 0$ . Suppose that for every real number  $x$  holds  $\text{Four}(a_1, a_2, a_3, a_4, a_5, x) = \text{Four0}(a_1, x_1, x_2, x_3, x_4, x)$ . Then  $\frac{a_1 \cdot x^4 + a_2 \cdot x^3 + a_3 \cdot x^2 + a_4 \cdot x + a_5}{a_1} = (((x^4 - (x_1 + x_2 + x_3 + x_4) \cdot x^3) + (x_1 \cdot x_2 + x_1 \cdot x_3 + x_1 \cdot x_4 + (x_2 \cdot x_3 + x_2 \cdot x_4) + x_3 \cdot x_4) \cdot x^2) - (x_1 \cdot x_2 \cdot x_3 + x_1 \cdot x_2 \cdot x_4 + x_1 \cdot x_3 \cdot x_4 + x_2 \cdot x_3 \cdot x_4) \cdot x) + x_1 \cdot x_2 \cdot x_3 \cdot x_4$ .
- (12) Let  $a_1, a_2, a_3, a_4, a_5, x_1, x_2, x_3, x_4$  be real numbers. Suppose  $a_1 \neq 0$  and for every real number  $x$  holds  $\text{Four}(a_1, a_2, a_3, a_4, a_5, x) = \text{Four0}(a_1, x_1, x_2, x_3, x_4, x)$ . Then  $\frac{a_2}{a_1} = -(x_1 + x_2 + x_3 + x_4)$  and  $\frac{a_3}{a_1} = x_1 \cdot x_2 + x_1 \cdot x_3 + x_1 \cdot x_4 + (x_2 \cdot x_3 + x_2 \cdot x_4) + x_3 \cdot x_4$  and  $\frac{a_4}{a_1} = -(x_1 \cdot x_2 \cdot x_3 + x_1 \cdot x_2 \cdot x_4 + x_1 \cdot x_3 \cdot x_4 + x_2 \cdot x_3 \cdot x_4)$  and  $\frac{a_5}{a_1} = x_1 \cdot x_2 \cdot x_3 \cdot x_4$ .
- (13) Let  $a, k, y$  be real numbers. Suppose  $a \neq 0$ . Suppose that for every real number  $x$  holds  $x^4 + a^4 = k \cdot a \cdot x \cdot (x^2 + a^2)$ . Then  $(y^4 - k \cdot y^3 - k \cdot y) + 1 = 0$ .

## REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinal1.html>.
- [2] Rafał Kwiatek. Factorial and Newton coefficients. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol2/newton.html>.
- [3] Jan Popiołek. Quadratic inequalities. *Journal of Formalized Mathematics*, 3, 1991. [http://mizar.org/JFM/Vol3/quin\\_1.html](http://mizar.org/JFM/Vol3/quin_1.html).
- [4] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers operations: min, max, square, and square root. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/square\\_1.html](http://mizar.org/JFM/Vol1/square_1.html).

Received February 3, 2003

Published January 2, 2004