

More on Multivariate Polynomials: Monomials and Constant Polynomials

Christoph Schwarzweller
University of Tuebingen

Summary. In this article we give some technical concepts for multivariate polynomials with arbitrary number of variables. Monomials and constant polynomials are introduced and their properties with respect to the eval functor are shown. In addition, the multiplication of polynomials with coefficients is defined and investigated.

MML Identifier: POLYNOM7.

WWW: <http://mizar.org/JFM/Vol13/polynom7.html>

The articles [18], [8], [22], [23], [24], [5], [10], [2], [7], [6], [9], [1], [13], [19], [3], [17], [20], [15], [4], [12], [14], [16], [11], and [21] provide the notation and terminology for this paper.

1. PRELIMINARIES

Let us observe that there exists a non empty zero structure which is non trivial.

Let us note that every zero structure which is non trivial is also non empty.

Let us mention that there exists a non trivial double loop structure which is Abelian, left zeroed, right zeroed, add-associative, right complementable, unital, associative, commutative, distributive, and integral domain-like.

Let R be a non trivial zero structure. Observe that there exists an element of R which is non-zero.

Let X be a set, let R be a non empty zero structure, and let p be a series of X, R . We say that p is non-zero if and only if:

(Def. 2)¹ $p \neq 0_X R$.

Let X be a set and let R be a non trivial zero structure. Note that there exists a series of X, R which is non-zero.

Let n be an ordinal number and let R be a non trivial zero structure. One can check that there exists a polynomial of n, R which is non-zero.

One can prove the following two propositions:

- (1) Let X be a set, R be a non empty zero structure, and s be a series of X, R . Then $s = 0_X R$ if and only if $\text{Support } s = \emptyset$.
- (2) Let X be a set and R be a non empty zero structure. Then R is non trivial if and only if there exists a series s of X, R such that $\text{Support } s \neq \emptyset$.

Let X be a set and let b be a bag of X . We say that b is univariate if and only if:

(Def. 3) There exists an element u of X such that $\text{support } b = \{u\}$.

¹ The definition (Def. 1) has been removed.

Let X be a non empty set. One can check that there exists a bag of X which is univariate.
 Let X be a non empty set. Note that every bag of X which is univariate is also non empty.

2. POLYNOMIALS WITHOUT VARIABLES

We now state three propositions:

- (3) For every bag b of \emptyset holds $b = \text{EmptyBag } \emptyset$.
- (4) Let L be a right zeroed add-associative right complementable well unital distributive non trivial double loop structure, p be a polynomial of \emptyset , L , and x be a function from \emptyset into L . Then $\text{eval}(p, x) = p(\text{EmptyBag } \emptyset)$.
- (5) Let L be a right zeroed add-associative right complementable well unital distributive non trivial double loop structure. Then $\text{Polynom-Ring}(\emptyset, L)$ is ring isomorphic to L .

3. MONOMIALS

Let X be a set, let L be a non empty zero structure, and let p be a series of X , L . We say that p is monomial-like if and only if:

(Def. 4) There exists a bag b of X such that for every bag b' of X such that $b' \neq b$ holds $p(b') = 0_L$.

Let X be a set and let L be a non empty zero structure. One can verify that there exists a series of X , L which is monomial-like.

Let X be a set and let L be a non empty zero structure. A monomial of X , L is a monomial-like series of X , L .

Let X be a set and let L be a non empty zero structure. One can check that every series of X , L which is monomial-like is also finite-Support.

One can prove the following proposition

- (6) Let X be a set, L be a non empty zero structure, and p be a series of X , L . Then p is a monomial of X , L if and only if $\text{Support } p = \emptyset$ or there exists a bag b of X such that $\text{Support } p = \{b\}$.

Let X be a set, let L be a non empty zero structure, let a be an element of L , and let b be a bag of X . The functor $\text{Monom}(a, b)$ yielding a monomial of X , L is defined as follows:

(Def. 5) $\text{Monom}(a, b) = 0_X L + \cdot (b, a)$.

Let X be a set, let L be a non empty zero structure, and let m be a monomial of X , L . The functor $\text{term } m$ yields a bag of X and is defined as follows:

(Def. 6) $m(\text{term } m) \neq 0_L$ or $\text{Support } m = \emptyset$ and $\text{term } m = \text{EmptyBag } X$.

Let X be a set, let L be a non empty zero structure, and let m be a monomial of X , L . The functor $\text{coefficient } m$ yielding an element of L is defined by:

(Def. 7) $\text{coefficient } m = m(\text{term } m)$.

We now state several propositions:

- (7) For every set X and for every non empty zero structure L and for every monomial m of X , L holds $\text{Support } m = \emptyset$ or $\text{Support } m = \{\text{term } m\}$.
- (8) For every set X and for every non empty zero structure L and for every bag b of X holds $\text{coefficient } \text{Monom}(0_L, b) = 0_L$ and $\text{term } \text{Monom}(0_L, b) = \text{EmptyBag } X$.
- (9) Let X be a set, L be a non empty zero structure, a be an element of L , and b be a bag of X . Then $\text{coefficient } \text{Monom}(a, b) = a$.

- (10) Let X be a set, L be a non trivial zero structure, a be a non-zero element of L , and b be a bag of X . Then $\text{term Monom}(a, b) = b$.
- (11) For every set X and for every non empty zero structure L and for every monomial m of X , L holds $\text{Monom}(\text{coefficient } m, \text{term } m) = m$.
- (12) Let n be an ordinal number, L be a right zeroed add-associative right complementable unital distributive non trivial double loop structure, m be a monomial of n , L , and x be a function from n into L . Then $\text{eval}(m, x) = \text{coefficient } m \cdot \text{eval}(\text{term } m, x)$.
- (13) Let n be an ordinal number, L be a right zeroed add-associative right complementable unital distributive non trivial double loop structure, a be an element of L , b be a bag of n , and x be a function from n into L . Then $\text{eval}(\text{Monom}(a, b), x) = a \cdot \text{eval}(b, x)$.

4. CONSTANT POLYNOMIALS

Let X be a set, let L be a non empty zero structure, and let p be a series of X, L . We say that p is constant if and only if:

- (Def. 8) For every bag b of X such that $b \neq \text{EmptyBag } X$ holds $p(b) = 0_L$.

Let X be a set and let L be a non empty zero structure. Note that there exists a series of X, L which is constant.

Let X be a set and let L be a non empty zero structure. A constant polynomial of X, L is a constant series of X, L .

Let X be a set and let L be a non empty zero structure. Note that every series of X, L which is constant is also monomial-like.

We now state the proposition

- (14) Let X be a set, L be a non empty zero structure, and p be a series of X, L . Then p is a constant polynomial of X, L if and only if $p = 0_X L$ or $\text{Support } p = \{\text{EmptyBag } X\}$.

Let X be a set and let L be a non empty zero structure. Observe that $0_X L$ is constant.

Let X be a set and let L be a unital non empty double loop structure. Note that $1_{-}(X, L)$ is constant.

We now state two propositions:

- (15) Let X be a set, L be a non empty zero structure, and c be a constant polynomial of X, L . Then $\text{Support } c = \emptyset$ or $\text{Support } c = \{\text{EmptyBag } X\}$.
- (16) Let X be a set, L be a non empty zero structure, and c be a constant polynomial of X, L . Then $\text{term } c = \text{EmptyBag } X$ and $\text{coefficient } c = c(\text{EmptyBag } X)$.

Let X be a set, let L be a non empty zero structure, and let a be an element of L . The functor $a_{-}(X, L)$ yielding a series of X, L is defined as follows:

- (Def. 9) $a_{-}(X, L) = 0_X L + \cdot (\text{EmptyBag } X, a)$.

Let X be a set, let L be a non empty zero structure, and let a be an element of L . Note that $a_{-}(X, L)$ is constant.

The following propositions are true:

- (17) Let X be a set, L be a non empty zero structure, and p be a series of X, L . Then p is a constant polynomial of X, L if and only if there exists an element a of L such that $p = a_{-}(X, L)$.
- (18) Let X be a set, L be a non empty multiplicative loop with zero structure, and a be an element of L . Then $(a_{-}(X, L))(\text{EmptyBag } X) = a$ and for every bag b of X such that $b \neq \text{EmptyBag } X$ holds $(a_{-}(X, L))(b) = 0_L$.
- (19) For every set X and for every non empty zero structure L holds $0_{L-}(X, L) = 0_X L$.

- (20) For every set X and for every unital non empty multiplicative loop with zero structure L holds $1_{L-}(X, L) = 1_-(X, L)$.
- (21) Let X be a set, L be a non empty zero structure, and a, b be elements of L . Then $a_-(X, L) = b_-(X, L)$ if and only if $a = b$.
- (22) For every set X and for every non empty zero structure L and for every element a of L holds $\text{Support}(a_-(X, L)) = \emptyset$ or $\text{Support}(a_-(X, L)) = \{\text{EmptyBag } X\}$.
- (23) For every set X and for every non empty zero structure L and for every element a of L holds term $a_-(X, L) = \text{EmptyBag } X$ and coefficient $(a_-(X, L)) = a$.
- (24) Let n be an ordinal number, L be a right zeroed add-associative right complementable unital distributive non trivial double loop structure, c be a constant polynomial of n, L , and x be a function from n into L . Then $\text{eval}(c, x) = \text{coefficient } c$.
- (25) Let n be an ordinal number, L be a right zeroed add-associative right complementable unital distributive non trivial double loop structure, a be an element of L , and x be a function from n into L . Then $\text{eval}(a_-(n, L), x) = a$.

5. MULTIPLICATION WITH COEFFICIENTS

Let X be a set, let L be a non empty multiplicative loop with zero structure, let p be a series of X, L , and let a be an element of L . The functor $a \cdot p$ yielding a series of X, L is defined by:

- (Def. 10) For every bag b of X holds $(a \cdot p)(b) = a \cdot p(b)$.

The functor $p \cdot a$ yielding a series of X, L is defined by:

- (Def. 11) For every bag b of X holds $(p \cdot a)(b) = p(b) \cdot a$.

Let X be a set, let L be a left zeroed right zeroed add-cancelable distributive non empty double loop structure, let p be a finite-Support series of X, L , and let a be an element of L . Observe that $a \cdot p$ is finite-Support and $p \cdot a$ is finite-Support.

One can prove the following propositions:

- (26) Let X be a set, L be a commutative non empty multiplicative loop with zero structure, p be a series of X, L , and a be an element of L . Then $a \cdot p = p \cdot a$.
- (27) Let n be an ordinal number, L be an add-associative right complementable right zeroed left distributive non empty double loop structure, p be a series of n, L , and a be an element of L . Then $a \cdot p = (a_-(n, L)) * p$.
- (28) Let n be an ordinal number, L be an add-associative right complementable right zeroed right distributive non empty double loop structure, p be a series of n, L , and a be an element of L . Then $p \cdot a = p * (a_-(n, L))$.
- (29) Let n be an ordinal number, L be an Abelian left zeroed right zeroed add-associative right complementable unital associative commutative distributive non trivial double loop structure, p be a polynomial of n, L , a be an element of L , and x be a function from n into L . Then $\text{eval}(a \cdot p, x) = a \cdot \text{eval}(p, x)$.
- (30) Let n be an ordinal number, L be a left zeroed right zeroed add-left-cancelable add-associative right complementable unital associative integral domain-like distributive non trivial double loop structure, p be a polynomial of n, L , a be an element of L , and x be a function from n into L . Then $\text{eval}(a \cdot p, x) = a \cdot \text{eval}(p, x)$.
- (31) Let n be an ordinal number, L be an Abelian left zeroed right zeroed add-associative right complementable unital associative commutative distributive non trivial double loop structure, p be a polynomial of n, L , a be an element of L , and x be a function from n into L . Then $\text{eval}(p \cdot a, x) = \text{eval}(p, x) \cdot a$.

- (32) Let n be an ordinal number, L be a left zeroed right zeroed add-left-cancelable add-associative right complementable unital associative commutative distributive integral domain-like non trivial double loop structure, p be a polynomial of n , L , a be an element of L , and x be a function from n into L . Then $\text{eval}(p \cdot a, x) = \text{eval}(p, x) \cdot a$.
- (33) Let n be an ordinal number, L be an Abelian left zeroed right zeroed add-associative right complementable unital associative commutative distributive non trivial double loop structure, p be a polynomial of n , L , a be an element of L , and x be a function from n into L . Then $\text{eval}((a \cdot (n, L)) * p, x) = a \cdot \text{eval}(p, x)$.
- (34) Let n be an ordinal number, L be an Abelian left zeroed right zeroed add-associative right complementable unital associative commutative distributive non trivial double loop structure, p be a polynomial of n , L , a be an element of L , and x be a function from n into L . Then $\text{eval}(p * (a \cdot (n, L)), x) = \text{eval}(p, x) \cdot a$.

REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/nat_1.html.
- [2] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Voll/ordinal1.html>.
- [3] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/finseq_1.html.
- [4] Józef Białas. Group and field definitions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Voll/realset1.html>.
- [5] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/funct_1.html.
- [6] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/funct_2.html.
- [7] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Voll/partfun1.html>.
- [8] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/zfmisc_1.html.
- [9] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/funct_4.html.
- [10] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/finset_1.html.
- [11] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/vectsp_1.html.
- [12] Michał Muzalewski. Construction of rings and left-, right-, and bi-modules over a ring. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/vectsp_2.html.
- [13] Michał Muzalewski and Wojciech Skaba. From loops to abelian multiplicative groups with zero. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/algstr_1.html.
- [14] Piotr Rudnicki and Andrzej Trybulec. Multivariate polynomials with arbitrary number of variables. *Journal of Formalized Mathematics*, 11, 1999. <http://mizar.org/JFM/Vol11/polynom1.html>.
- [15] Christoph Schwarzweller. The field of quotients over an integral domain. *Journal of Formalized Mathematics*, 10, 1998. <http://mizar.org/JFM/Vol10/quofield.html>.
- [16] Christoph Schwarzweller and Andrzej Trybulec. The evaluation of multivariate polynomials. *Journal of Formalized Mathematics*, 12, 2000. <http://mizar.org/JFM/Vol12/polynom2.html>.
- [17] Andrzej Trybulec. Binary operations applied to functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/funcop_1.html.
- [18] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [19] Wojciech A. Trybulec. Vectors in real linear space. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/rvect_1.html.
- [20] Wojciech A. Trybulec. Groups. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol1/group_1.html.
- [21] Wojciech A. Trybulec. Pigeon hole principle. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_4.html.

- [22] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [23] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.
- [24] Edmund Woronowicz. Relations defined on sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relset_1.html.

Received November 28, 2001

Published January 2, 2004
