

# Topological Spaces and Continuous Functions<sup>1</sup>

Beata Padlewska  
Warsaw University  
Białystok

Agata Darmochwał  
Warsaw University  
Białystok

**Summary.** The paper contains a definition of topological space. The following notions are defined: point of topological space, subset of topological space, subspace of topological space, and continuous function.

MML Identifier: PRE\_TOPC.

WWW: [http://mizar.org/JFM/Voll/pre\\_topc.html](http://mizar.org/JFM/Voll/pre_topc.html)

The articles [4], [2], [5], [6], [1], and [3] provide the notation and terminology for this paper.

We consider topological structures as extensions of 1-sorted structure as systems  $\langle a \text{ carrier, a topology } \rangle$ ,

where the carrier is a set and the topology is a family of subsets of the carrier.

In the sequel  $T$  is a topological structure.

Let  $I_1$  be a topological structure. We say that  $I_1$  is topological space-like if and only if the conditions (Def. 1) are satisfied.

(Def. 1)(i) The carrier of  $I_1 \in$  the topology of  $I_1$ ,

(ii) for every family  $a$  of subsets of  $I_1$  such that  $a \subseteq$  the topology of  $I_1$  holds  $\bigcup a \in$  the topology of  $I_1$ , and

(iii) for all subsets  $a, b$  of  $I_1$  such that  $a \in$  the topology of  $I_1$  and  $b \in$  the topology of  $I_1$  holds  $a \cap b \in$  the topology of  $I_1$ .

Let us note that there exists a topological structure which is non empty, strict, and topological space-like.

A topological space is a topological space-like topological structure.

Let  $S$  be a 1-sorted structure. A point of  $S$  is an element of  $S$ .

In the sequel  $G_1$  is a topological space.

Next we state the proposition

(5)<sup>1</sup>  $\emptyset \in$  the topology of  $G_1$ .

Let  $T$  be a 1-sorted structure. The functor  $\theta_T$  yields a subset of  $T$  and is defined as follows:

(Def. 2)  $\theta_T = \emptyset$ .

The functor  $\Omega_T$  yielding a subset of  $T$  is defined by:

(Def. 3)  $\Omega_T =$  the carrier of  $T$ .

---

<sup>1</sup>Supported by RPB.P.III-24.C1.

<sup>1</sup> The propositions (1)–(4) have been removed.

Let  $T$  be a 1-sorted structure. Observe that  $\emptyset_T$  is empty.  
One can prove the following proposition

(12)<sup>2</sup> For every 1-sorted structure  $T$  holds  $\Omega_T =$  the carrier of  $T$ .

Let  $T$  be a non empty 1-sorted structure. Note that  $\Omega_T$  is non empty.  
The following propositions are true:

- (13) For every non empty 1-sorted structure  $T$  and for every point  $p$  of  $T$  holds  $p \in \Omega_T$ .
- (14) For every 1-sorted structure  $T$  and for every subset  $P$  of  $T$  holds  $P \subseteq \Omega_T$ .
- (15) For every 1-sorted structure  $T$  and for every subset  $P$  of  $T$  holds  $P \cap \Omega_T = P$ .
- (16) For every 1-sorted structure  $T$  and for every set  $A$  such that  $A \subseteq \Omega_T$  holds  $A$  is a subset of  $T$ .
- (17) For every 1-sorted structure  $T$  and for every subset  $P$  of  $T$  holds  $P^c = \Omega_T \setminus P$ .
- (18) For every 1-sorted structure  $T$  and for every subset  $P$  of  $T$  holds  $P \cup P^c = \Omega_T$ .
- (19) For every 1-sorted structure  $T$  and for all subsets  $P, Q$  of  $T$  holds  $P \subseteq Q$  iff  $Q^c \subseteq P^c$ .
- (20) For every 1-sorted structure  $T$  and for every subset  $P$  of  $T$  holds  $P = (P^c)^c$ .
- (21) For every 1-sorted structure  $T$  and for all subsets  $P, Q$  of  $T$  holds  $P \subseteq Q^c$  iff  $P$  misses  $Q$ .
- (22) For every 1-sorted structure  $T$  and for every subset  $P$  of  $T$  holds  $\Omega_T \setminus (\Omega_T \setminus P) = P$ .
- (23) For every 1-sorted structure  $T$  and for every subset  $P$  of  $T$  holds  $P \neq \Omega_T$  iff  $\Omega_T \setminus P \neq \emptyset$ .
- (24) For every 1-sorted structure  $T$  and for all subsets  $P, Q$  of  $T$  such that  $\Omega_T \setminus P = Q$  holds  $\Omega_T = P \cup Q$ .
- (25) For every 1-sorted structure  $T$  and for all subsets  $P, Q$  of  $T$  such that  $\Omega_T = P \cup Q$  and  $P$  misses  $Q$  holds  $Q = \Omega_T \setminus P$ .
- (26) For every 1-sorted structure  $T$  and for every subset  $P$  of  $T$  holds  $P$  misses  $P^c$ .
- (27) For every 1-sorted structure  $T$  holds  $\Omega_T = (\emptyset_T)^c$ .

Let  $T$  be a topological structure and let  $P$  be a subset of  $T$ . We say that  $P$  is open if and only if:

(Def. 5)<sup>3</sup>  $P \in$  the topology of  $T$ .

Let  $T$  be a topological structure and let  $P$  be a subset of  $T$ . We say that  $P$  is closed if and only if:

(Def. 6)  $\Omega_T \setminus P$  is open.

Let  $T$  be a 1-sorted structure and let  $F$  be a family of subsets of  $T$ . Then  $\bigcup F$  is a subset of  $T$ .

Let  $T$  be a 1-sorted structure and let  $F$  be a family of subsets of  $T$ . Then  $\bigcap F$  is a subset of  $T$ .

Let  $T$  be a 1-sorted structure and let  $F$  be a family of subsets of  $T$ . We say that  $F$  is a cover of  $T$  if and only if:

(Def. 8)<sup>4</sup>  $\Omega_T = \bigcup F$ .

Let  $T$  be a topological structure. A topological structure is said to be a subspace of  $T$  if it satisfies the conditions (Def. 9).

<sup>2</sup> The propositions (6)–(11) have been removed.

<sup>3</sup> The definition (Def. 4) has been removed.

<sup>4</sup> The definition (Def. 7) has been removed.

(Def. 9)(i)  $\Omega_{it} \subseteq \Omega_T$ , and

(ii) for every subset  $P$  of it holds  $P \in$  the topology of it iff there exists a subset  $Q$  of  $T$  such that  $Q \in$  the topology of  $T$  and  $P = Q \cap \Omega_{it}$ .

Let  $T$  be a topological structure. Observe that there exists a subspace of  $T$  which is strict.

Let  $T$  be a non empty topological structure. Observe that there exists a subspace of  $T$  which is strict and non empty.

The scheme *SubFamExS* deals with a topological structure  $\mathcal{A}$  and a unary predicate  $\mathcal{P}$ , and states that:

There exists a family  $F$  of subsets of  $\mathcal{A}$  such that for every subset  $B$  of  $\mathcal{A}$  holds  $B \in F$   
iff  $\mathcal{P}[B]$

for all values of the parameters.

Let  $T$  be a topological space. One can check that every subspace of  $T$  is topological space-like.

Let  $T$  be a topological structure and let  $P$  be a subset of  $T$ . The functor  $T \upharpoonright P$  yielding a strict subspace of  $T$  is defined as follows:

(Def. 10)  $\Omega_{T \upharpoonright P} = P$ .

Let  $T$  be a non empty topological structure and let  $P$  be a non empty subset of  $T$ . Note that  $T \upharpoonright P$  is non empty.

Let  $T$  be a topological space. Note that there exists a subspace of  $T$  which is topological space-like and strict.

Let  $T$  be a topological space and let  $P$  be a subset of  $T$ . Note that  $T \upharpoonright P$  is topological space-like.

Let  $S, T$  be 1-sorted structures. A map from  $S$  into  $T$  is a function from the carrier of  $S$  into the carrier of  $T$ .

Let  $S, T$  be 1-sorted structures, let  $f$  be a function from the carrier of  $S$  into the carrier of  $T$ , and let  $P$  be a set. Then  $f \circ P$  is a subset of  $T$ .

Let  $S, T$  be 1-sorted structures, let  $f$  be a function from the carrier of  $S$  into the carrier of  $T$ , and let  $P$  be a set. Then  $f^{-1}(P)$  is a subset of  $S$ .

Let  $S, T$  be topological structures and let  $f$  be a map from  $S$  into  $T$ . We say that  $f$  is continuous if and only if:

(Def. 12)<sup>5</sup> For every subset  $P_1$  of  $T$  such that  $P_1$  is closed holds  $f^{-1}(P_1)$  is closed.

The scheme *TopAbstr* deals with a topological structure  $\mathcal{A}$  and a unary predicate  $\mathcal{P}$ , and states that:

There exists a subset  $P$  of  $\mathcal{A}$  such that for every set  $x$  such that  $x \in$  the carrier of  $\mathcal{A}$   
holds  $x \in P$  iff  $\mathcal{P}[x]$

for all values of the parameters.

Next we state three propositions:

(39)<sup>6</sup> For every subspace  $X'$  of  $T$  holds every subset of  $X'$  is a subset of  $T$ .

(41)<sup>7</sup> For every subset  $A$  of  $T$  such that  $A \neq \emptyset_T$  there exists a point  $x$  of  $T$  such that  $x \in A$ .

(42)  $\Omega_{(G_1)}$  is closed.

Let  $T$  be a topological space. One can verify that  $\Omega_T$  is closed.

Let  $T$  be a topological space. Note that there exists a subset of  $T$  which is closed.

Let  $T$  be a non empty topological space. One can verify that there exists a subset of  $T$  which is non empty and closed.

We now state two propositions:

(43) Let  $X'$  be a subspace of  $T$  and  $B$  be a subset of  $X'$ . Then  $B$  is closed if and only if there exists a subset  $C$  of  $T$  such that  $C$  is closed and  $C \cap \Omega_{X'} = B$ .

<sup>5</sup> The definition (Def. 11) has been removed.

<sup>6</sup> The propositions (28)–(38) have been removed.

<sup>7</sup> The proposition (40) has been removed.

- (44) Let  $F$  be a family of subsets of  $G_1$ . Suppose that for every subset  $A$  of  $G_1$  such that  $A \in F$  holds  $A$  is closed. Then  $\bigcap F$  is closed.

Let  $G_1$  be a topological structure and let  $A$  be a subset of  $G_1$ . The functor  $\bar{A}$  yields a subset of  $G_1$  and is defined by the condition (Def. 13).

- (Def. 13) Let  $p$  be a set. Suppose  $p \in$  the carrier of  $G_1$ . Then  $p \in \bar{A}$  if and only if for every subset  $G$  of  $G_1$  such that  $G$  is open holds if  $p \in G$ , then  $A$  meets  $G$ .

We now state a number of propositions:

- (45) Let  $A$  be a subset of  $T$  and  $p$  be a set. Suppose  $p \in$  the carrier of  $T$ . Then  $p \in \bar{A}$  if and only if for every subset  $C$  of  $T$  such that  $C$  is closed holds if  $A \subseteq C$ , then  $p \in C$ .
- (46) Let  $A$  be a subset of  $G_1$ . Then there exists a family  $F$  of subsets of  $G_1$  such that for every subset  $C$  of  $G_1$  holds  $C \in F$  iff  $C$  is closed and  $A \subseteq C$  and  $\bar{A} = \bigcap F$ .
- (47) For every subspace  $X'$  of  $T$  and for every subset  $A$  of  $T$  and for every subset  $A_1$  of  $X'$  such that  $A = A_1$  holds  $\bar{A}_1 = \bar{A} \cap \Omega_{X'}$ .
- (48) For every subset  $A$  of  $T$  holds  $A \subseteq \bar{A}$ .
- (49) For all subsets  $A, B$  of  $T$  such that  $A \subseteq B$  holds  $\bar{A} \subseteq \bar{B}$ .
- (50) For all subsets  $A, B$  of  $G_1$  holds  $\overline{A \cup B} = \bar{A} \cup \bar{B}$ .
- (51) For all subsets  $A, B$  of  $T$  holds  $\overline{A \cap B} \subseteq \bar{A} \cap \bar{B}$ .
- (52) Let  $A$  be a subset of  $T$ . Then
- (i) if  $A$  is closed, then  $\bar{A} = A$ , and
  - (ii) if  $T$  is topological space-like and  $\bar{A} = A$ , then  $A$  is closed.
- (53) Let  $A$  be a subset of  $T$ . Then
- (i) if  $A$  is open, then  $\overline{\Omega_T \setminus A} = \Omega_T \setminus A$ , and
  - (ii) if  $T$  is topological space-like and  $\overline{\Omega_T \setminus A} = \Omega_T \setminus A$ , then  $A$  is open.
- (54) Let  $A$  be a subset of  $T$  and  $p$  be a point of  $T$ . Then  $p \in \bar{A}$  if and only if the following conditions are satisfied:
- (i)  $T$  is non empty, and
  - (ii) for every subset  $G$  of  $T$  such that  $G$  is open holds if  $p \in G$ , then  $A$  meets  $G$ .

## REFERENCES

- [1] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/funct\\_2.html](http://mizar.org/JFM/Vol1/funct_2.html).
- [2] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/zfmisc\\_1.html](http://mizar.org/JFM/Vol1/zfmisc_1.html).
- [3] Beata Padlewska. Families of sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/setfam\\_1.html](http://mizar.org/JFM/Vol1/setfam_1.html).
- [4] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [5] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/subset\\_1.html](http://mizar.org/JFM/Vol1/subset_1.html).

- [6] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/relat\\_1.html](http://mizar.org/JFM/Vol1/relat_1.html).

*Received April 14, 1989*

*Published January 2, 2004*

---