

Some Basic Properties of Many Sorted Sets

Artur Korniłowicz
Institute of Mathematics
Warsaw University
Białystok

MML Identifier: PZFMISC1.

WWW: <http://mizar.org/JFM/Vol7/pzfmisc1.html>

The articles [8], [2], [10], [1], [4], [5], [3], [9], [7], and [6] provide the notation and terminology for this paper.

1. PRELIMINARIES

For simplicity, we follow the rules: i, I denote sets, f denotes a function, $x, x_1, x_2, y, A, B, X, Y, Z$ denote many sorted sets indexed by I , J denotes a non empty set, and N_1 denotes a many sorted set indexed by J .

Next we state three propositions:

- (1) For every set X and for every many sorted set M indexed by I such that $i \in I$ holds $\text{dom}(M + \cdot(i \rightarrow X)) = I$.
- (2) If $f = \emptyset$, then f is a many sorted set indexed by \emptyset .
- (3) If I is non empty, then there exists no X which is empty yielding and non-empty.

2. SINGLETON AND UNORDERED PAIRS

Let us consider I, A . The functor $\{A\}$ yielding a many sorted set indexed by I is defined by:

- (Def. 1) For every i such that $i \in I$ holds $\{A\}(i) = \{A(i)\}$.

Let us consider I, A . One can check that $\{A\}$ is non-empty and locally-finite.

Let us consider I, A, B . The functor $\{A, B\}$ yielding a many sorted set indexed by I is defined by:

- (Def. 2) For every i such that $i \in I$ holds $\{A, B\}(i) = \{A(i), B(i)\}$.

Let us notice that the functor $\{A, B\}$ is commutative.

Let us consider I, A, B . Note that $\{A, B\}$ is non-empty and locally-finite.

The following propositions are true:

- (4) $X = \{y\}$ iff for every x holds $x \in X$ iff $x = y$.
- (5) If for every x holds $x \in X$ iff $x = x_1$ or $x = x_2$, then $X = \{x_1, x_2\}$.
- (6) If $X = \{x_1, x_2\}$, then for every x such that $x = x_1$ or $x = x_2$ holds $x \in X$.

- (7) $\{N_1\} \neq \mathbf{0}_I$.
- (8) If $x \in \{A\}$, then $x = A$.
- (9) $x \in \{x\}$.
- (10) If $x = A$ or $x = B$, then $x \in \{A, B\}$.
- (11) $\{A\} \cup \{B\} = \{A, B\}$.
- (12) $\{x, x\} = \{x\}$.
- (14)¹ If $\{A\} \subseteq \{B\}$, then $A = B$.
- (15) If $\{x\} = \{y\}$, then $x = y$.
- (16) If $\{x\} = \{A, B\}$, then $x = A$ and $x = B$.
- (17) If $\{x\} = \{A, B\}$, then $A = B$.
- (18) $\{x\} \subseteq \{x, y\}$ and $\{y\} \subseteq \{x, y\}$.
- (19) If $\{x\} \cup \{y\} = \{x\}$ or $\{x\} \cup \{y\} = \{y\}$, then $x = y$.
- (20) $\{x\} \cup \{x, y\} = \{x, y\}$.
- (22)² If I is non empty and $\{x\} \cap \{y\} = \mathbf{0}_I$, then $x \neq y$.
- (23) If $\{x\} \cap \{y\} = \{x\}$ or $\{x\} \cap \{y\} = \{y\}$, then $x = y$.
- (24) $\{x\} \cap \{x, y\} = \{x\}$ and $\{y\} \cap \{x, y\} = \{y\}$.
- (25) If I is non empty and $\{x\} \setminus \{y\} = \{x\}$, then $x \neq y$.
- (26) If $\{x\} \setminus \{y\} = \mathbf{0}_I$, then $x = y$.
- (27) $\{x\} \setminus \{x, y\} = \mathbf{0}_I$ and $\{y\} \setminus \{x, y\} = \mathbf{0}_I$.
- (28) If $\{x\} \subseteq \{y\}$, then $\{x\} = \{y\}$.
- (29) If $\{x, y\} \subseteq \{A\}$, then $x = A$ and $y = A$.
- (30) If $\{x, y\} \subseteq \{A\}$, then $\{x, y\} = \{A\}$.
- (31) $2^{\{x\}} = \{\mathbf{0}_I, \{x\}\}$.
- (32) $\{A\} \subseteq 2^A$.
- (33) $\bigcup \{x\} = x$.
- (34) $\bigcup \{\{x\}, \{y\}\} = \{x, y\}$.
- (35) $\bigcup \{A, B\} = A \cup B$.
- (36) $\{x\} \subseteq X$ iff $x \in X$.
- (37) $\{x_1, x_2\} \subseteq X$ iff $x_1 \in X$ and $x_2 \in X$.
- (38) If $A = \mathbf{0}_I$ or $A = \{x_1\}$ or $A = \{x_2\}$ or $A = \{x_1, x_2\}$, then $A \subseteq \{x_1, x_2\}$.

¹ The proposition (13) has been removed.

² The proposition (21) has been removed.

3. SUM OF UNORDERED PAIRS (OR A SINGLETON) AND A SET

Next we state several propositions:

- (39) If $x \in A$ or $x = B$, then $x \in A \cup \{B\}$.
- (40) $A \cup \{x\} \subseteq B$ iff $x \in B$ and $A \subseteq B$.
- (41) If $\{x\} \cup X = X$, then $x \in X$.
- (42) If $x \in X$, then $\{x\} \cup X = X$.
- (43) $\{x, y\} \cup A = A$ iff $x \in A$ and $y \in A$.
- (44) If I is non empty, then $\{x\} \cup X \neq \mathbf{0}_I$.
- (45) If I is non empty, then $\{x, y\} \cup X \neq \mathbf{0}_I$.

4. INTERSECTION OF UNORDERED PAIRS (OR A SINGLETON) AND A SET

We now state several propositions:

- (46) If $X \cap \{x\} = \{x\}$, then $x \in X$.
- (47) If $x \in X$, then $X \cap \{x\} = \{x\}$.
- (48) $x \in X$ and $y \in X$ iff $\{x, y\} \cap X = \{x, y\}$.
- (49) If I is non empty and $\{x\} \cap X = \mathbf{0}_I$, then $x \notin X$.
- (50) If I is non empty and $\{x, y\} \cap X = \mathbf{0}_I$, then $x \notin X$ and $y \notin X$.

5. DIFFERENCE OF UNORDERED PAIRS (OR A SINGLETON) AND A SET

The following propositions are true:

- (51) If $y \in X \setminus \{x\}$, then $y \in X$.
- (52) If I is non empty and $y \in X \setminus \{x\}$, then $y \neq x$.
- (53) If I is non empty and $X \setminus \{x\} = X$, then $x \notin X$.
- (54) If I is non empty and $\{x\} \setminus X = \{x\}$, then $x \notin X$.
- (55) $\{x\} \setminus X = \mathbf{0}_I$ iff $x \in X$.
- (56) If I is non empty and $\{x, y\} \setminus X = \{x\}$, then $x \notin X$.
- (58)³ If I is non empty and $\{x, y\} \setminus X = \{x, y\}$, then $x \notin X$ and $y \notin X$.
- (59) $\{x, y\} \setminus X = \mathbf{0}_I$ iff $x \in X$ and $y \in X$.
- (60) If $X = \mathbf{0}_I$ or $X = \{x\}$ or $X = \{y\}$ or $X = \{x, y\}$, then $X \setminus \{x, y\} = \mathbf{0}_I$.

³ The proposition (57) has been removed.

6. CARTESIAN PRODUCT

One can prove the following propositions:

- (61) If $X = \mathbf{0}_I$ or $Y = \mathbf{0}_I$, then $\llbracket X, Y \rrbracket = \mathbf{0}_I$.
- (62) If X is non-empty and Y is non-empty and $\llbracket X, Y \rrbracket = \llbracket Y, X \rrbracket$, then $X = Y$.
- (63) If $\llbracket X, X \rrbracket = \llbracket Y, Y \rrbracket$, then $X = Y$.
- (64) If Z is non-empty and if $\llbracket X, Z \rrbracket \subseteq \llbracket Y, Z \rrbracket$ or $\llbracket Z, X \rrbracket \subseteq \llbracket Z, Y \rrbracket$, then $X \subseteq Y$.
- (65) If $X \subseteq Y$, then $\llbracket X, Z \rrbracket \subseteq \llbracket Y, Z \rrbracket$ and $\llbracket Z, X \rrbracket \subseteq \llbracket Z, Y \rrbracket$.
- (66) If $x_1 \subseteq A$ and $x_2 \subseteq B$, then $\llbracket x_1, x_2 \rrbracket \subseteq \llbracket A, B \rrbracket$.
- (67) $\llbracket X \cup Y, Z \rrbracket = \llbracket X, Z \rrbracket \cup \llbracket Y, Z \rrbracket$ and $\llbracket Z, X \cup Y \rrbracket = \llbracket Z, X \rrbracket \cup \llbracket Z, Y \rrbracket$.
- (68) $\llbracket x_1 \cup x_2, A \cup B \rrbracket = \llbracket x_1, A \rrbracket \cup \llbracket x_1, B \rrbracket \cup \llbracket x_2, A \rrbracket \cup \llbracket x_2, B \rrbracket$.
- (69) $\llbracket X \cap Y, Z \rrbracket = \llbracket X, Z \rrbracket \cap \llbracket Y, Z \rrbracket$ and $\llbracket Z, X \cap Y \rrbracket = \llbracket Z, X \rrbracket \cap \llbracket Z, Y \rrbracket$.
- (70) $\llbracket x_1 \cap x_2, A \cap B \rrbracket = \llbracket x_1, A \rrbracket \cap \llbracket x_2, B \rrbracket$.
- (71) If $A \subseteq X$ and $B \subseteq Y$, then $\llbracket A, Y \rrbracket \cap \llbracket X, B \rrbracket = \llbracket A, B \rrbracket$.
- (72) $\llbracket X \setminus Y, Z \rrbracket = \llbracket X, Z \rrbracket \setminus \llbracket Y, Z \rrbracket$ and $\llbracket Z, X \setminus Y \rrbracket = \llbracket Z, X \rrbracket \setminus \llbracket Z, Y \rrbracket$.
- (73) $\llbracket x_1, x_2 \rrbracket \setminus \llbracket A, B \rrbracket = \llbracket x_1 \setminus A, x_2 \rrbracket \cup \llbracket x_1, x_2 \setminus B \rrbracket$.
- (74) If $x_1 \cap x_2 = \mathbf{0}_I$ or $A \cap B = \mathbf{0}_I$, then $\llbracket x_1, A \rrbracket \cap \llbracket x_2, B \rrbracket = \mathbf{0}_I$.
- (75) If X is non-empty, then $\llbracket \{x\}, X \rrbracket$ is non-empty and $\llbracket X, \{x\} \rrbracket$ is non-empty.
- (76) $\llbracket \{x, y\}, X \rrbracket = \llbracket \{x\}, X \rrbracket \cup \llbracket \{y\}, X \rrbracket$ and $\llbracket X, \{x, y\} \rrbracket = \llbracket X, \{x\} \rrbracket \cup \llbracket X, \{y\} \rrbracket$.
- (77) If x_1 is non-empty and A is non-empty and $\llbracket x_1, A \rrbracket = \llbracket x_2, B \rrbracket$, then $x_1 = x_2$ and $A = B$.
- (78) If $X \subseteq \llbracket X, Y \rrbracket$ or $X \subseteq \llbracket Y, X \rrbracket$, then $X = \mathbf{0}_I$.
- (79) If $A \in \llbracket x, y \rrbracket$ and $A \in \llbracket X, Y \rrbracket$, then $A \in \llbracket x \cap X, y \cap Y \rrbracket$.
- (80) If $\llbracket x, X \rrbracket \subseteq \llbracket y, Y \rrbracket$ and $\llbracket x, X \rrbracket$ is non-empty, then $x \subseteq y$ and $X \subseteq Y$.
- (81) If $A \subseteq X$, then $\llbracket A, A \rrbracket \subseteq \llbracket X, X \rrbracket$.
- (82) If $X \cap Y = \mathbf{0}_I$, then $\llbracket X, Y \rrbracket \cap \llbracket Y, X \rrbracket = \mathbf{0}_I$.
- (83) If A is non-empty and if $\llbracket A, B \rrbracket \subseteq \llbracket X, Y \rrbracket$ or $\llbracket B, A \rrbracket \subseteq \llbracket Y, X \rrbracket$, then $B \subseteq Y$.
- (84) If $x \subseteq \llbracket A, B \rrbracket$ and $y \subseteq \llbracket X, Y \rrbracket$, then $x \cup y \subseteq \llbracket A \cup X, B \cup Y \rrbracket$.

REFERENCES

- [1] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol11/funct_1.html.
- [2] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [3] Czesław Byliński. A classical first order language. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol12/cqc_lang.html.
- [4] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol12/funct_4.html.
- [5] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.
- [6] Artur Korniłowicz. Definitions and basic properties of boolean and union of many sorted sets. *Journal of Formalized Mathematics*, 7, 1995. <http://mizar.org/JFM/Vol7/mboolean.html>.

- [7] Beata Madras. Products of many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/pralg_2.html.
- [8] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [9] Andrzej Trybulec. Many-sorted sets. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/pboole.html>.
- [10] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received September 29, 1995

Published January 2, 2004
