JOURNAL OF FORMALIZED MATHEMATICS
Volume 2, Released 1990, Published 2003
Inst. of Computer Science, Univ. of Białystok

Linear Independence in Right Module over Domain

Michał Muzalewski
Warsaw University
Białystok

Wojciech Skaba
Nicolaus Copernicus University
Toruń

MML Identifier: RMOD_5.

WWW: http://mizar.org/JFM/Vol2/rmod_5.html

The articles [9], [13], [3], [1], [2], [10], [11], [12], [4], [5], [8], [7], and [6] provide the notation and terminology for this paper.

For simplicity, we adopt the following rules: x denotes a set, R denotes a ring, V denotes a right module over R, v, v_1, v_2 denote vectors of V, and A, B denote subsets of V.

Let us consider R, let us consider V, and let I_1 be a subset of V. We say that I_1 is linearly independent if and only if:

(Def. 1) For every linear combination l of I_1 such that $\sum l = 0_V$ holds the support of $l = \emptyset$.

We introduce I_1 is linearly dependent as an antonym of I_1 is linearly independent.

One can prove the following propositions:

(2)[1] If $A \subseteq B$ and B is linearly independent, then A is linearly independent.

(3) If $0_R \neq \mathbf{1}_R$ and A is linearly independent, then $0_V \notin A$.

(4) $\emptyset_{\text{the carrier of } V}$ is linearly independent.

(5) If $0_R \neq \mathbf{1}_R$ and $\{v_1, v_2\}$ is linearly independent, then $v_1 \neq 0_V$ and $v_2 \neq 0_V$.

(6) If $0_R \neq \mathbf{1}_R$, then $\{v, 0_V\}$ is linearly dependent and $\{0_V, v\}$ is linearly dependent.

For simplicity, we use the following convention: R denotes an integral domain, V denotes a right module over R, A, B denote subsets of V, and l denotes a linear combination of A.

Let us consider R, let us consider V, and let us consider A. The functor $\mathrm{Lin}(A)$ yielding a strict submodule of V is defined by:

(Def. 2) The carrier of $\mathrm{Lin}(A) = \{\sum l\}$.

One can prove the following propositions:

(9)[2] $x \in \mathrm{Lin}(A)$ iff there exists l such that $x = \sum l$.

(10) If $x \in A$, then $x \in \mathrm{Lin}(A)$.

(11) $\mathrm{Lin}(\emptyset_{\text{the carrier of } V}) = \mathbf{0}_V$.

(12) If $\mathrm{Lin}(A) = \mathbf{0}_V$, then $A = \emptyset$ or $A = \{0_V\}$.

[1] The proposition (1) has been removed.
[2] The propositions (7) and (8) have been removed.

(13) For every strict submodule W of V such that $0_R \neq \mathbf{1}_R$ and $A =$ the carrier of W holds $\mathrm{Lin}(A) = W$.

(14) Let V be a strict right module over R and A be a subset of V. If $0_R \neq \mathbf{1}_R$ and $A =$ the carrier of V, then $\mathrm{Lin}(A) = V$.

(15) If $A \subseteq B$, then $\mathrm{Lin}(A)$ is a submodule of $\mathrm{Lin}(B)$.

(16) For every strict right module V over R and for all subsets A, B of V such that $\mathrm{Lin}(A) = V$ and $A \subseteq B$ holds $\mathrm{Lin}(B) = V$.

(17) $\mathrm{Lin}(A \cup B) = \mathrm{Lin}(A) + \mathrm{Lin}(B)$.

(18) $\mathrm{Lin}(A \cap B)$ is a submodule of $\mathrm{Lin}(A) \cap \mathrm{Lin}(B)$.

REFERENCES

[1] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.

[2] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.

[3] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.

[4] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/vectsp_1.html.

[5] Michał Muzalewski. Construction of rings and left-, right-, and bi-modules over a ring. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/vectsp_2.html.

[6] Michał Muzalewski and Wojciech Skaba. Linear combinations in right module over associative ring. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/rmod_4.html.

[7] Michał Muzalewski and Wojciech Skaba. Operations on submodules in right module over associative ring. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/rmod_3.html.

[8] Michał Muzalewski and Wojciech Skaba. Submodules and cosets of submodules in right module over associative ring. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/rmod_2.html.

[9] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.

[10] Andrzej Trybulec. Function domains and Frænkel operator. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/fraenkel.html.

[11] Wojciech A. Trybulec. Vectors in real linear space. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/rlvect_1.html.

[12] Wojciech A. Trybulec. Linear combinations in real linear space. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/rlvect_2.html.

[13] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

Received October 22, 1990

Published January 2, 2004