

Banach Space of Absolute Summable Real Sequences

Yasumasa Suzuki
2-14-18 Take, Yokosuka-shi
Kanagawa, Japan

Noboru Endou
Gifu National College of Technology

Yasunari Shidama
Shinshu University
Nagano

Summary. A continuation of [5]. As the example of real norm spaces, we introduce the arithmetic addition and multiplication in the set of absolute summable real sequences and introduce the norm also. This set has the structure of the Banach space.

MML Identifier: RSSPACE3.

WWW: <http://mizar.org/JFM/Vol15/rsspace3.html>

The articles [13], [17], [4], [1], [14], [7], [2], [3], [18], [16], [10], [15], [11], [9], [8], [12], and [6] provide the notation and terminology for this paper.

1. L1-SPACE: THE SPACE OF ABSOLUTE SUMMABLE REAL SEQUENCES

The subset the set of l1-real sequences of the linear space of real sequences is defined by the condition (Def. 1).

(Def. 1) Let x be a set. Then $x \in$ the set of l1-real sequences if and only if $x \in$ the set of real sequences and $\text{id}_{\text{seq}}(x)$ is absolutely summable.

One can verify that the set of l1-real sequences is non empty.

The following two propositions are true:

- (1) The set of l1-real sequences is linearly closed.
- (2) \langle the set of l1-real sequences, Zero_- (the set of l1-real sequences, the linear space of real sequences), Add_- (the set of l1-real sequences, the linear space of real sequences), Mult_- (the set of l1-real sequences, the linear space of real sequences) \rangle is a subspace of the linear space of real sequences.

Let us observe that \langle the set of l1-real sequences, Zero_- (the set of l1-real sequences, the linear space of real sequences), Add_- (the set of l1-real sequences, the linear space of real sequences), Mult_- (the set of l1-real sequences, the linear space of real sequences) \rangle is Abelian, add-associative, right zeroed, right complementable, and real linear space-like.

Next we state the proposition

- (3) \langle the set of l1-real sequences, Zero_- (the set of l1-real sequences, the linear space of real sequences), Add_- (the set of l1-real sequences, the linear space of real sequences), Mult_- (the set of l1-real sequences, the linear space of real sequences) \rangle is a real linear space.

The function norm_{seq} from the set of 11-real sequences into \mathbb{R} is defined as follows:

(Def. 2) For every set x such that $x \in$ the set of 11-real sequences holds $\text{norm}_{\text{seq}}(x) = \sum |\text{id}_{\text{seq}}(x)|$.

Let X be a non empty set, let Z be an element of X , let A be a binary operation on X , let M be a function from $[\mathbb{R}, X]$ into X , and let N be a function from X into \mathbb{R} . Observe that $\langle X, Z, A, M, N \rangle$ is non empty.

One can prove the following four propositions:

- (4) Let l be a normed structure. Suppose \langle the carrier of l , the zero of l , the addition of l , the external multiplication of l \rangle is a real linear space. Then l is a real linear space.
- (5) Let r_1 be a sequence of real numbers. Suppose that for every natural number n holds $r_1(n) = 0$. Then r_1 is absolutely summable and $\sum |r_1| = 0$.
- (6) Let r_1 be a sequence of real numbers. Suppose r_1 is absolutely summable and $\sum |r_1| = 0$. Let n be a natural number. Then $r_1(n) = 0$.
- (7) \langle the set of 11-real sequences, Zero_(the set of 11-real sequences, the linear space of real sequences), Add_(the set of 11-real sequences, the linear space of real sequences), Mult_(the set of 11-real sequences, the linear space of real sequences), norm_{seq} \rangle is a real linear space.

The non empty normed structure 11-Space is defined by the condition (Def. 3).

(Def. 3) 11-Space = \langle the set of 11-real sequences, Zero_(the set of 11-real sequences, the linear space of real sequences), Add_(the set of 11-real sequences, the linear space of real sequences), Mult_(the set of 11-real sequences, the linear space of real sequences), norm_{seq} \rangle .

2. L1_SPACE IS BANACH

We now state two propositions:

- (8) The carrier of 11-Space = the set of 11-real sequences and for every set x holds x is an element of 11-Space iff x is a sequence of real numbers and $\text{id}_{\text{seq}}(x)$ is absolutely summable and for every set x holds x is a vector of 11-Space iff x is a sequence of real numbers and $\text{id}_{\text{seq}}(x)$ is absolutely summable and $0_{11\text{-Space}} = \text{Zero}_{\text{seq}}$ and for every vector u of 11-Space holds $u = \text{id}_{\text{seq}}(u)$ and for all vectors u, v of 11-Space holds $u + v = \text{id}_{\text{seq}}(u) + \text{id}_{\text{seq}}(v)$ and for every real number r and for every vector u of 11-Space holds $r \cdot u = r \text{id}_{\text{seq}}(u)$ and for every vector u of 11-Space holds $-u = -\text{id}_{\text{seq}}(u)$ and $\text{id}_{\text{seq}}(-u) = -\text{id}_{\text{seq}}(u)$ and for all vectors u, v of 11-Space holds $u - v = \text{id}_{\text{seq}}(u) - \text{id}_{\text{seq}}(v)$ and for every vector v of 11-Space holds $\text{id}_{\text{seq}}(v)$ is absolutely summable and for every vector v of 11-Space holds $\|v\| = \sum |\text{id}_{\text{seq}}(v)|$.
- (9) Let x, y be points of 11-Space and a be a real number. Then $\|x\| = 0$ iff $x = 0_{11\text{-Space}}$ and $0 \leq \|x\|$ and $\|x + y\| \leq \|x\| + \|y\|$ and $\|a \cdot x\| = |a| \cdot \|x\|$.

Let us note that 11-Space is real normed space-like, real linear space-like, Abelian, add-associative, right zeroed, and right complementable.

Let X be a non empty normed structure and let x, y be points of X . The functor $\rho(x, y)$ yielding a real number is defined by:

(Def. 4) $\rho(x, y) = \|x - y\|$.

Let N_1 be a non empty normed structure and let s_1 be a sequence of N_1 . We say that s_1 is CCauchy if and only if the condition (Def. 5) is satisfied.

(Def. 5) Let r_2 be a real number. Suppose $r_2 > 0$. Then there exists a natural number k_1 such that for all natural numbers n_1, m_1 if $n_1 \geq k_1$ and $m_1 \geq k_1$, then $\rho(s_1(n_1), s_1(m_1)) < r_2$.

We introduce s_1 is Cauchy sequence by norm as a synonym of s_1 is CCauchy.

In the sequel N_1 denotes a non empty real normed space and s_2 denotes a sequence of N_1 .

Next we state two propositions:

- (10) s_2 is Cauchy sequence by norm if and only if for every real number r such that $r > 0$ there exists a natural number k such that for all natural numbers n, m such that $n \geq k$ and $m \geq k$ holds $\|s_2(n) - s_2(m)\| < r$.
- (11) For every sequence v_1 of l_1 -Space such that v_1 is Cauchy sequence by norm holds v_1 is convergent.

REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinal1.html>.
- [2] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [3] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [4] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [5] Noboru Endou, Yasumasa Suzuki, and Yasunari Shidama. Hilbert space of real sequences. *Journal of Formalized Mathematics*, 15, 2003. <http://mizar.org/JFM/Vol15/rsspace2.html>.
- [6] Noboru Endou, Yasumasa Suzuki, and Yasunari Shidama. Real linear space of real sequences. *Journal of Formalized Mathematics*, 15, 2003. <http://mizar.org/JFM/Vol15/rsspace.html>.
- [7] Krzysztof Hryniewiecki. Basic properties of real numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/real_1.html.
- [8] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/seqm_3.html.
- [9] Jarosław Kotowicz. Real sequences and basic operations on them. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/seq_1.html.
- [10] Jan Popiołek. Some properties of functions modul and signum. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/absvalue.html>.
- [11] Jan Popiołek. Real normed space. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/normsp_1.html.
- [12] Konrad Raczkowski and Andrzej Nędzusiak. Series. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/series_1.html.
- [13] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [14] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [15] Wojciech A. Trybulec. Subspaces and cosets of subspaces in real linear space. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/rlsub_1.html.
- [16] Wojciech A. Trybulec. Vectors in real linear space. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/rlvect_1.html.
- [17] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [18] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received August 8, 2003

Published January 2, 2004
