

Schemes¹

Stanisław T. Czuba
Warsaw University
Białystok

Summary. Some basic schemes of quantifier calculus are proved.

MML Identifier: SCHEMS_1.

WWW: http://mizar.org/JFM/Vol2/schems_1.html

In this paper a, b denote sets.

In this article we present several logical schemes. The scheme *Schemat0* concerns a unary predicate \mathcal{P} , and states that:

There exists a such that $\mathcal{P}[a]$

provided the following requirement is met:

- For every a holds $\mathcal{P}[a]$.

The scheme *Schemat1a* concerns a 0-ary predicate Q and a unary predicate \mathcal{P} , and states that:

For every a holds $\mathcal{P}[a]$ and Q

provided the parameters have the following property:

- For every a holds $\mathcal{P}[a]$ and Q .

The scheme *Schemat1b* concerns a 0-ary predicate Q and a unary predicate \mathcal{P} , and states that:

For every a holds $\mathcal{P}[a]$ and Q

provided the parameters meet the following requirement:

- For every a holds $\mathcal{P}[a]$ and Q .

The scheme *Schemat2a* concerns a 0-ary predicate Q and a unary predicate \mathcal{P} , and states that:

There exists a such that $\mathcal{P}[a]$ or Q

provided the following condition is met:

- There exists a such that $\mathcal{P}[a]$ or Q .

The scheme *Schemat2b* concerns a 0-ary predicate Q and a unary predicate \mathcal{P} , and states that:

There exists a such that $\mathcal{P}[a]$ or Q

provided the parameters meet the following condition:

- There exists a such that $\mathcal{P}[a]$ or Q .

The scheme *Schemat3* concerns a binary predicate \mathcal{P} , and states that:

For every b there exists a such that $\mathcal{P}[a, b]$

provided the following condition is met:

- There exists a such that for every b holds $\mathcal{P}[a, b]$.

The scheme *Schemat4a* concerns two unary predicates \mathcal{P}, Q , and states that:

There exists a such that $\mathcal{P}[a]$ or there exists a such that $Q[a]$

provided the following condition is satisfied:

- There exists a such that $\mathcal{P}[a]$ or $Q[a]$.

The scheme *Schemat4b* concerns two unary predicates \mathcal{P}, Q , and states that:

There exists a such that $\mathcal{P}[a]$ or $Q[a]$

¹Supported by RPB.P.III-24.C10.

provided the following requirement is met:

- There exists a such that $\mathcal{P}[a]$ or there exists a such that $Q[a]$.

The scheme *Schemat5* concerns two unary predicates \mathcal{P} , Q , and states that:

There exists a such that $\mathcal{P}[a]$ and there exists a such that $Q[a]$

provided the parameters satisfy the following condition:

- There exists a such that $\mathcal{P}[a]$ and $Q[a]$.

The scheme *Schemat6a* concerns two unary predicates \mathcal{P} , Q , and states that:

For every a holds $\mathcal{P}[a]$ and for every a holds $Q[a]$

provided the parameters satisfy the following condition:

- For every a holds $\mathcal{P}[a]$ and $Q[a]$.

The scheme *Schemat6b* concerns two unary predicates \mathcal{P} , Q , and states that:

For every a holds $\mathcal{P}[a]$ and $Q[a]$

provided the following condition is satisfied:

- For every a holds $\mathcal{P}[a]$ and for every a holds $Q[a]$.

The scheme *Schemat7* concerns two unary predicates \mathcal{P} , Q , and states that:

For every a holds $\mathcal{P}[a]$ or $Q[a]$

provided the following condition is met:

- For every a holds $\mathcal{P}[a]$ or for every a holds $Q[a]$.

The scheme *Schemat8* concerns two unary predicates \mathcal{P} , Q , and states that:

If for every a holds $\mathcal{P}[a]$, then for every a holds $Q[a]$

provided the parameters satisfy the following condition:

- For every a such that $\mathcal{P}[a]$ holds $Q[a]$.

The scheme *Schemat9* concerns two unary predicates \mathcal{P} , Q , and states that:

For every a holds $\mathcal{P}[a]$ iff for every a holds $Q[a]$

provided the parameters meet the following requirement:

- For every a holds $\mathcal{P}[a]$ iff $Q[a]$.

The scheme *Schemat10b* concerns a 0-ary predicate \mathcal{P} , and states that:

For every a holds \mathcal{P}

provided the following requirement is met:

- \mathcal{P} .

The scheme *Schemat11a* concerns a 0-ary predicate Q and a unary predicate \mathcal{P} , and states that:

For every a holds $\mathcal{P}[a]$ or Q

provided the parameters satisfy the following condition:

- For every a holds $\mathcal{P}[a]$ or Q .

The scheme *Schemat11b* concerns a 0-ary predicate Q and a unary predicate \mathcal{P} , and states that:

For every a holds $\mathcal{P}[a]$ or Q

provided the parameters satisfy the following condition:

- For every a holds $\mathcal{P}[a]$ or Q .

The scheme *Schemat12a* concerns a 0-ary predicate Q and a unary predicate \mathcal{P} , and states that:

There exists a such that Q and $\mathcal{P}[a]$

provided the parameters meet the following requirement:

- Q and there exists a such that $\mathcal{P}[a]$.

The scheme *Schemat12b* concerns a 0-ary predicate Q and a unary predicate \mathcal{P} , and states that:

Q and there exists a such that $\mathcal{P}[a]$

provided the parameters have the following property:

- There exists a such that Q and $\mathcal{P}[a]$.

The scheme *Schemat13a* concerns a 0-ary predicate Q and a unary predicate \mathcal{P} , and states that:

For every a such that Q holds $\mathcal{P}[a]$

provided the parameters meet the following requirement:

- If Q , then for every a holds $\mathcal{P}[a]$.

The scheme *Schemat13b* concerns a 0-ary predicate Q and a unary predicate \mathcal{P} , and states that:

If Q , then for every a holds $\mathcal{P}[a]$

provided the following requirement is met:

- For every a such that Q holds $\mathcal{P}[a]$.

The scheme *Schemat14* concerns a 0-ary predicate Q and a unary predicate \mathcal{P} , and states that:

There exists a such that if Q , then $\mathcal{P}[a]$

provided the following requirement is met:

- If Q , then there exists a such that $\mathcal{P}[a]$.

The scheme *Schemat17* concerns a 0-ary predicate Q and a unary predicate \mathcal{P} , and states that:

If for every a holds $\mathcal{P}[a]$, then Q .

provided the parameters meet the following requirement:

- For every a such that $\mathcal{P}[a]$ holds Q .

The scheme *Schemat18a* concerns two unary predicates \mathcal{P} , Q , and states that:

There exists a such that for every b holds $\mathcal{P}[a]$ or $Q[b]$

provided the following condition is met:

- There exists a such that $\mathcal{P}[a]$ or for every b holds $Q[b]$.

The scheme *Schemat18b* concerns two unary predicates \mathcal{P} , Q , and states that:

There exists a such that $\mathcal{P}[a]$ or for every b holds $Q[b]$

provided the parameters satisfy the following condition:

- There exists a such that for every b holds $\mathcal{P}[a]$ or $Q[b]$.

The scheme *Schemat19a* concerns two unary predicates \mathcal{P} , Q , and states that:

For every b there exists a such that $\mathcal{P}[a]$ or $Q[b]$

provided the following condition is met:

- There exists a such that $\mathcal{P}[a]$ or for every b holds $Q[b]$.

The scheme *Schemat19b* concerns two unary predicates \mathcal{P} , Q , and states that:

There exists a such that $\mathcal{P}[a]$ or for every b holds $Q[b]$

provided the following requirement is met:

- For every b there exists a such that $\mathcal{P}[a]$ or $Q[b]$.

The scheme *Schemat20b* concerns two unary predicates \mathcal{P} , Q , and states that:

There exists a such that for every b holds $\mathcal{P}[a]$ or $Q[b]$

provided the following requirement is met:

- For every b there exists a such that $\mathcal{P}[a]$ or $Q[b]$.

The scheme *Schemat21a* concerns two unary predicates \mathcal{P} , Q , and states that:

There exists a such that for every b holds $\mathcal{P}[a]$ and $Q[b]$

provided the parameters have the following property:

- There exists a such that $\mathcal{P}[a]$ and for every b holds $Q[b]$.

The scheme *Schemat21b* concerns two unary predicates \mathcal{P} , Q , and states that:

There exists a such that $\mathcal{P}[a]$ and for every b holds $Q[b]$

provided the following condition is met:

- There exists a such that for every b holds $\mathcal{P}[a]$ and $Q[b]$.

The scheme *Schemat22a* concerns two unary predicates \mathcal{P} , Q , and states that:

For every b there exists a such that $\mathcal{P}[a]$ and $Q[b]$

provided the following requirement is met:

- There exists a such that $\mathcal{P}[a]$ and for every b holds $Q[b]$.

The scheme *Schemat22b* concerns two unary predicates \mathcal{P} , Q , and states that:

There exists a such that $\mathcal{P}[a]$ and for every b holds $Q[b]$

provided the following requirement is met:

- For every b there exists a such that $\mathcal{P}[a]$ and $Q[b]$.

The scheme *Schemat23b* concerns two unary predicates \mathcal{P} , Q , and states that:

There exists a such that for every b holds $\mathcal{P}[a]$ and $Q[b]$

provided the parameters meet the following requirement:

- For every b there exists a such that $\mathcal{P}[a]$ and $Q[b]$.

The scheme *Schemat24a* concerns a unary predicate Q and a binary predicate \mathcal{P} , and states that:

For every a there exists b such that if $\mathcal{P}[a, b]$, then $Q[a]$

provided the parameters meet the following requirement:

- For every a such that for every b holds $\mathcal{P}[a, b]$ holds $Q[a]$.

The scheme *Schemat24b* concerns a unary predicate Q and a binary predicate \mathcal{P} , and states that:

For every a such that for every b holds $\mathcal{P}[a, b]$ holds $Q[a]$

provided the following requirement is met:

- For every a there exists b such that if $\mathcal{P}[a, b]$, then $Q[a]$.

The scheme *Schemat25a* concerns a unary predicate Q and a binary predicate \mathcal{P} , and states that:

For all a, b such that $\mathcal{P}[a, b]$ holds $Q[a]$

provided the following condition is satisfied:

- For every a such that there exists b such that $\mathcal{P}[a, b]$ holds $Q[a]$.

The scheme *Schemat25b* concerns a unary predicate Q and a binary predicate \mathcal{P} , and states that:

For every a such that there exists b such that $\mathcal{P}[a, b]$ holds $Q[a]$

provided the parameters meet the following condition:

- For all a, b such that $\mathcal{P}[a, b]$ holds $Q[a]$.

The scheme *Schemat27* concerns a binary predicate \mathcal{P} , and states that:

For every a holds $\mathcal{P}[a, a]$

provided the parameters satisfy the following condition:

- For all a, b holds $\mathcal{P}[a, b]$.

The scheme *Schemat28* concerns a binary predicate \mathcal{P} , and states that:

There exists b such that for every a holds $\mathcal{P}[a, b]$

provided the following condition is satisfied:

- For all a, b holds $\mathcal{P}[a, b]$.

The scheme *Schemat30* concerns a binary predicate \mathcal{P} , and states that:

There exists a such that $\mathcal{P}[a, a]$

provided the parameters satisfy the following condition:

- There exists a such that for every b holds $\mathcal{P}[a, b]$.

The scheme *Schemat31* concerns a binary predicate \mathcal{P} , and states that:

For every a there exists b such that $\mathcal{P}[b, a]$

provided the following requirement is met:

- For every a holds $\mathcal{P}[a, a]$.

The scheme *Schemat33* concerns a binary predicate \mathcal{P} , and states that:

For every a there exists b such that $\mathcal{P}[a, b]$

provided the following requirement is met:

- For every a holds $\mathcal{P}[a, a]$.

The scheme *Schemat36* concerns a binary predicate \mathcal{P} , and states that:

There exist a, b such that $\mathcal{P}[a, b]$

provided the following condition is met:

- For every b there exists a such that $\mathcal{P}[a, b]$.

The scheme *Schemat37* concerns a binary predicate \mathcal{P} , and states that:

There exist a, b such that $\mathcal{P}[a, b]$

provided the parameters satisfy the following condition:

- There exists a such that $\mathcal{P}[a, a]$.

Received December 17, 1990

Published January 2, 2004
