

The Construction of SCM over Ring

Artur Kornilowicz
University of Białystok

MML Identifier: SCMRING1.

WWW: <http://mizar.org/JFM/Vol10/scmring1.html>

The articles [16], [15], [7], [23], [11], [24], [5], [6], [14], [21], [20], [12], [2], [17], [19], [3], [1], [4], [18], [22], [8], [9], [10], and [13] provide the notation and terminology for this paper.

For simplicity, we use the following convention: i, k are natural numbers, I is an element of \mathbb{Z}_8 , i_1 is an element of $\text{Instr-Loc}_{\text{SCM}}$, d_1 is an element of $\text{Data-Loc}_{\text{SCM}}$, and S is a non empty 1-sorted structure.

Let us observe that every set which is infinite is also non trivial and every 1-sorted structure which is infinite is also non trivial.

Let us mention that every non empty loop structure which is trivial is also Abelian, add-associative, right zeroed, and right complementable and every non empty double loop structure which is trivial is also right unital and right distributive.

Let us mention that every element of $\text{Data-Loc}_{\text{SCM}}$ is natural.

Let us observe that $\text{Instr}_{\text{SCM}}$ is non trivial and $\text{Instr-Loc}_{\text{SCM}}$ is infinite.

Let S be a non empty 1-sorted structure. The functor $\text{Instr}_{\text{SCM}}(S)$ yielding a subset of $[\mathbb{Z}_8, (\bigcup\{\text{the carrier of } S\} \cup \mathbb{N})^*]$ is defined by the condition (Def. 1).

(Def. 1) $\text{Instr}_{\text{SCM}}(S) = \{\langle 0, \emptyset \rangle\} \cup \{\langle I, \langle a, b \rangle \rangle; I \text{ ranges over elements of } \mathbb{Z}_8, a \text{ ranges over elements of } \text{Data-Loc}_{\text{SCM}}, b \text{ ranges over elements of } \text{Data-Loc}_{\text{SCM}}: I \in \{1, 2, 3, 4\}\} \cup \{\langle 6, \langle i \rangle \rangle : i \text{ ranges over elements of } \text{Instr-Loc}_{\text{SCM}}\} \cup \{\langle 7, \langle i, a \rangle \rangle : i \text{ ranges over elements of } \text{Instr-Loc}_{\text{SCM}}, a \text{ ranges over elements of } \text{Data-Loc}_{\text{SCM}}\} \cup \{\langle 5, \langle a, r \rangle \rangle : a \text{ ranges over elements of } \text{Data-Loc}_{\text{SCM}}, r \text{ ranges over elements of } S\}$.

Let S be a non empty 1-sorted structure. One can check that $\text{Instr}_{\text{SCM}}(S)$ is non trivial.

Let S be a non empty 1-sorted structure. We say that S is good if and only if:

(Def. 2) The carrier of $S \neq \text{Instr-Loc}_{\text{SCM}}$ and the carrier of $S \neq \text{Instr}_{\text{SCM}}(S)$.

Let us note that every non empty 1-sorted structure which is trivial is also good.

One can check that there exists a 1-sorted structure which is strict, trivial, and non empty.

Let us note that there exists a double loop structure which is strict, trivial, and non empty.

Let us observe that there exists a ring which is strict and trivial.

In the sequel G is a good non empty 1-sorted structure.

Let S be a non empty 1-sorted structure. The functor $\text{OK}_{\text{SCM}}(S)$ yields a function from \mathbb{N} into $\{\text{the carrier of } S\} \cup \{\text{Instr}_{\text{SCM}}(S), \text{Instr-Loc}_{\text{SCM}}\}$ and is defined as follows:

(Def. 3) $(\text{OK}_{\text{SCM}}(S))(0) = \text{Instr-Loc}_{\text{SCM}}$ and for every natural number k holds $(\text{OK}_{\text{SCM}}(S))(2 \cdot k + 1) = \text{the carrier of } S$ and $(\text{OK}_{\text{SCM}}(S))(2 \cdot k + 2) = \text{Instr}_{\text{SCM}}(S)$.

Let S be a non empty 1-sorted structure. An **SCM**-state over S is an element of $\prod \text{OK}_{\text{SCM}}(S)$.

Next we state several propositions:

- (1) $\text{Instr-Loc}_{\text{SCM}} \neq \text{Instr}_{\text{SCM}}(S)$.
- (2) $(\text{OK}_{\text{SCM}}(G))(i) = \text{Instr-Loc}_{\text{SCM}}$ iff $i = 0$.
- (3) $(\text{OK}_{\text{SCM}}(G))(i) = \text{the carrier of } G$ iff there exists k such that $i = 2 \cdot k + 1$.
- (4) $(\text{OK}_{\text{SCM}}(G))(i) = \text{Instr}_{\text{SCM}}(G)$ iff there exists k such that $i = 2 \cdot k + 2$.
- (5) $(\text{OK}_{\text{SCM}}(G))(d_1) = \text{the carrier of } G$.
- (6) $(\text{OK}_{\text{SCM}}(G))(i_1) = \text{Instr}_{\text{SCM}}(G)$.
- (7) $\pi_0 \prod \text{OK}_{\text{SCM}}(S) = \text{Instr-Loc}_{\text{SCM}}$.
- (8) $\pi_{d_1} \prod \text{OK}_{\text{SCM}}(G) = \text{the carrier of } G$.
- (9) $\pi_{i_1} \prod \text{OK}_{\text{SCM}}(G) = \text{Instr}_{\text{SCM}}(G)$.

Let S be a non empty 1-sorted structure and let s be an **SCM**-state over S . The functor \mathbf{IC}_s yielding an element of $\text{Instr-Loc}_{\text{SCM}}$ is defined as follows:

(Def. 4) $\mathbf{IC}_s = s(0)$.

Let R be a good non empty 1-sorted structure, let s be an **SCM**-state over R , and let u be an element of $\text{Instr-Loc}_{\text{SCM}}$. The functor $\text{Chg}_{\text{SCM}}(s, u)$ yielding an **SCM**-state over R is defined by:

(Def. 5) $\text{Chg}_{\text{SCM}}(s, u) = s + \cdot (0 \dashrightarrow u)$.

Next we state three propositions:

- (10) For every **SCM**-state s over G and for every element u of $\text{Instr-Loc}_{\text{SCM}}$ holds $(\text{Chg}_{\text{SCM}}(s, u))(0) = u$.
- (11) For every **SCM**-state s over G and for every element u of $\text{Instr-Loc}_{\text{SCM}}$ and for every element m_1 of $\text{Data-Loc}_{\text{SCM}}$ holds $(\text{Chg}_{\text{SCM}}(s, u))(m_1) = s(m_1)$.
- (12) For every **SCM**-state s over G and for all elements u, v of $\text{Instr-Loc}_{\text{SCM}}$ holds $(\text{Chg}_{\text{SCM}}(s, u))(v) = s(v)$.

Let R be a good non empty 1-sorted structure, let s be an **SCM**-state over R , let t be an element of $\text{Data-Loc}_{\text{SCM}}$, and let u be an element of R . The functor $\text{Chg}_{\text{SCM}}(s, t, u)$ yields an **SCM**-state over R and is defined by:

(Def. 6) $\text{Chg}_{\text{SCM}}(s, t, u) = s + \cdot (t \dashrightarrow u)$.

Next we state four propositions:

- (13) For every **SCM**-state s over G and for every element t of $\text{Data-Loc}_{\text{SCM}}$ and for every element u of G holds $(\text{Chg}_{\text{SCM}}(s, t, u))(0) = s(0)$.
- (14) For every **SCM**-state s over G and for every element t of $\text{Data-Loc}_{\text{SCM}}$ and for every element u of G holds $(\text{Chg}_{\text{SCM}}(s, t, u))(t) = u$.
- (15) Let s be an **SCM**-state over G , t be an element of $\text{Data-Loc}_{\text{SCM}}$, u be an element of G , and m_1 be an element of $\text{Data-Loc}_{\text{SCM}}$. If $m_1 \neq t$, then $(\text{Chg}_{\text{SCM}}(s, t, u))(m_1) = s(m_1)$.
- (16) Let s be an **SCM**-state over G , t be an element of $\text{Data-Loc}_{\text{SCM}}$, u be an element of G , and v be an element of $\text{Instr-Loc}_{\text{SCM}}$. Then $(\text{Chg}_{\text{SCM}}(s, t, u))(v) = s(v)$.

Let R be a good non empty 1-sorted structure, let s be an **SCM**-state over R , and let a be an element of $\text{Data-Loc}_{\text{SCM}}$. Then $s(a)$ is an element of R .

Let S be a non empty 1-sorted structure and let x be an element of $\text{Instr}_{\text{SCM}}(S)$. Let us assume that there exist elements m_1, m_2 of $\text{Data-Loc}_{\text{SCM}}$ and I such that $x = \langle I, \langle m_1, m_2 \rangle \rangle$. The functor $x \text{address}_1$ yields an element of $\text{Data-Loc}_{\text{SCM}}$ and is defined by:

(Def. 7) There exists a finite sequence f of elements of $\text{Data-Loc}_{\text{SCM}}$ such that $f = x_2$ and $x\text{address}_1 = f_1$.

The functor $x\text{address}_2$ yields an element of $\text{Data-Loc}_{\text{SCM}}$ and is defined as follows:

(Def. 8) There exists a finite sequence f of elements of $\text{Data-Loc}_{\text{SCM}}$ such that $f = x_2$ and $x\text{address}_2 = f_2$.

Next we state the proposition

(17) For every element x of $\text{Instr}_{\text{SCM}}(S)$ and for all elements m_1, m_2 of $\text{Data-Loc}_{\text{SCM}}$ such that $x = \langle I, \langle m_1, m_2 \rangle \rangle$ holds $x\text{address}_1 = m_1$ and $x\text{address}_2 = m_2$.

Let R be a non empty 1-sorted structure and let x be an element of $\text{Instr}_{\text{SCM}}(R)$. Let us assume that there exist an element m_1 of $\text{Instr-Loc}_{\text{SCM}}$ and I such that $x = \langle I, \langle m_1 \rangle \rangle$. The functor $x\text{address}_j$ yields an element of $\text{Instr-Loc}_{\text{SCM}}$ and is defined as follows:

(Def. 9) There exists a finite sequence f of elements of $\text{Instr-Loc}_{\text{SCM}}$ such that $f = x_2$ and $x\text{address}_j = f_1$.

Next we state the proposition

(18) For every element x of $\text{Instr}_{\text{SCM}}(S)$ and for every element m_1 of $\text{Instr-Loc}_{\text{SCM}}$ such that $x = \langle I, \langle m_1 \rangle \rangle$ holds $x\text{address}_j = m_1$.

Let S be a non empty 1-sorted structure and let x be an element of $\text{Instr}_{\text{SCM}}(S)$. Let us assume that there exist an element m_1 of $\text{Instr-Loc}_{\text{SCM}}$, an element m_2 of $\text{Data-Loc}_{\text{SCM}}$, and I such that $x = \langle I, \langle m_1, m_2 \rangle \rangle$. The functor $x\text{address}_j$ yields an element of $\text{Instr-Loc}_{\text{SCM}}$ and is defined by:

(Def. 10) There exists an element m_1 of $\text{Instr-Loc}_{\text{SCM}}$ and there exists an element m_2 of $\text{Data-Loc}_{\text{SCM}}$ such that $\langle m_1, m_2 \rangle = x_2$ and $x\text{address}_j = \langle m_1, m_2 \rangle_1$.

The functor $x\text{address}_c$ yields an element of $\text{Data-Loc}_{\text{SCM}}$ and is defined as follows:

(Def. 11) There exists an element m_1 of $\text{Instr-Loc}_{\text{SCM}}$ and there exists an element m_2 of $\text{Data-Loc}_{\text{SCM}}$ such that $\langle m_1, m_2 \rangle = x_2$ and $x\text{address}_c = \langle m_1, m_2 \rangle_2$.

We now state the proposition

(19) Let x be an element of $\text{Instr}_{\text{SCM}}(S)$, m_1 be an element of $\text{Instr-Loc}_{\text{SCM}}$, and m_2 be an element of $\text{Data-Loc}_{\text{SCM}}$. If $x = \langle I, \langle m_1, m_2 \rangle \rangle$, then $x\text{address}_j = m_1$ and $x\text{address}_c = m_2$.

Let S be a non empty 1-sorted structure, let d be an element of $\text{Data-Loc}_{\text{SCM}}$, and let s be an element of S . Then $\langle d, s \rangle$ is a finite sequence of elements of $\text{Data-Loc}_{\text{SCM}} \cup$ the carrier of S .

Let S be a non empty 1-sorted structure and let x be an element of $\text{Instr}_{\text{SCM}}(S)$. Let us assume that there exist an element m_1 of $\text{Data-Loc}_{\text{SCM}}$, an element r of S , and I such that $x = \langle I, \langle m_1, r \rangle \rangle$. The functor $x\text{const_address}$ yielding an element of $\text{Data-Loc}_{\text{SCM}}$ is defined as follows:

(Def. 12) There exists a finite sequence f of elements of $\text{Data-Loc}_{\text{SCM}} \cup$ the carrier of S such that $f = x_2$ and $x\text{const_address} = f_1$.

The functor $x\text{const_value}$ yielding an element of S is defined by:

(Def. 13) There exists a finite sequence f of elements of $\text{Data-Loc}_{\text{SCM}} \cup$ the carrier of S such that $f = x_2$ and $x\text{const_value} = f_2$.

Next we state the proposition

(20) Let x be an element of $\text{Instr}_{\text{SCM}}(S)$, m_1 be an element of $\text{Data-Loc}_{\text{SCM}}$, and r be an element of S . If $x = \langle I, \langle m_1, r \rangle \rangle$, then $x\text{const_address} = m_1$ and $x\text{const_value} = r$.

Let R be a good ring, let x be an element of $\text{Instr}_{\text{SCM}}(R)$, and let s be an **SCM**-state over R . The functor $\text{Exec-Res}_{\text{SCM}}(x, s)$ yielding an **SCM**-state over R is defined as follows:

$$(\text{Def. 14}) \quad \text{Exec-Ress}_{\text{SCM}}(x, s) = \begin{cases} \text{Chg}_{\text{SCM}}(\text{Chg}_{\text{SCM}}(s, x \text{ address}_1, s(x \text{ address}_2)), \text{Next}(\mathbf{IC}_s)), & \text{if there exist elements } m_1, m_2 \text{ of } \\ \text{Chg}_{\text{SCM}}(\text{Chg}_{\text{SCM}}(s, x \text{ address}_1, s(x \text{ address}_1) + s(x \text{ address}_2)), \text{Next}(\mathbf{IC}_s)), & \text{if there exist elements } \\ \text{Chg}_{\text{SCM}}(\text{Chg}_{\text{SCM}}(s, x \text{ address}_1, s(x \text{ address}_1) - s(x \text{ address}_2)), \text{Next}(\mathbf{IC}_s)), & \text{if there exist elements } \\ \text{Chg}_{\text{SCM}}(\text{Chg}_{\text{SCM}}(s, x \text{ address}_1, s(x \text{ address}_1) \cdot s(x \text{ address}_2)), \text{Next}(\mathbf{IC}_s)), & \text{if there exist elements } \\ \text{Chg}_{\text{SCM}}(s, x \text{ address}_j), & \text{if there exists an element } m_1 \text{ of } \text{Instr-Loc}_{\text{SCM}} \text{ such that } x = \langle 6, \langle m_1 \rangle \rangle, \\ \text{Chg}_{\text{SCM}}(s, (s(x \text{ address}_c) = 0_R \rightarrow x \text{ address}_j, \text{Next}(\mathbf{IC}_s))), & \text{if there exists an element } m_1 \text{ of } \text{Instr-Loc}_{\text{SCM}} \\ \text{Chg}_{\text{SCM}}(\text{Chg}_{\text{SCM}}(s, x \text{ const_address}, x \text{ const_value}), \text{Next}(\mathbf{IC}_s)), & \text{if there exists an element } m_1 \text{ of } \text{Instr-Loc}_{\text{SCM}} \\ s, & \text{otherwise.} \end{cases}$$

Let S be a non empty 1-sorted structure, let f be a function from $\text{Instr}_{\text{SCM}}(S)$ into $(\prod \text{OK}_{\text{SCM}}(S))^{\prod \text{OK}_{\text{SCM}}(S)}$, and let x be an element of $\text{Instr}_{\text{SCM}}(S)$. One can verify that $f(x)$ is function-like and relation-like.

Let R be a good ring. The functor $\text{Exec}_{\text{SCM}}(R)$ yielding a function from $\text{Instr}_{\text{SCM}}(R)$ into $(\prod \text{OK}_{\text{SCM}}(R))^{\prod \text{OK}_{\text{SCM}}(R)}$ is defined as follows:

(Def. 15) For every element x of $\text{Instr}_{\text{SCM}}(R)$ and for every **SCM**-state y over R holds $(\text{Exec}_{\text{SCM}}(R))(x)(y) = \text{Exec-Ress}_{\text{SCM}}(x, y)$.

REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/nat_1.html.
- [2] Grzegorz Bancerek. Sequences of ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/ordinal2.html>.
- [3] Grzegorz Bancerek. König's theorem. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/card_3.html.
- [4] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.
- [5] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [6] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [7] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [8] Czesław Byliński. A classical first order language. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/cqc_lang.html.
- [9] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/funct_4.html.
- [10] Czesław Byliński. Subcategories and products of categories. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/cat_2.html.
- [11] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.
- [12] Eugeniusz Kusak, Wojciech Leoniczuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/vectsp_1.html.
- [13] Yatsuka Nakamura and Andrzej Trybulec. On a mathematical model of programs. *Journal of Formalized Mathematics*, 4, 1992. http://mizar.org/JFM/Vol4/ami_2.html.
- [14] Dariusz Surowik. Cyclic groups and some of their properties — part I. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/gr_cy_1.html.
- [15] Andrzej Trybulec. Enumerated sets. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/enumset1.html>.
- [16] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [17] Andrzej Trybulec. Tuples, projections and Cartesian products. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/mcart_1.html.
- [18] Andrzej Trybulec. Function domains and Fränkel operator. *Journal of Formalized Mathematics*, 2, 1990. <http://mizar.org/JFM/Vol2/fraenkel.html>.
- [19] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [20] Wojciech A. Trybulec. Vectors in real linear space. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/rlvect_1.html.

- [21] Wojciech A. Trybulec. Groups. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/group_1.html.
- [22] Wojciech A. Trybulec. Pigeon hole principle. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_4.html.
- [23] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [24] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received November 29, 1998

Published January 2, 2004
