

Some Properties of Special Polygonal Curves

Andrzej Trybulec
University of Białystok

Yatsuka Nakamura
Shinshu University
Nagano

Summary. In the paper some auxiliary theorems are proved, needed in the proof of the second part of the Jordan curve theorem for special polygons. They deal mostly with characteristic points of plane non empty compacts introduced in [9], operation *mid* introduced in [22] and the predicate “*f* is in the area of *g*” (*f* and *g* : finite sequences of points of the plane) introduced in [30].

MML Identifier: SPRECT_3.

WWW: http://mizar.org/JFM/Vol10/sprect_3.html

The articles [28], [27], [7], [33], [3], [14], [2], [24], [1], [6], [4], [32], [8], [15], [16], [17], [26], [34], [10], [25], [5], [11], [12], [9], [18], [19], [20], [23], [29], [22], [13], [21], [31], and [30] provide the notation and terminology for this paper.

1. PRELIMINARIES

In this paper *i, j, k, n* are natural numbers.

One can prove the following propositions:

- (2)¹ For all sets *A, B, C, p* such that $A \subseteq B$ and $B \cap C = \{p\}$ and $p \in A$ holds $A \cap C = \{p\}$.
- (3) For all real numbers *q, r, s, t* such that $t \geq 0$ and $t \leq 1$ and $s = (1 - t) \cdot q + t \cdot r$ and $q \leq s$ and $r < s$ holds $t = 0$.
- (4) For all real numbers *q, r, s, t* such that $t \geq 0$ and $t \leq 1$ and $s = (1 - t) \cdot q + t \cdot r$ and $q \geq s$ and $r > s$ holds $t = 0$.
- (5) If $i -' k \leq j$, then $i \leq j + k$.
- (6) If $i \leq j + k$, then $i -' k \leq j$.
- (7) If $i \leq j -' k$ and $k \leq j$, then $i + k \leq j$.
- (8) If $j + k \leq i$, then $k \leq i -' j$.
- (9) If $k \leq i$ and $i < j$, then $i -' k < j -' k$.
- (10) If $i < j$ and $k < j$, then $i -' k < j -' k$.
- (11) Let *D* be a non empty set, *f* be a non empty finite sequence of elements of *D*, and *g* be a finite sequence of elements of *D*. Then $(g \hat{\ } f)_{\text{len}(g \hat{\ } f)} = f_{\text{len} f}$.
- (12) For all sets *a, b, c, d* holds the indices of $\begin{pmatrix} a & b \\ c & d \end{pmatrix} = \{\langle 1, 1 \rangle, \langle 1, 2 \rangle, \langle 2, 1 \rangle, \langle 2, 2 \rangle\}$.

¹ The proposition (1) has been removed.

2. EUCLIDEAN SPACE

The following four propositions are true:

- (13) For all points p, q of \mathcal{E}_T^n and for every real number r such that $0 < r$ and $p = (1-r) \cdot p + r \cdot q$ holds $p = q$.
- (14) For all points p, q of \mathcal{E}_T^n and for every real number r such that $r < 1$ and $p = (1-r) \cdot q + r \cdot p$ holds $p = q$.
- (15) For all points p, q of \mathcal{E}_T^n such that $p = \frac{1}{2} \cdot (p+q)$ holds $p = q$.
- (16) For all points p, q, r of \mathcal{E}_T^n such that $q \in \mathcal{L}(p, r)$ and $r \in \mathcal{L}(p, q)$ holds $q = r$.

3. EUCLIDEAN PLANE

We now state several propositions:

- (17) Let A be a non empty subset of \mathcal{E}_T^2 , p be an element of \mathcal{E}^2 , and r be a real number. If $A = \text{Ball}(p, r)$, then A is connected.
- (18) For all subsets A, B of \mathcal{E}_T^2 such that A is open and B is a component of A holds B is open.
- (19) For all points p, q, r of \mathcal{E}_T^2 such that $\mathcal{L}(p, q)$ is horizontal and $r \in \mathcal{L}(p, q)$ holds $p_2 = r_2$.
- (20) For all points p, q, r of \mathcal{E}_T^2 such that $\mathcal{L}(p, q)$ is vertical and $r \in \mathcal{L}(p, q)$ holds $p_1 = r_1$.
- (21) For all points p, q, r, s of \mathcal{E}_T^2 such that $\mathcal{L}(p, q)$ is horizontal and $\mathcal{L}(r, s)$ is horizontal and $\mathcal{L}(p, q)$ meets $\mathcal{L}(r, s)$ holds $p_2 = r_2$.
- (22) For all points p, q, r of \mathcal{E}_T^2 such that $\mathcal{L}(p, q)$ is vertical and $\mathcal{L}(q, r)$ is horizontal holds $\mathcal{L}(p, q) \cap \mathcal{L}(q, r) = \{q\}$.
- (23) For all points p, q, r, s of \mathcal{E}_T^2 such that $\mathcal{L}(p, q)$ is horizontal and $\mathcal{L}(s, r)$ is vertical and $r \in \mathcal{L}(p, q)$ holds $\mathcal{L}(p, q) \cap \mathcal{L}(s, r) = \{r\}$.

4. MISCELLANEOUS

In the sequel p, q denote points of \mathcal{E}_T^2 and G denotes a Go-board.

We now state two propositions:

- (24) If $1 \leq j$ and $j \leq k$ and $k \leq \text{width } G$ and $1 \leq i$ and $i \leq \text{len } G$, then $(G \circ (i, j))_2 \leq (G \circ (i, k))_2$.
- (25) If $1 \leq j$ and $j \leq \text{width } G$ and $1 \leq i$ and $i \leq k$ and $k \leq \text{len } G$, then $(G \circ (i, j))_1 \leq (G \circ (i, k))_1$.

In the sequel C denotes a subset of \mathcal{E}_T^2 .

One can prove the following propositions:

- (26) $\mathcal{L}(\text{NW-corner}(C), \text{NE-corner}(C)) \subseteq \tilde{\mathcal{L}}(\text{SpStSeq } C)$.
- (27) $N_{\text{most}}(C) \subseteq \mathcal{L}(\text{NW-corner}(C), \text{NE-corner}(C))$.
- (28) For every non empty compact subset C of \mathcal{E}_T^2 holds $N_{\text{min}}(C) \in \mathcal{L}(\text{NW-corner}(C), \text{NE-corner}(C))$.
- (29) $\mathcal{L}(\text{NW-corner}(C), \text{NE-corner}(C))$ is horizontal.
- (31)² Let g be a finite sequence of elements of \mathcal{E}_T^2 and p be a point of \mathcal{E}_T^2 . Suppose $g_1 \neq p$ and $(g_1)_1 = p_1$ or $(g_1)_2 = p_2$ and g is a special sequence and $\mathcal{L}(p, g_1) \cap \tilde{\mathcal{L}}(g) = \{g_1\}$. Then $\langle p \rangle \hat{\ } g$ is a special sequence.

² The proposition (30) has been removed.

- (33)³ Let f be a S-sequence in \mathbb{R}^2 and p be a point of \mathcal{E}_T^2 . If $1 < j$ and $j \leq \text{len } f$ and $p \in \tilde{\mathcal{L}}(\text{mid}(f, 1, j))$, then LE $p, f_j, \tilde{\mathcal{L}}(f), f_1, f_{\text{len } f}$.
- (34) For every finite sequence h of elements of \mathcal{E}_T^2 such that $i \in \text{dom } h$ and $j \in \text{dom } h$ holds $\tilde{\mathcal{L}}(\text{mid}(h, i, j)) \subseteq \tilde{\mathcal{L}}(h)$.
- (35) If $1 \leq i$ and $i < j$, then for every finite sequence f of elements of \mathcal{E}_T^2 such that $j \leq \text{len } f$ holds $\tilde{\mathcal{L}}(\text{mid}(f, i, j)) = \mathcal{L}(f, i) \cup \tilde{\mathcal{L}}(\text{mid}(f, i+1, j))$.
- (36) Let f be a finite sequence of elements of \mathcal{E}_T^2 . If $1 \leq i$, then if $i < j$ and $j \leq \text{len } f$, then $\tilde{\mathcal{L}}(\text{mid}(f, i, j)) = \tilde{\mathcal{L}}(\text{mid}(f, i, j-1)) \cup \mathcal{L}(f, j-1)$.
- (38)⁴ Let f, g be finite sequences of elements of \mathcal{E}_T^2 . Suppose that
- (i) f is a special sequence,
 - (ii) g is a special sequence,
 - (iii) $(f_{\text{len } f})_1 = (g_1)_1$ or $(f_{\text{len } f})_2 = (g_1)_2$,
 - (iv) $\tilde{\mathcal{L}}(f)$ misses $\tilde{\mathcal{L}}(g)$,
 - (v) $\mathcal{L}(f_{\text{len } f}, g_1) \cap \tilde{\mathcal{L}}(f) = \{f_{\text{len } f}\}$, and
 - (vi) $\mathcal{L}(f_{\text{len } f}, g_1) \cap \tilde{\mathcal{L}}(g) = \{g_1\}$.
- Then $f \wedge g$ is a special sequence.
- (39) For every S-sequence f in \mathbb{R}^2 and for every point p of \mathcal{E}_T^2 such that $p \in \tilde{\mathcal{L}}(f)$ holds $(\perp f, p)_1 = f_1$.
- (40) Let f be a S-sequence in \mathbb{R}^2 and p, q be points of \mathcal{E}_T^2 . If $1 \leq j$ and $j < \text{len } f$ and $p \in \mathcal{L}(f, j)$ and $q \in \mathcal{L}(f_j, p)$, then LE $q, p, \tilde{\mathcal{L}}(f), f_1, f_{\text{len } f}$.

5. SPECIAL CIRCULAR SEQUENCES

The following proposition is true

- (41) For every non constant standard special circular sequence f holds $\text{LeftComp}(f)$ is open and $\text{RightComp}(f)$ is open.

Let f be a non constant standard special circular sequence. One can verify the following observations:

- * $\tilde{\mathcal{L}}(f)$ is non vertical and non horizontal,
- * $\text{LeftComp}(f)$ is region, and
- * $\text{RightComp}(f)$ is region.

One can prove the following propositions:

- (42) For every non constant standard special circular sequence f holds $\text{RightComp}(f)$ misses $\tilde{\mathcal{L}}(f)$.
- (43) For every non constant standard special circular sequence f holds $\text{LeftComp}(f)$ misses $\tilde{\mathcal{L}}(f)$.
- (44) For every non constant standard special circular sequence f holds $i_{\text{WN}} f < i_{\text{EN}} f$.
- (45) Let f be a non constant standard special circular sequence. Then there exists i such that $1 \leq i$ and $i < \text{len } f$ the Go-board of f and $N_{\min}(\tilde{\mathcal{L}}(f)) =$ the Go-board of $f \circ (i, \text{width the Go-board of } f)$.

³ The proposition (32) has been removed.

⁴ The proposition (37) has been removed.

- (46) Let f be a clockwise oriented non constant standard special circular sequence. Suppose $i \in \text{dom}$ the Go-board of f and $f_1 =$ the Go-board of $f \circ (i, \text{width the Go-board of } f)$ and $f_1 = N_{\min}(\tilde{\mathcal{L}}(f))$. Then $f_2 =$ the Go-board of $f \circ (i+1, \text{width the Go-board of } f)$ and $f_{\text{len } f - 1} =$ the Go-board of $f \circ (i, \text{width the Go-board of } f - 1)$.
- (47) Let f be a non constant standard special circular sequence. If $1 \leq i$ and $i < j$ and $j \leq \text{len } f$ and $f_1 \in \tilde{\mathcal{L}}(\text{mid}(f, i, j))$, then $i = 1$ or $j = \text{len } f$.
- (48) Let f be a clockwise oriented non constant standard special circular sequence. If $f_1 = N_{\min}(\tilde{\mathcal{L}}(f))$, then $\mathcal{L}(f_1, f_2) \subseteq \tilde{\mathcal{L}}(\text{SpStSeq } \tilde{\mathcal{L}}(f))$.

6. RECTANGULAR SEQUENCES

Next we state the proposition

- (49) Let f be a rectangular finite sequence of elements of \mathcal{E}_T^2 and p be a point of \mathcal{E}_T^2 . If $p \in \tilde{\mathcal{L}}(f)$, then $p_1 = \text{W-bound}(\tilde{\mathcal{L}}(f))$ or $p_1 = \text{E-bound}(\tilde{\mathcal{L}}(f))$ or $p_2 = \text{S-bound}(\tilde{\mathcal{L}}(f))$ or $p_2 = \text{N-bound}(\tilde{\mathcal{L}}(f))$.

Let us observe that there exists a special circular sequence which is rectangular.

One can prove the following propositions:

- (50) Let f be a rectangular special circular sequence and g be a S-sequence in \mathbb{R}^2 . If $g_1 \in \text{LeftComp}(f)$ and $g_{\text{len } g} \in \text{RightComp}(f)$, then $\tilde{\mathcal{L}}(f)$ meets $\tilde{\mathcal{L}}(g)$.
- (51) For every rectangular special circular sequence f holds $\text{SpStSeq } \tilde{\mathcal{L}}(f) = f$.
- (52) Let f be a rectangular special circular sequence. Then $\tilde{\mathcal{L}}(f) = \{p; p \text{ ranges over points of } \mathcal{E}_T^2: p_1 = \text{W-bound}(\tilde{\mathcal{L}}(f)) \wedge p_2 \leq \text{N-bound}(\tilde{\mathcal{L}}(f)) \wedge p_2 \geq \text{S-bound}(\tilde{\mathcal{L}}(f)) \vee p_1 \leq \text{E-bound}(\tilde{\mathcal{L}}(f)) \wedge p_1 \geq \text{W-bound}(\tilde{\mathcal{L}}(f)) \wedge p_2 = \text{N-bound}(\tilde{\mathcal{L}}(f)) \vee p_1 \leq \text{E-bound}(\tilde{\mathcal{L}}(f)) \wedge p_1 \geq \text{W-bound}(\tilde{\mathcal{L}}(f)) \wedge p_2 = \text{S-bound}(\tilde{\mathcal{L}}(f)) \vee p_1 = \text{E-bound}(\tilde{\mathcal{L}}(f)) \wedge p_2 \leq \text{N-bound}(\tilde{\mathcal{L}}(f)) \wedge p_2 \geq \text{S-bound}(\tilde{\mathcal{L}}(f))\}$.
- (53) For every rectangular special circular sequence f holds the Go-board of $f = \begin{pmatrix} f_4 & f_1 \\ f_3 & f_2 \end{pmatrix}$.
- (54) Let f be a rectangular special circular sequence. Then $\text{LeftComp}(f) = \{p : \text{W-bound}(\tilde{\mathcal{L}}(f)) \not\leq p_1 \vee p_1 \not\leq \text{E-bound}(\tilde{\mathcal{L}}(f)) \vee \text{S-bound}(\tilde{\mathcal{L}}(f)) \not\leq p_2 \vee p_2 \not\leq \text{N-bound}(\tilde{\mathcal{L}}(f))\}$ and $\text{RightComp}(f) = \{q : \text{W-bound}(\tilde{\mathcal{L}}(f)) < q_1 \wedge q_1 < \text{E-bound}(\tilde{\mathcal{L}}(f)) \wedge \text{S-bound}(\tilde{\mathcal{L}}(f)) < q_2 \wedge q_2 < \text{N-bound}(\tilde{\mathcal{L}}(f))\}$.

Let us mention that there exists a rectangular special circular sequence which is clockwise oriented.

Let us mention that every rectangular special circular sequence is clockwise oriented.

Next we state four propositions:

- (55) Let f be a rectangular special circular sequence and g be a S-sequence in \mathbb{R}^2 . If $g_1 \in \text{LeftComp}(f)$ and $g_{\text{len } g} \in \text{RightComp}(f)$, then $\text{LPoint}(\tilde{\mathcal{L}}(g), g_1, g_{\text{len } g}, \tilde{\mathcal{L}}(f)) \neq \text{NW-corner}(\tilde{\mathcal{L}}(f))$.
- (56) Let f be a rectangular special circular sequence and g be a S-sequence in \mathbb{R}^2 . If $g_1 \in \text{LeftComp}(f)$ and $g_{\text{len } g} \in \text{RightComp}(f)$, then $\text{LPoint}(\tilde{\mathcal{L}}(g), g_1, g_{\text{len } g}, \tilde{\mathcal{L}}(f)) \neq \text{SE-corner}(\tilde{\mathcal{L}}(f))$.
- (57) Let f be a rectangular special circular sequence and p be a point of \mathcal{E}_T^2 . If $\text{W-bound}(\tilde{\mathcal{L}}(f)) > p_1$ or $p_1 > \text{E-bound}(\tilde{\mathcal{L}}(f))$ or $\text{S-bound}(\tilde{\mathcal{L}}(f)) > p_2$ or $p_2 > \text{N-bound}(\tilde{\mathcal{L}}(f))$, then $p \in \text{LeftComp}(f)$.
- (58) For every clockwise oriented non constant standard special circular sequence f such that $f_1 = N_{\min}(\tilde{\mathcal{L}}(f))$ holds $\text{LeftComp}(\text{SpStSeq } \tilde{\mathcal{L}}(f)) \subseteq \text{LeftComp}(f)$.

7. IN THE AREA

The following propositions are true:

- (59) Let f be a finite sequence of elements of \mathcal{E}_T^2 and p, q be points of \mathcal{E}_T^2 . Then $\langle p, q \rangle$ is in the area of f if and only if $\langle p \rangle$ is in the area of f and $\langle q \rangle$ is in the area of f .
- (60) Let f be a rectangular finite sequence of elements of \mathcal{E}_T^2 and p be a point of \mathcal{E}_T^2 . Suppose $\langle p \rangle$ is in the area of f but $p_1 = \text{W-bound}(\tilde{\mathcal{L}}(f))$ or $p_1 = \text{E-bound}(\tilde{\mathcal{L}}(f))$ or $p_2 = \text{S-bound}(\tilde{\mathcal{L}}(f))$ or $p_2 = \text{N-bound}(\tilde{\mathcal{L}}(f))$. Then $p \in \tilde{\mathcal{L}}(f)$.
- (61) Let f be a finite sequence of elements of \mathcal{E}_T^2 , p, q be points of \mathcal{E}_T^2 , and r be a real number. Suppose $0 \leq r$ and $r \leq 1$ and $\langle p, q \rangle$ is in the area of f . Then $\langle (1-r) \cdot p + r \cdot q \rangle$ is in the area of f .
- (62) Let f, g be finite sequences of elements of \mathcal{E}_T^2 . If g is in the area of f and $i \in \text{dom } g$, then $\langle g_i \rangle$ is in the area of f .
- (63) Let f, g be finite sequences of elements of \mathcal{E}_T^2 and p be a point of \mathcal{E}_T^2 . If g is in the area of f and $p \in \tilde{\mathcal{L}}(g)$, then $\langle p \rangle$ is in the area of f .
- (64) Let f be a rectangular finite sequence of elements of \mathcal{E}_T^2 and p, q be points of \mathcal{E}_T^2 . If $q \notin \tilde{\mathcal{L}}(f)$ and $\langle p, q \rangle$ is in the area of f , then $\mathcal{L}(p, q) \cap \tilde{\mathcal{L}}(f) \subseteq \{p\}$.
- (65) Let f be a rectangular finite sequence of elements of \mathcal{E}_T^2 and p, q be points of \mathcal{E}_T^2 . If $p \in \tilde{\mathcal{L}}(f)$ and $q \notin \tilde{\mathcal{L}}(f)$ and $\langle q \rangle$ is in the area of f , then $\mathcal{L}(p, q) \cap \tilde{\mathcal{L}}(f) = \{p\}$.
- (66) Let f be a non constant standard special circular sequence. Suppose $1 \leq i$ and $i \leq \text{len the Go-board of } f$ and $1 \leq j$ and $j \leq \text{width the Go-board of } f$. Then $\langle \text{the Go-board of } f \circ (i, j) \rangle$ is in the area of f .
- (67) Let g be a finite sequence of elements of \mathcal{E}_T^2 and p, q be points of \mathcal{E}_T^2 . If $\langle p, q \rangle$ is in the area of g , then $\langle \frac{1}{2} \cdot (p + q) \rangle$ is in the area of g .
- (68) For all finite sequences f, g of elements of \mathcal{E}_T^2 such that g is in the area of f holds $\text{Rev}(g)$ is in the area of f .
- (69) Let f, g be finite sequences of elements of \mathcal{E}_T^2 and p be a point of \mathcal{E}_T^2 . Suppose g is in the area of f and $\langle p \rangle$ is in the area of f and g is a special sequence and $p \in \tilde{\mathcal{L}}(g)$. Then $\downarrow g, p$ is in the area of f .
- (70) Let f be a non constant standard special circular sequence and g be a finite sequence of elements of \mathcal{E}_T^2 . Then g is in the area of f if and only if g is in the area of $\text{SpStSeq } \tilde{\mathcal{L}}(f)$.
- (71) Let f be a rectangular special circular sequence and g be a S-sequence in \mathbb{R}^2 . If $g_1 \in \text{LeftComp}(f)$ and $g_{\text{len } g} \in \text{RightComp}(f)$, then $\downarrow \text{LPoint}(\tilde{\mathcal{L}}(g), g_1, g_{\text{len } g}, \tilde{\mathcal{L}}(f)), g$ is in the area of f .
- (72) Let f be a non constant standard special circular sequence. Suppose $1 \leq i$ and $i < \text{len the Go-board of } f$ and $1 \leq j$ and $j < \text{width the Go-board of } f$. Then $\text{Intcell}(\text{the Go-board of } f, i, j)$ misses $\tilde{\mathcal{L}}(\text{SpStSeq } \tilde{\mathcal{L}}(f))$.
- (73) Let f, g be finite sequences of elements of \mathcal{E}_T^2 and p be a point of \mathcal{E}_T^2 . Suppose g is in the area of f and $\langle p \rangle$ is in the area of f and g is a special sequence and $p \in \tilde{\mathcal{L}}(g)$. Then $\downarrow p, g$ is in the area of f .

REFERENCES

- [1] Grzegorz Bancerek. Cardinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/card_1.html.
- [2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/nat_1.html.
- [3] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Voll/ordinal1.html>.
- [4] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/finseq_1.html.
- [5] Leszek Borys. Paracompact and metrizable spaces. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/pcomps_1.html.
- [6] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/funct_1.html.
- [7] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/zfmisc_1.html.
- [8] Czesław Byliński. Some properties of restrictions of finite sequences. *Journal of Formalized Mathematics*, 7, 1995. http://mizar.org/JFM/Vol7/finseq_5.html.
- [9] Czesław Byliński and Piotr Rudnicki. Bounding boxes for compact sets in \mathcal{E}^2 . *Journal of Formalized Mathematics*, 9, 1997. http://mizar.org/JFM/Vol9/pscomp_1.html.
- [10] Agata Darmochwał. Compact spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/compts_1.html.
- [11] Agata Darmochwał. The Euclidean space. *Journal of Formalized Mathematics*, 3, 1991. <http://mizar.org/JFM/Vol3/euclid.html>.
- [12] Agata Darmochwał and Yatsuka Nakamura. The topological space \mathcal{E}_T^2 . Arcs, line segments and special polygonal arcs. *Journal of Formalized Mathematics*, 3, 1991. <http://mizar.org/JFM/Vol3/topreall.html>.
- [13] Adam Grabowski and Yatsuka Nakamura. The ordering of points on a curve. Part II. *Journal of Formalized Mathematics*, 9, 1997. <http://mizar.org/JFM/Vol9/jordan5c.html>.
- [14] Krzysztof Hryniewiecki. Basic properties of real numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/real_1.html.
- [15] Katarzyna Jankowska. Matrices. Abelian group of matrices. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/matrix_1.html.
- [16] Katarzyna Jankowska. Transpose matrices and groups of permutations. *Journal of Formalized Mathematics*, 4, 1992. http://mizar.org/JFM/Vol4/matrix_2.html.
- [17] Stanisława Kanas, Adam Lecko, and Mariusz Startek. Metric spaces. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/metric_1.html.
- [18] Jarosław Kotowicz and Yatsuka Nakamura. Introduction to Go-Board — part I. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/goboard1.html>.
- [19] Jarosław Kotowicz and Yatsuka Nakamura. Introduction to Go-Board — part II. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/goboard2.html>.
- [20] Yatsuka Nakamura and Czesław Byliński. Extremal properties of vertices on special polygons, part I. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/sppol_1.html.
- [21] Yatsuka Nakamura and Adam Grabowski. Bounding boxes for special sequences in \mathcal{E}^2 . *Journal of Formalized Mathematics*, 10, 1998. <http://mizar.org/JFM/Vol10/jordan5d.html>.
- [22] Yatsuka Nakamura and Roman Matuszewski. Reconstructions of special sequences. *Journal of Formalized Mathematics*, 8, 1996. <http://mizar.org/JFM/Vol8/jordan3.html>.
- [23] Yatsuka Nakamura and Andrzej Trybulec. Decomposing a Go-Board into cells. *Journal of Formalized Mathematics*, 7, 1995. <http://mizar.org/JFM/Vol7/goboard5.html>.
- [24] Takaya Nishiyama and Yasuho Mizuhara. Binary arithmetics. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/binarith.html>.
- [25] Beata Padlewska. Connected spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/connsp_1.html.
- [26] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/pre_topc.html.
- [27] Andrzej Trybulec. Enumerated sets. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Voll/enumset1.html>.
- [28] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [29] Andrzej Trybulec. Left and right component of the complement of a special closed curve. *Journal of Formalized Mathematics*, 7, 1995. <http://mizar.org/JFM/Vol7/goboard9.html>.

- [30] Andrzej Trybulec and Yatsuka Nakamura. On the order on a special polygon. *Journal of Formalized Mathematics*, 9, 1997. http://mizar.org/JFM/Vol9/sprect_2.html.
- [31] Andrzej Trybulec and Yatsuka Nakamura. On the rectangular finite sequences of the points of the plane. *Journal of Formalized Mathematics*, 9, 1997. http://mizar.org/JFM/Vol9/sprect_1.html.
- [32] Wojciech A. Trybulec. Pigeon hole principle. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_4.html.
- [33] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [34] Mirosław Wysocki and Agata Darmochwał. Subsets of topological spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/tops_1.html.

Received October 22, 1998

Published January 2, 2004
