

Sequences in \mathcal{E}_T^N

Agnieszka Sakowicz
Warsaw University
Białystok

Jarosław Gryko
Warsaw University
Białystok

Adam Grabowski
Warsaw University
Białystok

MML Identifier: TOPRNS_1.

WWW: http://mizar.org/JFM/Vol6/toprns_1.html

The articles [10], [11], [12], [2], [5], [6], [8], [9], [1], [3], [4], and [7] provide the notation and terminology for this paper.

Let N be a natural number. A sequence in \mathcal{E}_T^N is a sequence of \mathcal{E}_T^N .

For simplicity, we follow the rules: N, n, m are natural numbers, q, r, r_1, r_2 are real numbers, x is a set, w, w_1, w_2, g are points of \mathcal{E}_T^N , and s_1, s_2, s_3, s_4, s'_1 are sequences in \mathcal{E}_T^N .

Next we state the proposition

(2)¹ Let f be a function. Then f is a sequence in \mathcal{E}_T^N if and only if $\text{dom } f = \mathbb{N}$ and for every n holds $f(n)$ is a point of \mathcal{E}_T^N .

Let us consider N and let I_1 be a sequence in \mathcal{E}_T^N . We say that I_1 is non-zero if and only if:

(Def. 1) $\text{rng } I_1 \subseteq (\text{the carrier of } \mathcal{E}_T^N) \setminus \{0_{\mathcal{E}_T^N}\}$.

One can prove the following propositions:

(3) s_1 is non-zero iff for every x such that $x \in \mathbb{N}$ holds $s_1(x) \neq 0_{\mathcal{E}_T^N}$.

(4) s_1 is non-zero iff for every n holds $s_1(n) \neq 0_{\mathcal{E}_T^N}$.

(5) For all N, s_1, s_2 such that for every x such that $x \in \mathbb{N}$ holds $s_1(x) = s_2(x)$ holds $s_1 = s_2$.

(6) For all N, s_1, s_2 such that for every n holds $s_1(n) = s_2(n)$ holds $s_1 = s_2$.

The scheme *ExTopRealNSeq* deals with a natural number \mathcal{A} and a unary functor \mathcal{F} yielding a point of $\mathcal{E}_T^{\mathcal{A}}$, and states that:

There exists a sequence s_1 in $\mathcal{E}_T^{\mathcal{A}}$ such that for every n holds $s_1(n) = \mathcal{F}(n)$ for all values of the parameters.

Let us consider N, s_2, s_3 . The functor $s_2 + s_3$ yielding a sequence in \mathcal{E}_T^N is defined as follows:

(Def. 2) For every n holds $(s_2 + s_3)(n) = s_2(n) + s_3(n)$.

Let us consider r, N, s_1 . The functor $r \cdot s_1$ yields a sequence in \mathcal{E}_T^N and is defined as follows:

(Def. 3) For every n holds $(r \cdot s_1)(n) = r \cdot s_1(n)$.

Let us consider N, s_1 . The functor $-s_1$ yielding a sequence in \mathcal{E}_T^N is defined as follows:

¹ The proposition (1) has been removed.

(Def. 4) For every n holds $(-s_1)(n) = -s_1(n)$.

Let us consider N, s_2, s_3 . The functor $s_2 - s_3$ yields a sequence in \mathcal{E}_T^N and is defined by:

(Def. 5) $s_2 - s_3 = s_2 + -s_3$.

Let us consider N and let x be a point of \mathcal{E}_T^N . The functor $|x|$ yielding a real number is defined by:

(Def. 6) There exists a finite sequence y of elements of \mathbb{R} such that $x = y$ and $|x| = |y|$.

Let us consider N, s_1 . The functor $|s_1|$ yields a sequence of real numbers and is defined as follows:

(Def. 7) For every n holds $|s_1|(n) = |s_1(n)|$.

One can prove the following propositions:

$$(8)^2 \quad |r| \cdot |w| = |r \cdot w|.$$

$$(9) \quad |r \cdot s_1| = |r| |s_1|.$$

$$(10) \quad s_2 + s_3 = s_3 + s_2.$$

$$(11) \quad (s_2 + s_3) + s_4 = s_2 + (s_3 + s_4).$$

$$(12) \quad -s_1 = (-1) \cdot s_1.$$

$$(13) \quad r \cdot (s_2 + s_3) = r \cdot s_2 + r \cdot s_3.$$

$$(14) \quad (r \cdot q) \cdot s_1 = r \cdot (q \cdot s_1).$$

$$(15) \quad r \cdot (s_2 - s_3) = r \cdot s_2 - r \cdot s_3.$$

$$(16) \quad s_2 - (s_3 + s_4) = s_2 - s_3 - s_4.$$

$$(17) \quad 1 \cdot s_1 = s_1.$$

$$(18) \quad --s_1 = s_1.$$

$$(19) \quad s_2 - -s_3 = s_2 + s_3.$$

$$(20) \quad s_2 - (s_3 - s_4) = (s_2 - s_3) + s_4.$$

$$(21) \quad s_2 + (s_3 - s_4) = (s_2 + s_3) - s_4.$$

(22) If $r \neq 0$ and s_1 is non-zero, then $r \cdot s_1$ is non-zero.

(23) If s_1 is non-zero, then $-s_1$ is non-zero.

$$(24) \quad |0_{\mathcal{E}_T^N}| = 0.$$

(25) If $|w| = 0$, then $w = 0_{\mathcal{E}_T^N}$.

$$(26) \quad |w| \geq 0.$$

$$(27) \quad |-w| = |w|.$$

$$(28) \quad |w_1 - w_2| = |w_2 - w_1|.$$

$$(29) \quad |w_1 - w_2| = 0 \text{ iff } w_1 = w_2.$$

$$(30) \quad |w_1 + w_2| \leq |w_1| + |w_2|.$$

$$(31) \quad |w_1 - w_2| \leq |w_1| + |w_2|.$$

² The proposition (7) has been removed.

$$(32) \quad |w_1| - |w_2| \leq |w_1 + w_2|.$$

$$(33) \quad |w_1| - |w_2| \leq |w_1 - w_2|.$$

$$(34) \quad \text{If } w_1 \neq w_2, \text{ then } |w_1 - w_2| > 0.$$

$$(35) \quad |w_1 - w_2| \leq |w_1 - w| + |w - w_2|.$$

$$(36) \quad \text{If } 0 \leq r_1 \text{ and } |w_1| < |w_2| \text{ and } r_1 < r_2, \text{ then } |w_1| \cdot r_1 < |w_2| \cdot r_2.$$

$$(38)^3 \quad -|w| < r \text{ and } r < |w| \text{ iff } |r| < |w|.$$

Let us consider N and let I_1 be a sequence in \mathcal{E}_T^N . We say that I_1 is bounded if and only if:

(Def. 8) There exists r such that for every n holds $|I_1(n)| < r$.

The following proposition is true

$$(39) \quad \text{For every } n \text{ there exists } r \text{ such that } 0 < r \text{ and for every } m \text{ such that } m \leq n \text{ holds } |s_1(m)| < r.$$

Let us consider N and let I_1 be a sequence in \mathcal{E}_T^N . We say that I_1 is convergent if and only if:

(Def. 9) There exists g such that for every r such that $0 < r$ there exists n such that for every m such that $n \leq m$ holds $|I_1(m) - g| < r$.

Let us consider N, s_1 . Let us assume that s_1 is convergent. The functor $\lim s_1$ yielding a point of \mathcal{E}_T^N is defined by:

(Def. 10) For every r such that $0 < r$ there exists n such that for every m such that $n \leq m$ holds $|s_1(m) - \lim s_1| < r$.

One can prove the following propositions:

$$(41)^4 \quad \text{If } s_1 \text{ is convergent and } s'_1 \text{ is convergent, then } s_1 + s'_1 \text{ is convergent.}$$

$$(42) \quad \text{If } s_1 \text{ is convergent and } s'_1 \text{ is convergent, then } \lim(s_1 + s'_1) = \lim s_1 + \lim s'_1.$$

$$(43) \quad \text{If } s_1 \text{ is convergent, then } r \cdot s_1 \text{ is convergent.}$$

$$(44) \quad \text{If } s_1 \text{ is convergent, then } \lim(r \cdot s_1) = r \cdot \lim s_1.$$

$$(45) \quad \text{If } s_1 \text{ is convergent, then } -s_1 \text{ is convergent.}$$

$$(46) \quad \text{If } s_1 \text{ is convergent, then } \lim(-s_1) = -\lim s_1.$$

$$(47) \quad \text{If } s_1 \text{ is convergent and } s'_1 \text{ is convergent, then } s_1 - s'_1 \text{ is convergent.}$$

$$(48) \quad \text{If } s_1 \text{ is convergent and } s'_1 \text{ is convergent, then } \lim(s_1 - s'_1) = \lim s_1 - \lim s'_1.$$

$$(50)^5 \quad \text{If } s_1 \text{ is convergent, then } s_1 \text{ is bounded.}$$

$$(51) \quad \text{If } s_1 \text{ is convergent, then if } \lim s_1 \neq 0_{\mathcal{E}_T^N}, \text{ then there exists } n \text{ such that for every } m \text{ such that } n \leq m \text{ holds } \frac{|\lim s_1|}{2} < |s_1(m)|.$$

³ The proposition (37) has been removed.

⁴ The proposition (40) has been removed.

⁵ The proposition (49) has been removed.

REFERENCES

- [1] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.
- [2] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [3] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_2.html.
- [4] Agata Darmochwał. The Euclidean space. *Journal of Formalized Mathematics*, 3, 1991. <http://mizar.org/JFM/Vol3/euclid.html>.
- [5] Krzysztof Hryniewiecki. Basic properties of real numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/real_1.html.
- [6] Jarosław Kotowicz. Real sequences and basic operations on them. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/seq_1.html.
- [7] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/pre_topc.html.
- [8] Jan Popiołek. Some properties of functions modul and signum. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/absvalue.html>.
- [9] Jan Popiołek. Real normed space. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/normsp_1.html.
- [10] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [11] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [12] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received May 10, 1994

Published January 2, 2004
