## The Scott Topology. Part II<sup>1</sup>

Czesław Byliński Warsaw University Białystok

Piotr Rudnicki University of Alberta Edmonton

**Summary.** Mizar formalization of pp. 105–108 of [10] which continues [27]. We found a simplification for the proof of Corollary 1.15, in the last case, see the proof in the Mizar article for details.

MML Identifier: WAYBEL14.

WWW: http://mizar.org/JFM/Vol9/waybel14.html

The articles [23], [7], [29], [18], [9], [22], [6], [28], [2], [1], [20], [19], [24], [21], [8], [3], [11], [12], [13], [25], [26], [4], [14], [5], [30], [16], [17], [15], and [27] provide the notation and terminology for this paper.

## 1. PRELIMINARIES

One can prove the following propositions:

- (1) Let X be a set and F be a finite family of subsets of X. Then there exists a finite family G of subsets of X such that  $G \subseteq F$  and  $\bigcup G = \bigcup F$  and for every subset g of X such that  $g \in G$  holds  $g \not\subseteq \bigcup (G \setminus \{g\})$ .
- (2) For every 1-sorted structure S and for every subset X of S holds  $X^c$  = the carrier of S iff X is empty.
- (3) Let *R* be an antisymmetric transitive non empty relational structure with g.l.b.'s and *x*, *y* be elements of *R*. Then  $\downarrow (x \sqcap y) = \downarrow x \cap \downarrow y$ .
- (4) Let *R* be an antisymmetric transitive non empty relational structure with l.u.b.'s and *x*, *y* be elements of *R*. Then  $\uparrow(x \sqcup y) = \uparrow x \cap \uparrow y$ .
- (5) Let *L* be a complete antisymmetric non empty relational structure and *X* be a lower subset of *L*. If  $\sup X \in X$ , then  $X = \bigcup \sup X$ .
- (6) Let *L* be a complete antisymmetric non empty relational structure and *X* be an upper subset of *L*. If  $\inf X \in X$ , then  $X = \uparrow \inf X$ .
- (7) Let *R* be a non empty reflexive transitive relational structure and *x*, *y* be elements of *R*. Then  $x \ll y$  if and only if  $\uparrow y \subseteq \uparrow x$ .
- (8) Let R be a non empty reflexive transitive relational structure and x, y be elements of R. Then  $x \ll y$  if and only if  $\downarrow x \subseteq \mbox{$\downarrow$} y$ .

<sup>&</sup>lt;sup>1</sup>This work was partially supported by NSERC Grant OGP9207 and NATO CRG 951368.

- (9) Let R be a complete reflexive antisymmetric non empty relational structure and x be an element of R. Then  $\sup x \le x$  and  $x \le \inf x$ .
- (10) For every lower-bounded antisymmetric non empty relational structure L holds  $\uparrow(\bot_L)$  = the carrier of L.
- (11) For every upper-bounded antisymmetric non empty relational structure L holds  $\downarrow(\top_L)$  = the carrier of L.
- (12) For every poset *P* with l.u.b.'s and for all elements x, y of *P* holds  $\uparrow x \sqcup \uparrow y \subseteq \uparrow (x \sqcup y)$ .
- (13) For every poset *P* with g.l.b.'s and for all elements x, y of *P* holds  $\downarrow x \sqcap \downarrow y \subseteq \downarrow (x \sqcap y)$ .
- (14) Let *R* be a non empty poset with l.u.b.'s and *l* be an element of *R*. Then *l* is co-prime if and only if for all elements x, y of R such that  $l \le x \sqcup y$  holds  $l \le x$  or  $l \le y$ .
- (15) For every complete non empty poset P and for every non empty subset V of P holds  $\downarrow \inf V = \bigcap \{ \downarrow u; u \text{ ranges over elements of } P: u \in V \}.$
- (16) For every complete non empty poset P and for every non empty subset V of P holds  $\uparrow \sup V = \bigcap \{ \uparrow u; u \text{ ranges over elements of } P: u \in V \}.$

Let L be a sup-semilattice and let x be an element of L. Observe that compactbelow(x) is directed.

We now state four propositions:

- (17) Let T be a non empty topological space, S be an irreducible subset of T, and V be an element of  $\langle$  the topology of T,  $\subseteq \rangle$ . If  $V = S^c$ , then V is prime.
- (18) Let T be a non empty topological space and x, y be elements of  $\langle$  the topology of T,  $\subseteq \rangle$ . Then  $x \sqcup y = x \cup y$  and  $x \sqcap y = x \cap y$ .
- (19) Let T be a non empty topological space and V be an element of  $\langle$  the topology of  $T, \subseteq \rangle$ . Then V is prime if and only if for all elements X, Y of  $\langle$  the topology of T,  $\subseteq \rangle$  such that  $X \cap Y \subseteq V$  holds  $X \subseteq V$  or  $Y \subseteq V$ .
- (20) Let T be a non empty topological space and V be an element of  $\langle$  the topology of  $T, \subseteq \rangle$ . Then V is co-prime if and only if for all elements X, Y of  $\langle$  the topology of T,  $\subseteq \rangle$  such that  $V \subseteq X \cup Y$  holds  $V \subseteq X$  or  $V \subseteq Y$ .

Let T be a non empty topological space. Observe that  $\langle$  the topology of  $T, \subseteq \rangle$  is distributive. Next we state two propositions:

- (21) Let T be a non empty topological space, L be a top-lattice, t be a point of T, t be a point of L, and X be a family of subsets of L. Suppose the topological structure of T = the topological structure of L and t = t and t is a basis of t.
- (22) Let *L* be a top-lattice and *x* be an element of *L*. Suppose that for every subset *X* of *L* such that *X* is open holds *X* is upper. Then  $\uparrow x$  is compact.

## 2. The Scott topology

For simplicity, we use the following convention: L denotes a complete Scott top-lattice, x denotes an element of L, X, Y denote subsets of L, V, W denote elements of  $\langle \sigma(L), \subseteq \rangle$ , and  $V_1$  denotes a subset of  $\langle \sigma(L), \subseteq \rangle$ .

Let *L* be a complete lattice. Observe that  $\sigma(L)$  is non empty.

Next we state four propositions:

1

 $<sup>^{1}</sup>$   $\sigma(L) = \text{sigma } L$ , as defined in [27, p. 316, Def. 12] and  $\sqcup_{L} = \text{sup\_op}(L)$ , as defined in [14, p. 163, Def. 51]

- (23)  $\sigma(L)$  = the topology of L.
- (24)  $X \in \sigma(L)$  iff X is open.
- (25) For every filtered subset *X* of *L* such that  $V_1 = \{(\downarrow x)^c : x \in X\}$  holds  $V_1$  is directed.
- (26) If *X* is open and  $x \in X$ , then  $\inf X \ll x$ .

Let R be a non empty reflexive relational structure and let f be a map from [:R,R:] into R. We say that f is jointly Scott-continuous if and only if the condition (Def. 1) is satisfied.

(Def. 1) Let T be a non empty topological space. Suppose the topological structure of T = ConvergenceSpace(the Scott convergence of R). Then there exists a map  $f_1$  from [:T,T:] into T such that  $f_1 = f$  and  $f_1$  is continuous.

We now state a number of propositions:

- (27) If V = X, then V is co-prime iff X is filtered and upper.
- (28) If V = X and there exists x such that  $X = (\downarrow x)^c$ , then V is prime and  $V \neq$  the carrier of L.
- (29) If V = X and  $\sqcup_L$  is jointly Scott-continuous and V is prime and  $V \neq$  the carrier of L, then there exists x such that  $X = (\downarrow x)^c$ .
- (30) If *L* is continuous, then  $\sqcup_L$  is jointly Scott-continuous.
- (31) If  $\sqcup_L$  is jointly Scott-continuous, then *L* is sober.
- (32) If L is continuous, then L is compact, locally-compact, sober, and Baire.
- (33) If *L* is continuous and  $X \in \sigma(L)$ , then  $X = \bigcup \{ \uparrow x : x \in X \}$ .
- (34) If for every *X* such that  $X \in \sigma(L)$  holds  $X = \bigcup \{\uparrow x : x \in X\}$ , then *L* is continuous.
- (35) If L is continuous, then there exists a basis B of x such that for every X such that  $X \in B$  holds X is open and filtered.
- (36) If *L* is continuous, then  $\langle \sigma(L), \subseteq \rangle$  is continuous.
- (37) Suppose for every x there exists a basis B of x such that for every Y such that  $Y \in B$  holds Y is open and filtered and  $\langle \sigma(L), \subseteq \rangle$  is continuous. Then  $x = \bigsqcup_{L} \{\inf X : x \in X \land X \in \sigma(L)\}$ .
- (38) If for every *x* holds  $x = \bigsqcup_{L} \{\inf X : x \in X \land X \in \sigma(L)\}$ , then *L* is continuous.
- (39) The following statements are equivalent
  - (i) for every x there exists a basis B of x such that for every Y such that  $Y \in B$  holds Y is open and filtered,
- (ii) for every V there exists  $V_1$  such that  $V = \sup V_1$  and for every W such that  $W \in V_1$  holds W is co-prime.
- (40) For every V there exists  $V_1$  such that  $V = \sup V_1$  and for every W such that  $W \in V_1$  holds W is co-prime and  $\langle \sigma(L), \subseteq \rangle$  is continuous if and only if  $\langle \sigma(L), \subseteq \rangle$  is completely-distributive.
- (41)  $\langle \sigma(L), \subseteq \rangle$  is completely-distributive iff  $\langle \sigma(L), \subseteq \rangle$  is continuous and  $(\langle \sigma(L), \subseteq \rangle)^{op}$  is continuous.
- (42) If L is algebraic, then there exists a basis B of L such that  $B = \{\uparrow x : x \in \text{the carrier of } CompactSublatt(L)\}$ .
- (43) Given a basis B of L such that  $B = \{\uparrow x : x \in \text{the carrier of CompactSublatt}(L)\}$ . Then  $\langle \sigma(L), \subseteq \rangle$  is algebraic and for every V there exists  $V_1$  such that  $V = \sup V_1$  and for every W such that  $W \in V_1$  holds W is co-prime.

- (44) Suppose  $\langle \sigma(L), \subseteq \rangle$  is algebraic and for every V there exists  $V_1$  such that  $V = \sup V_1$  and for every W such that  $W \in V_1$  holds W is co-prime. Then there exists a basis B of L such that  $B = \{ \uparrow x : x \in \text{the carrier of CompactSublatt}(L) \}$ .
- (45) If there exists a basis B of L such that  $B = \{ \uparrow x : x \in \text{the carrier of CompactSublatt}(L) \}$ , then L is algebraic.

## REFERENCES

- [1] Grzegorz Bancerek. Cardinal numbers. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/card\_1.html.
- [2] Grzegorz Bancerek. Complete lattices. Journal of Formalized Mathematics, 4, 1992. http://mizar.org/JFM/Vol4/lattice3.html.
- [3] Grzegorz Bancerek. Bounds in posets and relational substructures. Journal of Formalized Mathematics, 8, 1996. http://mizar.org/ JFM/Vol8/yellow\_0.html.
- [4] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. Journal of Formalized Mathematics, 8, 1996. http://mizar.org/ JFM/Vol8/waybel\_0.html.
- [5] Grzegorz Bancerek. The "way-below" relation. Journal of Formalized Mathematics, 8, 1996. http://mizar.org/JFM/Vol8/waybel\_ 3.html.
- [6] Czesław Byliński. Functions and their basic properties. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/funct\_1.html.
- [7] Czesław Byliński. Some basic properties of sets. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/zfmisc\_1.html.
- [8] Agata Darmochwał. Compact spaces. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/compts\_1.html.
- [9] Agata Darmochwał. Finite sets. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/finset\_1.html.
- [10] G. Gierz, K.H. Hofmann, K. Keimel, J.D. Lawson, M. Mislove, and D.S. Scott. A Compendium of Continuous Lattices. Springer-Verlag, Berlin, Heidelberg, New York, 1980.
- [11] Adam Grabowski and Robert Milewski. Boolean posets, posets under inclusion and products of relational structures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow\_1.html.
- [12] Artur Korniłowicz. Cartesian products of relations and relational structures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow\_3.html.
- [13] Artur Korniłowicz. Definitions and properties of the join and meet of subsets. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow\_4.html.
- [14] Artur Korniłowicz. Meet continuous lattices. Journal of Formalized Mathematics, 8, 1996. http://mizar.org/JFM/Vol8/waybel\_ 2.html.
- [15] Artur Komiłowicz. On the topological properties of meet-continuous lattices. Journal of Formalized Mathematics, 8, 1996. http://mizar.org/JFM/Vol8/waybel\_9.html.
- [16] Beata Madras. Irreducible and prime elements. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel\_6.html.
- $[17] \begin{tabular}{l} Robert Milewski. Algebraic lattices. {\it Journal of Formalized Mathematics}, 8, 1996. http://mizar.org/JFM/Vol8/waybel_8.html. \\ \end{tabular}$
- [18] Beata Padlewska. Families of sets. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/setfam\_1.html.
- [19] Beata Padlewska. Locally connected spaces. Journal of Formalized Mathematics, 2, 1990. http://mizar.org/JFM/Vol2/connsp\_2.html.
- [20] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/pre\_topc.html.
- [21] Alexander Yu. Shibakov and Andrzej Trybulec. The Cantor set. Journal of Formalized Mathematics, 7, 1995. http://mizar.org/ JFM/Vol7/cantor\_1.html.
- [22] Andrzej Trybulec. Domains and their Cartesian products. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/domain\_1.html.
- [23] Andrzej Trybulec. Tarski Grothendieck set theory. Journal of Formalized Mathematics, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.
- [24] Andrzej Trybulec. A Borsuk theorem on homotopy types. Journal of Formalized Mathematics, 3, 1991. http://mizar.org/JFM/Vol3/borsuk\_1.html.
- [25] Andrzej Trybulec. Moore-Smith convergence. Journal of Formalized Mathematics, 8, 1996. http://mizar.org/JFM/Vol8/yellow\_6.html.
- [26] Andrzej Trybulec. Baire spaces, Sober spaces. Journal of Formalized Mathematics, 9, 1997. http://mizar.org/JFM/Vol9/yellow\_8.html.

- [27] Andrzej Trybulec. Scott topology. Journal of Formalized Mathematics, 9, 1997. http://mizar.org/JFM/Vol9/waybelll.html.
- [28] Wojciech A. Trybulec. Partially ordered sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/orders\_1.html.
- $[29] \ \ \textbf{Zinaida Trybulec. Properties of subsets.} \ \textbf{\textit{Journal of Formalized Mathematics}}, 1, 1989. \ \texttt{http://mizar.org/JFM/Vol1/subset\_1.html}.$
- [30] Mariusz Żynel. The equational characterization of continuous lattices. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel\_5.html.

Received August 27, 1997

Published January 2, 2004