

Scott-Continuous Functions¹

Adam Grabowski
University of Białystok

Summary. The article is a translation of [10, pp. 112–113].

MML Identifier: WAYBEL17.

WWW: <http://mizar.org/JFM/Vol10/waybel17.html>

The articles [18], [9], [23], [21], [1], [25], [24], [6], [8], [7], [17], [13], [22], [2], [3], [11], [4], [12], [26], [19], [5], [16], [15], [20], and [14] provide the notation and terminology for this paper.

1. PRELIMINARIES

Let S be a non empty set and let a, b be elements of S . The functor a, b, \dots yielding a function from \mathbb{N} into S is defined by the condition (Def. 1).

(Def. 1) Let i be a natural number. Then

- (i) if there exists a natural number k such that $i = 2 \cdot k$, then $(a, b, \dots)(i) = a$, and
- (ii) if it is not true that there exists a natural number k such that $i = 2 \cdot k$, then $(a, b, \dots)(i) = b$.

One can prove the following three propositions:

- (1) Let S, T be non empty reflexive relational structures, f be a map from S into T , and P be a lower subset of T . If f is monotone, then $f^{-1}(P)$ is lower.
- (2) Let S, T be non empty reflexive relational structures, f be a map from S into T , and P be an upper subset of T . If f is monotone, then $f^{-1}(P)$ is upper.
- (3) Let S, T be reflexive antisymmetric non empty relational structures and f be a map from S into T . If f is directed-sups-preserving, then f is monotone.

Let S, T be reflexive antisymmetric non empty relational structures. Note that every map from S into T which is directed-sups-preserving is also monotone.

Next we state the proposition

- (4) Let S, T be up-complete Scott top-lattices and f be a map from S into T . If f is continuous, then f is monotone.

Let S, T be up-complete Scott top-lattices. One can verify that every map from S into T which is continuous is also monotone.

¹This work has been supported by KBN Grant 8 T11C 018 12.

2. POSET OF CONTINUOUS MAPS

Let S be a set and let T be a reflexive relational structure. One can check that $S \mapsto T$ is reflexive-yielding.

Let S be a non empty set and let T be a complete lattice. Observe that T^S is complete.

Let S, T be up-complete Scott top-lattices. The functor $\text{SCMaps}(S, T)$ yielding a strict full relational substructure of $\text{MonMaps}(S, T)$ is defined as follows:

(Def. 2) For every map f from S into T holds $f \in \text{the carrier of } \text{SCMaps}(S, T)$ iff f is continuous.

Let S, T be up-complete Scott top-lattices. Observe that $\text{SCMaps}(S, T)$ is non empty.

3. SOME SPECIAL NETS

Let S be a non empty relational structure and let a, b be elements of S . The functor $\text{NetStr}(a, b)$ yields a strict non empty net structure over S and is defined by the conditions (Def. 3).

(Def. 3)(i) The carrier of $\text{NetStr}(a, b) = \mathbb{N}$,

(ii) the mapping of $\text{NetStr}(a, b) = a, b, \dots$, and

(iii) for all elements i, j of $\text{NetStr}(a, b)$ and for all natural numbers i', j' such that $i = i'$ and $j = j'$ holds $i \leq j$ iff $i' \leq j'$.

Let S be a non empty relational structure and let a, b be elements of S . Observe that $\text{NetStr}(a, b)$ is reflexive, transitive, directed, and antisymmetric.

Next we state four propositions:

(5) Let S be a non empty relational structure, a, b be elements of S , and i be an element of $\text{NetStr}(a, b)$. Then $(\text{NetStr}(a, b))(i) = a$ or $(\text{NetStr}(a, b))(i) = b$.

(6) Let S be a non empty relational structure, a, b be elements of S , i, j be elements of $\text{NetStr}(a, b)$, and i', j' be natural numbers such that $i' = i$ and $j' = i' + 1$ and $j' = j$. Then

(i) if $(\text{NetStr}(a, b))(i) = a$, then $(\text{NetStr}(a, b))(j) = b$, and

(ii) if $(\text{NetStr}(a, b))(i) = b$, then $(\text{NetStr}(a, b))(j) = a$.

(7) For every poset S with g.l.b.'s and for all elements a, b of S holds $\liminf \text{NetStr}(a, b) = a \sqcap b$.

(8) Let S, T be posets with g.l.b.'s, a, b be elements of S , and f be a map from S into T . Then $\liminf(f \cdot \text{NetStr}(a, b)) = f(a) \sqcap f(b)$.

Let S be a non empty relational structure and let D be a non empty subset of S . The functor $\text{NetStr}(D)$ yielding a strict net structure over S is defined by:

(Def. 4) $\text{NetStr}(D) = \langle D, (\text{the internal relation of } S)^2 \upharpoonright D, \text{id}_{\text{the carrier of } S \upharpoonright D} \rangle$.

One can prove the following proposition

(9) Let S be a non empty reflexive relational structure and D be a non empty subset of S . Then $\text{NetStr}(D) = \text{NetStr}(D, \text{id}_{\text{the carrier of } S \upharpoonright D})$.

Let S be a non empty reflexive relational structure and let D be a directed non empty subset of S . Observe that $\text{NetStr}(D)$ is non empty, directed, and reflexive.

Let S be a non empty reflexive transitive relational structure and let D be a directed non empty subset of S . One can verify that $\text{NetStr}(D)$ is transitive.

Let S be a non empty reflexive relational structure and let D be a directed non empty subset of S . Observe that $\text{NetStr}(D)$ is monotone.

The following proposition is true

(10) For every up-complete lattice S and for every directed non empty subset D of S holds $\liminf \text{NetStr}(D) = \sup D$.

4. MONOTONE MAPS

The following propositions are true:

- (11) Let S, T be lattices and f be a map from S into T . If for every net N in S holds $f(\liminf N) \leq \liminf(f \cdot N)$, then f is monotone.
- (12) Let S, T be continuous lower-bounded lattices and f be a map from S into T . Suppose f is directed-sups-preserving. Let x be an element of S . Then $f(x) = \bigsqcup_T \{f(w); w \text{ ranges over elements of } S: w \ll x\}$.
- (13) Let S be a lattice, T be an up-complete lower-bounded lattice, and f be a map from S into T . Suppose that for every element x of S holds $f(x) = \bigsqcup_T \{f(w); w \text{ ranges over elements of } S: w \ll x\}$. Then f is monotone.
- (14) Let S be an up-complete lower-bounded lattice, T be a continuous lower-bounded lattice, and f be a map from S into T . Suppose that for every element x of S holds $f(x) = \bigsqcup_T \{f(w); w \text{ ranges over elements of } S: w \ll x\}$. Let x be an element of S and y be an element of T . Then $y \ll f(x)$ if and only if there exists an element w of S such that $w \ll x$ and $y \ll f(w)$.
- (15) Let S, T be non empty relational structures, D be a subset of S , and f be a map from S into T . Suppose that
- (i) $\sup D$ exists in S and $\sup f^\circ D$ exists in T , or
 - (ii) S is complete and antisymmetric and T is complete and antisymmetric.
- If f is monotone, then $\sup(f^\circ D) \leq f(\sup D)$.
- (16) Let S, T be non empty reflexive antisymmetric relational structures, D be a directed non empty subset of S , and f be a map from S into T . Suppose $\sup D$ exists in S and $\sup f^\circ D$ exists in T or S is up-complete and T is up-complete. If f is monotone, then $\sup(f^\circ D) \leq f(\sup D)$.
- (17) Let S, T be non empty relational structures, D be a subset of S , and f be a map from S into T . Suppose that
- (i) $\inf D$ exists in S and $\inf f^\circ D$ exists in T , or
 - (ii) S is complete and antisymmetric and T is complete and antisymmetric.
- If f is monotone, then $f(\inf D) \leq \inf(f^\circ D)$.
- (18) Let S, T be up-complete lattices, f be a map from S into T , and N be a monotone non empty net structure over S . If f is monotone, then $f \cdot N$ is monotone.

Let S, T be up-complete lattices, let f be a monotone map from S into T , and let N be a monotone non empty net structure over S . Note that $f \cdot N$ is monotone.

One can prove the following two propositions:

- (19) Let S, T be up-complete lattices and f be a map from S into T . Suppose that for every net N in S holds $f(\liminf N) \leq \liminf(f \cdot N)$. Let D be a directed non empty subset of S . Then $\sup(f^\circ D) = f(\sup D)$.
- (20) Let S, T be complete lattices, f be a map from S into T , N be a net in S , j be an element of N , and j' be an element of $f \cdot N$. Suppose $j' = j$. Suppose f is monotone. Then $f(\bigsqcap_S \{N(k); k \text{ ranges over elements of } N: k \geq j\}) \leq \bigsqcap_T \{(f \cdot N)(l); l \text{ ranges over elements of } f \cdot N: l \geq j'\}$.

5. NECESSARY AND SUFFICIENT CONDITIONS OF SCOTT-CONTINUITY

One can prove the following propositions:

- (21) Let S, T be complete Scott top-lattices and f be a map from S into T . Then f is continuous if and only if for every net N in S holds $f(\liminf N) \leq \liminf(f \cdot N)$.

- (22) Let S, T be complete Scott top-lattices and f be a map from S into T . Then f is continuous if and only if f is directed-sups-preserving.

Let S, T be complete Scott top-lattices. Observe that every map from S into T which is continuous is also directed-sups-preserving and every map from S into T which is directed-sups-preserving is also continuous.

The following propositions are true:

- (23) Let S, T be continuous complete Scott top-lattices and f be a map from S into T . Then f is continuous if and only if for every element x of S and for every element y of T holds $y \ll f(x)$ iff there exists an element w of S such that $w \ll x$ and $y \ll f(w)$.
- (24) Let S, T be continuous complete Scott top-lattices and f be a map from S into T . Then f is continuous if and only if for every element x of S holds $f(x) = \bigsqcup_T \{f(w); w \text{ ranges over elements of } S: w \ll x\}$.
- (25) Let S be a lattice, T be a complete lattice, and f be a map from S into T . Suppose that for every element x of S holds $f(x) = \bigsqcup_T \{f(w); w \text{ ranges over elements of } S: w \leq x \wedge w \text{ is compact}\}$. Then f is monotone.
- (26) Let S, T be complete Scott top-lattices and f be a map from S into T . Suppose that for every element x of S holds $f(x) = \bigsqcup_T \{f(w); w \text{ ranges over elements of } S: w \leq x \wedge w \text{ is compact}\}$. Let x be an element of S . Then $f(x) = \bigsqcup_T \{f(w); w \text{ ranges over elements of } S: w \ll x\}$.
- (27) Let S, T be complete Scott top-lattices and f be a map from S into T . Suppose S is algebraic and T is algebraic. Then f is continuous if and only if for every element x of S and for every element k of T such that $k \in \text{carrier of CompactSublatt}(T)$ holds $k \leq f(x)$ iff there exists an element j of S such that $j \in \text{carrier of CompactSublatt}(S)$ and $j \leq x$ and $k \leq f(j)$.
- (28) Let S, T be complete Scott top-lattices and f be a map from S into T . Suppose S is algebraic and T is algebraic. Then f is continuous if and only if for every element x of S holds $f(x) = \bigsqcup_T \{f(w); w \text{ ranges over elements of } S: w \leq x \wedge w \text{ is compact}\}$.
- (29) Let S, T be up-complete Scott non empty reflexive transitive antisymmetric topological space-like FR-structures and f be a map from S into T . If f is directed-sups-preserving, then f is continuous.

REFERENCES

- [1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/nat_1.html.
- [2] Grzegorz Bancerek. Complete lattices. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/lattice3.html>.
- [3] Grzegorz Bancerek. Bounds in posets and relational substructures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_0.html.
- [4] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_0.html.
- [5] Grzegorz Bancerek. The “way-below” relation. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_3.html.
- [6] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [7] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [8] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/partfun1.html>.
- [9] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [10] G. Gierz, K.H. Hofmann, K. Keimel, J.D. Lawson, M. Mislove, and D.S. Scott. *A Compendium of Continuous Lattices*. Springer-Verlag, Berlin, Heidelberg, New York, 1980.
- [11] Adam Grabowski. On the category of posets. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/orders_3.html.

- [12] Adam Grabowski and Robert Milewski. Boolean posets, posets under inclusion and products of relational structures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_1.html.
- [13] Krzysztof Hryniewiecki. Relations of tolerance. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/toler_1.html.
- [14] Artur Korniłowicz. Meet – continuous lattices. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_2.html.
- [15] Artur Korniłowicz. On the topological properties of meet-continuous lattices. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_9.html.
- [16] Robert Milewski. Algebraic lattices. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_8.html.
- [17] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/pre_topc.html.
- [18] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [19] Andrzej Trybulec. Natural transformations. Discrete categories. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/nattra_1.html.
- [20] Andrzej Trybulec. Scott topology. *Journal of Formalized Mathematics*, 9, 1997. <http://mizar.org/JFM/Vol9/waybel11.html>.
- [21] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [22] Wojciech A. Trybulec. Partially ordered sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/orders_1.html.
- [23] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [24] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.
- [25] Edmund Woronowicz. Relations defined on sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relset_1.html.
- [26] Mariusz Żynel and Czesław Byliński. Properties of relational structures, posets, lattices and maps. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_2.html.

Received February 13, 1998

Published January 2, 2004
