On the Order-consistent Topology of Complete and Uncomplete Lattices

Ewa Grądzka University of Białystok

Summary. This paper is a continuation of the formalisation of [9] pp. 108–109. Order-consistent and upper topologies are defined. The theorem that the Scott and the upper topologies are order-consistent is proved. Remark 1.4 and example 1.5(2) are generalized for proving this theorem.

MML Identifier: WAYBEL32.

WWW: http://mizar.org/JFM/Vol12/waybel32.html

The articles [15], [7], [20], [21], [5], [8], [6], [18], [1], [13], [12], [23], [19], [2], [3], [14], [10], [16], [11], [22], [17], and [4] provide the notation and terminology for this paper.

Let *T* be a non empty FR-structure. We say that *T* is upper if and only if:

(Def. 1) $\{(\downarrow x)^c : x \text{ ranges over elements of } T\}$ is a prebasis of T.

One can check that there exists a top-lattice which is Scott, up-complete, and strict. Let *T* be a topological space-like non empty reflexive FR-structure. We say that *T* is order consistent if and only if the condition (Def. 2) is satisfied.

- (Def. 2) Let x be an element of T. Then
 - (i) $\downarrow x = \{x\}$, and
 - (ii) for every eventually-directed net N in T such that $x = \sup N$ and for every neighbourhood V of x holds N is eventually in V.

Let us note that every non empty reflexive topological space-like FR-structure which is trivial is also upper.

Let us mention that there exists a top-lattice which is upper, trivial, up-complete, and strict. One can prove the following propositions:

- (1) For every upper up-complete non empty top-poset *T* and for every subset *A* of *T* such that *A* is open holds *A* is upper.
- (2) For every up-complete non empty top-poset *T* such that *T* is upper holds *T* is order consistent.
- $(7)^{l}$ For every up-complete non empty reflexive transitive antisymmetric relational structure R holds there exists a topological augmentation of R which is Scott.
- (8) Let *R* be an up-complete non empty poset and *T* be a topological augmentation of *R*. If *T* is Scott, then *T* is correct.

1

¹ The propositions (3)–(6) have been removed.

Let *R* be an up-complete non empty reflexive transitive antisymmetric relational structure. Note that every topological augmentation of *R* which is Scott is also correct.

Let *R* be an up-complete non empty reflexive transitive antisymmetric relational structure. Observe that there exists a topological augmentation of *R* which is Scott and correct.

We now state several propositions:

- (9) Let *T* be a Scott up-complete non empty reflexive transitive antisymmetric FR-structure and *x* be an element of *T*. Then $\overline{\{x\}} = \downarrow x$.
- (10) Every up-complete Scott non empty top-poset is order consistent.
- (11) Let R be an inf-complete semilattice, Z be a net in R, and D be a subset of R. Suppose $D = \{ \bigcap_R \{Z(k); k \text{ ranges over elements of } Z : k \ge j \} : j \text{ ranges over elements of } Z \}$. Then D is non empty and directed.
- (12) Let *R* be an inf-complete semilattice, *S* be a subset of *R*, and *a* be an element of *R*. If $a \in S$, then $\bigcap_R S \leq a$.
- (13) For every inf-complete semilattice R and for every monotone reflexive net N in R holds $\lim \inf N = \sup N$.
- (14) Let R be an inf-complete semilattice and S be a subset of R. Then $S \in$ the topology of ConvergenceSpace(the Scott convergence of R) if and only if S is inaccessible and upper.
- (15) Let R be an inf-complete up-complete semilattice and T be a topological augmentation of R. If the topology of $T = \sigma(R)$, then T is Scott.

Let *R* be an inf-complete up-complete semilattice. Note that there exists a topological augmentation of *R* which is strict, Scott, and correct.

The following two propositions are true:

- (16) Let *S* be an up-complete inf-complete semilattice and *T* be a Scott topological augmentation of *S*. Then $\sigma(S)$ = the topology of *T*.
- (17) Every Scott up-complete non empty reflexive transitive antisymmetric FR-structure is a T_0 -space.

Let *R* be an up-complete non empty reflexive transitive antisymmetric relational structure. Note that every topological augmentation of *R* is up-complete.

One can prove the following propositions:

- (18) Let R be an up-complete non empty reflexive transitive antisymmetric relational structure, T be a Scott topological augmentation of R, x be an element of T, and A be an upper subset of T. If $x \notin A$, then $(\downarrow x)^c$ is a neighbourhood of A.
- (19) Let R be an up-complete non empty reflexive transitive antisymmetric FR-structure, T be a Scott topological augmentation of R, and S be an upper subset of T. Then there exists a family F of subsets of T such that $S = \bigcap F$ and for every subset X of T such that $X \in F$ holds X is a neighbourhood of S.
- (20) Let *T* be a Scott up-complete non empty reflexive transitive antisymmetric FR-structure and *S* be a subset of *T*. Then *S* is open if and only if *S* is upper and property(S).
- (21) Let R be an up-complete non empty reflexive transitive antisymmetric FR-structure, S be a non empty directed subset of R, and a be an element of R. If $a \in S$, then $a \le \bigsqcup_R S$.

Let T be an up-complete non empty reflexive transitive antisymmetric FR-structure. One can check that every subset of T which is lower is also property(S).

Next we state three propositions:

(22) For every finite up-complete non empty poset T holds every subset of T is inaccessible.

- (23) Let R be a complete connected lattice, T be a Scott topological augmentation of R, and x be an element of T. Then $(\downarrow x)^c$ is open.
- (24) Let R be a complete connected lattice, T be a Scott topological augmentation of R, and S be a subset of T. Then S is open if and only if one of the following conditions is satisfied:
 - (i) S =the carrier of T, or
- (ii) $S \in \{(\downarrow x)^c : x \text{ ranges over elements of } T\}.$

Let *R* be an up-complete non empty poset. Note that there exists a correct topological augmentation of *R* which is order consistent.

Let us mention that there exists a top-lattice which is order consistent and complete.

One can prove the following propositions:

- (25) Let *R* be a non empty FR-structure and *A* be a subset of *R*. Suppose that for every element x of *R* holds $\downarrow x = \overline{\{x\}}$. If *A* is open, then *A* is upper.
- (26) Let *R* be a non empty FR-structure and *A* be a subset of *R*. Suppose that for every element x of *R* holds $\downarrow x = \overline{\{x\}}$. Let *A* be a subset of *R*. If *A* is closed, then *A* is lower.
- (27) For every up-complete inf-complete lattice T and for every net N in T and for every element i of N holds $\liminf(N \upharpoonright i) = \liminf N$.

Let S be a non empty 1-sorted structure, let R be a non empty relational structure, and let f be a function from the carrier of R into the carrier of S. The functor R * f yields a strict non empty net structure over S and is defined as follows:

(Def. 3) The relational structure of R * f = the relational structure of R and the mapping of R * f = f.

Let S be a non empty 1-sorted structure, let R be a non empty transitive relational structure, and let f be a function from the carrier of R into the carrier of S. Observe that R * f is transitive.

Let S be a non empty 1-sorted structure, let R be a non empty directed relational structure, and let f be a function from the carrier of R into the carrier of S. Note that R * f is directed.

Let R be a non empty relational structure and let N be a prenet over R. The functor inf_net N yielding a strict prenet over R is defined by:

(Def. 4) There exists a map f from N into R such that $\inf_{n} \text{net } N = N * f$ and for every element i of N holds $f(i) = \bigcap_{R} \{N(k); k \text{ ranges over elements of } N: k \ge i\}$.

Let R be a non empty relational structure and let N be a net in R. One can check that $\inf_{n} N$ is transitive.

Let R be a non empty relational structure and let N be a net in R. Observe that $\inf_{n} \operatorname{net} N$ is directed.

Let R be an inf-complete non empty reflexive antisymmetric relational structure and let N be a net in R. One can check that inf_net N is monotone.

Let R be an inf-complete non empty reflexive antisymmetric relational structure and let N be a net in R. Observe that inf_net N is eventually-directed.

Next we state several propositions:

- (28) Let R be a non empty relational structure and N be a net in R. Then rng (the mapping of inf_net N) = { $\bigcap_R \{N(i); i \text{ ranges over elements of } N: i \ge j\}$: j ranges over elements of N}.
- (29) For every up-complete inf-complete lattice R and for every net N in R holds sup inf_net $N = \liminf N$.
- (30) For every up-complete inf-complete lattice R and for every net N in R and for every element i of N holds sup inf_net $N = \liminf(N \mid i)$.
- (31) Let R be an inf-complete semilattice, N be a net in R, and V be an upper subset of R. If inf_net N is eventually in V, then N is eventually in V.

- (32) Let R be an inf-complete semilattice, N be a net in R, and V be a lower subset of R. If N is eventually in V, then inf_net N is eventually in V.
- (33) Let R be an order consistent up-complete inf-complete non empty top-lattice, N be a net in R, and x be an element of R. If $x \le \liminf N$, then x is a cluster point of N.
- (34) Let R be an order consistent up-complete inf-complete non empty top-lattice, N be an eventually-directed net in R, and x be an element of R. Then $x \le \liminf N$ if and only if x is a cluster point of N.

ACKNOWLEDGMENTS

I would like to thank Dr. Grzegorz Bancerek for his help in the preparation of this article.

REFERENCES

- [1] Grzegorz Bancerek. Complete lattices. Journal of Formalized Mathematics, 4, 1992. http://mizar.org/JFM/Vol4/lattice3.html.
- [2] Grzegorz Bancerek. Bounds in posets and relational substructures. Journal of Formalized Mathematics, 8, 1996. http://mizar.org/ JFM/Vol8/yellow_0.html.
- [3] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. Journal of Formalized Mathematics, 8, 1996. http://mizar.org/ JFM/Vol8/waybel_0.html.
- [4] Grzegorz Bancerek. Bases and refinements of topologies. Journal of Formalized Mathematics, 10, 1998. http://mizar.org/JFM/ Vol10/yellow_9.html.
- [5] Czesław Byliński. Functions and their basic properties. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/funct_1.html.
- [6] Czesław Byliński. Functions from a set to a set. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [7] Czesław Byliński. Some basic properties of sets. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/zfmisc_1.html.
- [8] Agata Darmochwał. Finite sets. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/finset_1.html.
- [9] G. Gierz, K.H. Hofmann, K. Keimel, J.D. Lawson, M. Mislove, and D.S. Scott. A Compendium of Continuous Lattices. Springer-Verlag, Berlin, Heidelberg, New York, 1980.
- [10] Artur Korniłowicz. Meet continuous lattices. Journal of Formalized Mathematics, 8, 1996. http://mizar.org/JFM/Vol8/waybel_ 2.html.
- [11] Artur Korniłowicz. On the topological properties of meet-continuous lattices. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel 9.html.
- [12] Beata Padlewska. Locally connected spaces. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/connsp_2.html.
- [13] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/pre_topo.html.
- [14] Alexander Yu. Shibakov and Andrzej Trybulec. The Cantor set. Journal of Formalized Mathematics, 7, 1995. http://mizar.org/ JFM/Vol7/cantor_1.html.
- [15] Andrzej Trybulec. Tarski Grothendieck set theory. Journal of Formalized Mathematics, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.
- [16] Andrzej Trybulec. Moore-Smith convergence. Journal of Formalized Mathematics, 8, 1996. http://mizar.org/JFM/Vol8/yellow_6.html.
- [17] Andrzej Trybulec. Scott topology. Journal of Formalized Mathematics, 9, 1997. http://mizar.org/JFM/Vo19/waybell1.html.
- [18] Wojciech A. Trybulec. Partially ordered sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/orders_1.html.
- [19] Wojciech A. Trybulec. Groups. Journal of Formalized Mathematics, 2, 1990. http://mizar.org/JFM/Vo12/group_1.html.
- [20] Zinaida Trybulec. Properties of subsets. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [21] Edmund Woronowicz. Relations and their basic properties. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/relat_1.html.
- [22] Edmund Woronowicz. Relations defined on sets. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/relset_1.html.

[23] Mariusz Żynel and Adam Guzowski. T_0 topological spaces. Journal of Formalized Mathematics, 6, 1994. http://mizar.org/JFM/Vol6/t_Otopsp.html.

Received May 23, 2000

Published January 2, 2004