

# Morphisms Into Chains. Part I

Artur Kornilowicz  
University of Białystok

**Summary.** This work is the continuation of formalization of [10]. Items from 2.1 to 2.8 of Chapter 4 are proved.

MML Identifier: WAYBEL35.

WWW: <http://mizar.org/JFM/Vol15/waybel35.html>

The articles [13], [7], [17], [14], [4], [15], [16], [1], [18], [20], [19], [5], [6], [2], [12], [21], [3], [8], [11], and [9] provide the notation and terminology for this paper.

## 1. PRELIMINARIES

Let  $X$  be a set. Note that there exists a subset of  $X$  which is trivial.

Let  $X$  be a trivial set. Observe that every subset of  $X$  is trivial.

Let  $L$  be a 1-sorted structure. Note that there exists a subset of  $L$  which is trivial.

Let  $L$  be a relational structure. Note that there exists a subset of  $L$  which is trivial.

Let  $L$  be a non empty 1-sorted structure. Note that there exists a subset of  $L$  which is non empty and trivial.

Let  $L$  be a non empty relational structure. Observe that there exists a subset of  $L$  which is non empty and trivial.

One can prove the following propositions:

- (1) For every set  $X$  holds  $\subseteq_X$  is reflexive in  $X$ .
- (2) For every set  $X$  holds  $\subseteq_X$  is transitive in  $X$ .
- (3) For every set  $X$  holds  $\subseteq_X$  is antisymmetric in  $X$ .

## 2. MAIN PART

Let  $L$  be a relational structure. One can verify that there exists a binary relation on  $L$  which is auxiliary(i).

Let  $L$  be a transitive relational structure. Note that there exists a binary relation on  $L$  which is auxiliary(i) and auxiliary(ii).

Let  $L$  be an antisymmetric relational structure with l.u.b.'s. Observe that there exists a binary relation on  $L$  which is auxiliary(iii).

Let  $L$  be a non empty lower-bounded antisymmetric relational structure. Note that there exists a binary relation on  $L$  which is auxiliary(iv).

Let  $L$  be a non empty relational structure and let  $R$  be a binary relation on  $L$ . We say that  $R$  is extra-order if and only if:

(Def. 1)  $R$  is auxiliary(i), auxiliary(ii), and auxiliary(iv).

Let  $L$  be a non empty relational structure. Note that every binary relation on  $L$  which is extra-order is also auxiliary(i), auxiliary(ii), and auxiliary(iv) and every binary relation on  $L$  which is auxiliary(i), auxiliary(ii), and auxiliary(iv) is also extra-order.

Let  $L$  be a non empty relational structure. Observe that every binary relation on  $L$  which is extra-order and auxiliary(iii) is also auxiliary and every binary relation on  $L$  which is auxiliary is also extra-order.

Let  $L$  be a lower-bounded antisymmetric transitive non empty relational structure. Note that there exists a binary relation on  $L$  which is extra-order.

Let  $L$  be a lower-bounded poset with l.u.b.'s and let  $R$  be an auxiliary(ii) binary relation on  $L$ . The functor  $R\text{-LowerMap}$  yields a map from  $L$  into  $\langle \text{LOWER}L, \subseteq \rangle$  and is defined as follows:

(Def. 2) For every element  $x$  of  $L$  holds  $R\text{-LowerMap}(x) = \downarrow_R x$ .

Let  $L$  be a lower-bounded poset with l.u.b.'s and let  $R$  be an auxiliary(ii) binary relation on  $L$ . Observe that  $R\text{-LowerMap}$  is monotone.

Let  $L$  be a 1-sorted structure and let  $R$  be a binary relation on the carrier of  $L$ . A subset of  $L$  is called a strict chain of  $R$  if:

(Def. 3) For all sets  $x, y$  such that  $x \in \text{it}$  and  $y \in \text{it}$  holds  $\langle x, y \rangle \in R$  or  $x = y$  or  $\langle y, x \rangle \in R$ .

We now state the proposition

(4) Let  $L$  be a 1-sorted structure,  $C$  be a trivial subset of  $L$ , and  $R$  be a binary relation on the carrier of  $L$ . Then  $C$  is a strict chain of  $R$ .

Let  $L$  be a non empty 1-sorted structure and let  $R$  be a binary relation on the carrier of  $L$ . Observe that there exists a strict chain of  $R$  which is non empty and trivial.

Next we state four propositions:

(5) Let  $L$  be an antisymmetric relational structure,  $R$  be an auxiliary(i) binary relation on  $L$ ,  $C$  be a strict chain of  $R$ , and  $x, y$  be elements of  $L$ . If  $x \in C$  and  $y \in C$  and  $x < y$ , then  $\langle x, y \rangle \in R$ .

(6) Let  $L$  be an antisymmetric relational structure,  $R$  be an auxiliary(i) binary relation on  $L$ , and  $x, y$  be elements of  $L$ . If  $\langle x, y \rangle \in R$  and  $\langle y, x \rangle \in R$ , then  $x = y$ .

(7) Let  $L$  be a non empty lower-bounded antisymmetric relational structure,  $x$  be an element of  $L$ , and  $R$  be an auxiliary(iv) binary relation on  $L$ . Then  $\{\perp_L, x\}$  is a strict chain of  $R$ .

(8) Let  $L$  be a non empty lower-bounded antisymmetric relational structure,  $R$  be an auxiliary(iv) binary relation on  $L$ , and  $C$  be a strict chain of  $R$ . Then  $C \cup \{\perp_L\}$  is a strict chain of  $R$ .

Let  $L$  be a 1-sorted structure, let  $R$  be a binary relation on the carrier of  $L$ , and let  $C$  be a strict chain of  $R$ . We say that  $C$  is maximal if and only if:

(Def. 4) For every strict chain  $D$  of  $R$  such that  $C \subseteq D$  holds  $C = D$ .

Let  $L$  be a 1-sorted structure, let  $R$  be a binary relation on the carrier of  $L$ , and let  $C$  be a set. The functor  $\text{StrictChains}(R, C)$  is defined by:

(Def. 5) For every set  $x$  holds  $x \in \text{StrictChains}(R, C)$  iff  $x$  is a strict chain of  $R$  and  $C \subseteq x$ .

Let  $L$  be a 1-sorted structure, let  $R$  be a binary relation on the carrier of  $L$ , and let  $C$  be a strict chain of  $R$ . Note that  $\text{StrictChains}(R, C)$  is non empty.

Let  $R$  be a binary relation and let  $X$  be a set. We introduce  $X$  is inductive w.r.t.  $R$  as a synonym of  $X$  has the upper Zorn property w.r.t.  $R$ .

We now state several propositions:

(9) Let  $L$  be a 1-sorted structure,  $R$  be a binary relation on the carrier of  $L$ , and  $C$  be a strict chain of  $R$ . Then  $\text{StrictChains}(R, C)$  is inductive w.r.t.  $\subseteq_{\text{StrictChains}(R, C)}$  and there exists a set  $D$  such that  $D$  is maximal in  $\subseteq_{\text{StrictChains}(R, C)}$  and  $C \subseteq D$ .

- (10) Let  $L$  be a non empty transitive relational structure,  $C$  be a non empty subset of  $L$ , and  $X$  be a subset of  $C$ . Suppose  $\sup X$  exists in  $L$  and  $\sqcup_L X \in C$ . Then  $\sup X$  exists in  $\text{sub}(C)$  and  $\sqcup_L X = \sqcup_{\text{sub}(C)} X$ .
- (11) Let  $L$  be a non empty poset,  $R$  be an auxiliary(i) auxiliary(ii) binary relation on  $L$ ,  $C$  be a non empty strict chain of  $R$ , and  $X$  be a subset of  $C$ . If  $\sup X$  exists in  $L$  and  $C$  is maximal, then  $\sup X$  exists in  $\text{sub}(C)$ .
- (12) Let  $L$  be a non empty poset,  $R$  be an auxiliary(i) auxiliary(ii) binary relation on  $L$ ,  $C$  be a non empty strict chain of  $R$ , and  $X$  be a subset of  $C$ . Suppose  $\inf \uparrow \sqcup_L X \cap C$  exists in  $L$  and  $\sup X$  exists in  $L$  and  $C$  is maximal. Then  $\sqcup_{\text{sub}(C)} X = \prod_L(\uparrow \sqcup_L X \cap C)$  and if  $\sqcup_L X \notin C$ , then  $\sqcup_L X < \prod_L(\uparrow \sqcup_L X \cap C)$ .
- (13) Let  $L$  be a complete non empty poset,  $R$  be an auxiliary(i) auxiliary(ii) binary relation on  $L$ , and  $C$  be a non empty strict chain of  $R$ . If  $C$  is maximal, then  $\text{sub}(C)$  is complete.
- (14) Let  $L$  be a non empty lower-bounded antisymmetric relational structure,  $R$  be an auxiliary(iv) binary relation on  $L$ , and  $C$  be a strict chain of  $R$ . If  $C$  is maximal, then  $\perp_L \in C$ .
- (15) Let  $L$  be a non empty upper-bounded poset,  $R$  be an auxiliary(ii) binary relation on  $L$ ,  $C$  be a strict chain of  $R$ , and  $m$  be an element of  $L$ . Suppose  $C$  is maximal and  $m$  is a maximum of  $C$  and  $\langle m, \top_L \rangle \in R$ . Then  $\langle \top_L, \top_L \rangle \in R$  and  $m = \top_L$ .

Let  $L$  be a relational structure, let  $C$  be a set, and let  $R$  be a binary relation on the carrier of  $L$ . We say that  $R$  satisfies SIC on  $C$  if and only if the condition (Def. 6) is satisfied.

- (Def. 6) Let  $x, z$  be elements of  $L$ . Suppose  $x \in C$  and  $z \in C$  and  $\langle x, z \rangle \in R$  and  $x \neq z$ . Then there exists an element  $y$  of  $L$  such that  $y \in C$  and  $\langle x, y \rangle \in R$  and  $\langle y, z \rangle \in R$  and  $x \neq y$ .

Let  $L$  be a relational structure, let  $R$  be a binary relation on the carrier of  $L$ , and let  $C$  be a strict chain of  $R$ . We say that  $C$  satisfies SIC if and only if:

- (Def. 7)  $R$  satisfies SIC on  $C$ .

We introduce  $C$  satisfies the interpolation property and  $C$  satisfies the interpolation property as synonyms of  $C$  satisfies SIC.

Next we state the proposition

- (16) Let  $L$  be a relational structure,  $C$  be a set, and  $R$  be an auxiliary(i) binary relation on  $L$ . Suppose  $R$  satisfies SIC on  $C$ . Let  $x, z$  be elements of  $L$ . Suppose  $x \in C$  and  $z \in C$  and  $\langle x, z \rangle \in R$  and  $x \neq z$ . Then there exists an element  $y$  of  $L$  such that  $y \in C$  and  $\langle x, y \rangle \in R$  and  $\langle y, z \rangle \in R$  and  $x < y$ .

Let  $L$  be a relational structure and let  $R$  be a binary relation on the carrier of  $L$ . One can verify that every strict chain of  $R$  which is trivial satisfies also SIC.

Let  $L$  be a non empty relational structure and let  $R$  be a binary relation on the carrier of  $L$ . Observe that there exists a strict chain of  $R$  which is non empty and trivial.

One can prove the following proposition

- (17) Let  $L$  be a lower-bounded poset with l.u.b.'s,  $R$  be an auxiliary(i) auxiliary(ii) binary relation on  $L$ , and  $C$  be a strict chain of  $R$ . Suppose  $C$  is maximal and  $R$  satisfies strong interpolation property. Then  $R$  satisfies SIC on  $C$ .

Let  $R$  be a binary relation and let  $C, y$  be sets. The functor  $\text{SetBelow}(R, C, y)$  is defined by:

- (Def. 8)  $\text{SetBelow}(R, C, y) = R^{-1}(\{y\}) \cap C$ .

The following proposition is true

- (18) For every binary relation  $R$  and for all sets  $C, x, y$  holds  $x \in \text{SetBelow}(R, C, y)$  iff  $\langle x, y \rangle \in R$  and  $x \in C$ .

Let  $L$  be a 1-sorted structure, let  $R$  be a binary relation on the carrier of  $L$ , and let  $C, y$  be sets. Then  $\text{SetBelow}(R, C, y)$  is a subset of  $L$ .

One can prove the following propositions:

- (19) Let  $L$  be a relational structure,  $R$  be an auxiliary(i) binary relation on  $L$ ,  $C$  be a set, and  $y$  be an element of  $L$ . Then  $\text{SetBelow}(R, C, y) \leq y$ .
- (20) Let  $L$  be a reflexive transitive relational structure,  $R$  be an auxiliary(ii) binary relation on  $L$ ,  $C$  be a subset of  $L$ , and  $x, y$  be elements of  $L$ . If  $x \leq y$ , then  $\text{SetBelow}(R, C, x) \subseteq \text{SetBelow}(R, C, y)$ .
- (21) Let  $L$  be a relational structure,  $R$  be an auxiliary(i) binary relation on  $L$ ,  $C$  be a set, and  $x$  be an element of  $L$ . If  $x \in C$  and  $\langle x, x \rangle \in R$  and  $\sup \text{SetBelow}(R, C, x)$  exists in  $L$ , then  $x = \sup \text{SetBelow}(R, C, x)$ .

Let  $L$  be a relational structure and let  $C$  be a subset of  $L$ . We say that  $C$  is sup-closed if and only if:

(Def. 9) For every subset  $X$  of  $C$  such that  $\sup X$  exists in  $L$  holds  $\bigsqcup_L X = \bigsqcup_{\text{sub}(C)} X$ .

Next we state three propositions:

- (22) Let  $L$  be a complete non empty poset,  $R$  be an extra-order binary relation on  $L$ ,  $C$  be a strict chain of  $R$  satisfying SIC, and  $p, q$  be elements of  $L$ . Suppose  $p \in C$  and  $q \in C$  and  $p < q$ . Then there exists an element  $y$  of  $L$  such that  $p < y$  and  $\langle y, q \rangle \in R$  and  $y = \sup \text{SetBelow}(R, C, y)$ .
- (23) Let  $L$  be a lower-bounded non empty poset,  $R$  be an extra-order binary relation on  $L$ , and  $C$  be a non empty strict chain of  $R$ . Suppose that
  - (i)  $C$  is sup-closed,
  - (ii) for every element  $c$  of  $L$  such that  $c \in C$  holds  $\sup \text{SetBelow}(R, C, c)$  exists in  $L$ , and
  - (iii)  $R$  satisfies SIC on  $C$ .

Let  $c$  be an element of  $L$ . If  $c \in C$ , then  $c = \sup \text{SetBelow}(R, C, c)$ .

- (24) Let  $L$  be a non empty reflexive antisymmetric relational structure,  $R$  be an auxiliary(i) binary relation on  $L$ , and  $C$  be a strict chain of  $R$ . Suppose that for every element  $c$  of  $L$  such that  $c \in C$  holds  $\sup \text{SetBelow}(R, C, c)$  exists in  $L$  and  $c = \sup \text{SetBelow}(R, C, c)$ . Then  $R$  satisfies SIC on  $C$ .

Let  $L$  be a non empty relational structure, let  $R$  be a binary relation on the carrier of  $L$ , and let  $C$  be a set. The functor  $\text{SupBelow}(R, C)$  is defined as follows:

(Def. 10) For every set  $y$  holds  $y \in \text{SupBelow}(R, C)$  iff  $y = \sup \text{SetBelow}(R, C, y)$ .

Let  $L$  be a non empty relational structure, let  $R$  be a binary relation on the carrier of  $L$ , and let  $C$  be a set. Then  $\text{SupBelow}(R, C)$  is a subset of  $L$ .

One can prove the following propositions:

- (25) Let  $L$  be a non empty reflexive transitive relational structure,  $R$  be an auxiliary(i) auxiliary(ii) binary relation on  $L$ , and  $C$  be a strict chain of  $R$ . Suppose that for every element  $c$  of  $L$  holds  $\sup \text{SetBelow}(R, C, c)$  exists in  $L$ . Then  $\text{SupBelow}(R, C)$  is a strict chain of  $R$ .
- (26) Let  $L$  be a non empty poset,  $R$  be an auxiliary(i) auxiliary(ii) binary relation on  $L$ , and  $C$  be a subset of  $L$ . Suppose that for every element  $c$  of  $L$  holds  $\sup \text{SetBelow}(R, C, c)$  exists in  $L$ . Then  $\text{SupBelow}(R, C)$  is sup-closed.
- (27) Let  $L$  be a complete non empty poset,  $R$  be an extra-order binary relation on  $L$ ,  $C$  be a strict chain of  $R$  satisfying SIC, and  $d$  be an element of  $L$ . If  $d \in \text{SupBelow}(R, C)$ , then  $d = \bigsqcup_L \{b; b \text{ ranges over elements of } L: b \in \text{SupBelow}(R, C) \wedge \langle b, d \rangle \in R\}$ .

- (28) Let  $L$  be a complete non empty poset,  $R$  be an extra-order binary relation on  $L$ , and  $C$  be a strict chain of  $R$  satisfying SIC. Then  $R$  satisfies SIC on  $\text{SupBelow}(R, C)$ .
- (29) Let  $L$  be a complete non empty poset,  $R$  be an extra-order binary relation on  $L$ ,  $C$  be a strict chain of  $R$  satisfying SIC, and  $a, b$  be elements of  $L$ . Suppose  $a \in C$  and  $b \in C$  and  $a < b$ . Then there exists an element  $d$  of  $L$  such that  $d \in \text{SupBelow}(R, C)$  and  $a < d$  and  $\langle d, b \rangle \in R$ .

## REFERENCES

- [1] Grzegorz Bancerek. Complete lattices. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/lattice3.html>.
- [2] Grzegorz Bancerek. Bounds in posets and relational substructures. *Journal of Formalized Mathematics*, 8, 1996. [http://mizar.org/JFM/Vol8/yellow\\_0.html](http://mizar.org/JFM/Vol8/yellow_0.html).
- [3] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. *Journal of Formalized Mathematics*, 8, 1996. [http://mizar.org/JFM/Vol8/waybel\\_0.html](http://mizar.org/JFM/Vol8/waybel_0.html).
- [4] Józef Białas. Group and field definitions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/realset1.html>.
- [5] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/funct\\_1.html](http://mizar.org/JFM/Vol1/funct_1.html).
- [6] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/funct\\_2.html](http://mizar.org/JFM/Vol1/funct_2.html).
- [7] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/zfmisc\\_1.html](http://mizar.org/JFM/Vol1/zfmisc_1.html).
- [8] Czesław Byliński. Galois connections. *Journal of Formalized Mathematics*, 8, 1996. [http://mizar.org/JFM/Vol8/waybel\\_1.html](http://mizar.org/JFM/Vol8/waybel_1.html).
- [9] Marek Dudzicz. Representation theorem for finite distributive lattices. *Journal of Formalized Mathematics*, 12, 2000. <http://mizar.org/JFM/Vol12/lattice7.html>.
- [10] G. Gierz, K.H. Hofmann, K. Keimel, J.D. Lawson, M. Mislove, and D.S. Scott. *A Compendium of Continuous Lattices*. Springer-Verlag, Berlin, Heidelberg, New York, 1980.
- [11] Adam Grabowski. Auxiliary and approximating relations. *Journal of Formalized Mathematics*, 8, 1996. [http://mizar.org/JFM/Vol8/waybel\\_4.html](http://mizar.org/JFM/Vol8/waybel_4.html).
- [12] Adam Grabowski and Robert Milewski. Boolean posets, posets under inclusion and products of relational structures. *Journal of Formalized Mathematics*, 8, 1996. [http://mizar.org/JFM/Vol8/yellow\\_1.html](http://mizar.org/JFM/Vol8/yellow_1.html).
- [13] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [14] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. <http://mizar.org/JFM/Addenda/numbers.html>.
- [15] Wojciech A. Trybulec. Partially ordered sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/orders\\_1.html](http://mizar.org/JFM/Vol1/orders_1.html).
- [16] Wojciech A. Trybulec and Grzegorz Bancerek. Kuratowski - Zorn lemma. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/orders\\_2.html](http://mizar.org/JFM/Vol1/orders_2.html).
- [17] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/subset\\_1.html](http://mizar.org/JFM/Vol1/subset_1.html).
- [18] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/relat\\_1.html](http://mizar.org/JFM/Vol1/relat_1.html).
- [19] Edmund Woronowicz. Relations defined on sets. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/relset\\_1.html](http://mizar.org/JFM/Vol1/relset_1.html).
- [20] Edmund Woronowicz and Anna Zalewska. Properties of binary relations. *Journal of Formalized Mathematics*, 1, 1989. [http://mizar.org/JFM/Vol1/relat\\_2.html](http://mizar.org/JFM/Vol1/relat_2.html).
- [21] Mariusz Żynel and Czesław Byliński. Properties of relational structures, posets, lattices and maps. *Journal of Formalized Mathematics*, 8, 1996. [http://mizar.org/JFM/Vol8/yellow\\_2.html](http://mizar.org/JFM/Vol8/yellow_2.html).

*Received February 6, 2003*

*Published January 2, 2004*