Directed Sets, Nets, Ideals, Filters, and Maps¹

Grzegorz Bancerek Institute of Mathematics Polish Academy of Sciences

Summary. Notation and facts necessary to start with the formalization of continuous lattices according to [8] are introduced. The article contains among other things, the definition of directed and filtered subsets of a poset (see 1.1 in [8, p. 2]), the definition of nets on the poset (see 1.2 in [8, p. 2]), the definition of ideals and filters and the definition of maps preserving arbitrary and directed sups and arbitrary and filtered infs (1.9 also in [8, p. 4]). The concepts of semilattices, sup-semiletices and poset lattices (1.8 in [8, p. 4]) are also introduced. A number of facts concerning the above notion and including remarks 1.4, 1.5, and 1.10 from [8, pp. 3–5] is presented.

MML Identifier: WAYBEL_0.

WWW: http://mizar.org/JFM/Vol8/waybel_0.html

The articles [12], [6], [14], [15], [17], [16], [7], [4], [5], [11], [3], [10], [1], [2], [13], and [9] provide the notation and terminology for this paper.

1. DIRECTED SUBSETS

Let L be a relational structure and let X be a subset of L. We say that X is directed if and only if:

(Def. 1) For all elements x, y of L such that $x \in X$ and $y \in X$ there exists an element z of L such that $z \in X$ and $x \le z$ and $y \le z$.

We say that *X* is filtered if and only if:

(Def. 2) For all elements x, y of L such that $x \in X$ and $y \in X$ there exists an element z of L such that $z \in X$ and $z \le x$ and $z \le y$.

One can prove the following two propositions:

- (1) Let L be a non empty transitive relational structure and X be a subset of L. Then X is non empty and directed if and only if for every finite subset Y of X there exists an element x of L such that $x \in X$ and $x \ge Y$.
- (2) Let L be a non empty transitive relational structure and X be a subset of L. Then X is non empty and filtered if and only if for every finite subset Y of X there exists an element x of L such that $x \in X$ and $x \le Y$.

Let *L* be a relational structure. Observe that \emptyset_L is directed and filtered.

Let L be a relational structure. One can verify that there exists a subset of L which is directed and filtered.

We now state three propositions:

1

¹This work has been partially supported by Office of Naval Research Grant N00014-95-1-1336.

- (3) Let L_1 , L_2 be relational structures. Suppose the relational structure of L_1 = the relational structure of L_2 . Let X_1 be a subset of L_1 and X_2 be a subset of L_2 . If $X_1 = X_2$ and X_1 is directed, then X_2 is directed.
- (4) Let L_1 , L_2 be relational structures. Suppose the relational structure of L_1 = the relational structure of L_2 . Let X_1 be a subset of L_1 and X_2 be a subset of L_2 . If $X_1 = X_2$ and X_1 is filtered, then X_2 is filtered.
- (5) For every non empty reflexive relational structure L and for every element x of L holds $\{x\}$ is directed and filtered.

Let L be a non empty reflexive relational structure. One can verify that there exists a subset of L which is directed, filtered, non empty, and finite.

Let L be a relational structure with l.u.b.'s. Observe that Ω_L is directed.

Let L be an upper-bounded non empty relational structure. Note that Ω_L is directed.

Let L be a relational structure with g.l.b.'s. Observe that Ω_L is filtered.

Let L be a lower-bounded non empty relational structure. Note that Ω_L is filtered.

Let *L* be a non empty relational structure and let *S* be a relational substructure of *L*. We say that *S* is filtered-infs-inheriting if and only if:

(Def. 3) For every filtered subset X of S such that $X \neq \emptyset$ and inf X exists in L holds $\bigcap_L X \in$ the carrier of S.

We say that *S* is directed-sups-inheriting if and only if:

(Def. 4) For every directed subset X of S such that $X \neq \emptyset$ and sup X exists in L holds $\bigsqcup_L X \in$ the carrier of S.

Let L be a non empty relational structure. Note that every relational substructure of L which is infs-inheriting is also filtered-infs-inheriting and every relational substructure of L which is sups-inheriting is also directed-sups-inheriting.

Let L be a non empty relational structure. Observe that there exists a relational substructure of L which is infs-inheriting, sups-inheriting, non empty, full, and strict.

Next we state two propositions:

- (6) Let *L* be a non empty transitive relational structure, *S* be a filtered-infs-inheriting non empty full relational substructure of *L*, and *X* be a filtered subset of *S*. Suppose $X \neq \emptyset$ and inf *X* exists in *L*. Then inf *X* exists in *S* and $\bigcap_S X = \bigcap_L X$.
- (7) Let L be a non empty transitive relational structure, S be a directed-sups-inheriting non empty full relational substructure of L, and X be a directed subset of S. Suppose $X \neq \emptyset$ and sup X exists in L. Then sup X exists in S and $\bigsqcup_{S} X = \bigsqcup_{L} X$.

2. Nets

Let L_1 , L_2 be non empty 1-sorted structures, let f be a map from L_1 into L_2 , and let x be an element of L_1 . Then f(x) is an element of L_2 .

Let L_1 , L_2 be relational structures and let f be a map from L_1 into L_2 . We say that f is antitone if and only if:

(Def. 5) For all elements x, y of L_1 such that $x \le y$ and for all elements a, b of L_2 such that a = f(x) and b = f(y) holds $a \ge b$.

Let L be a 1-sorted structure. We consider net structures over L as extensions of relational structure as systems

⟨ a carrier, an internal relation, a mapping ⟩,

where the carrier is a set, the internal relation is a binary relation on the carrier, and the mapping is a function from the carrier into the carrier of L.

Let *L* be a 1-sorted structure, let *X* be a non empty set, let *O* be a binary relation on *X*, and let *F* be a function from *X* into the carrier of *L*. Note that $\langle X, O, F \rangle$ is non empty.

Let *N* be a relational structure. We say that *N* is directed if and only if:

(Def. 6) Ω_N is directed.

Let L be a 1-sorted structure. One can verify that there exists a strict net structure over L which is non empty, reflexive, transitive, antisymmetric, and directed.

Let L be a 1-sorted structure. A prenet over L is a directed non empty net structure over L.

Let L be a 1-sorted structure. A net in L is a transitive prenet over L.

Let L be a non empty 1-sorted structure and let N be a non empty net structure over L. The functor netmap(N,L) yielding a map from N into L is defined as follows:

(Def. 7) netmap(N, L) = the mapping of N.

Let i be an element of N. The functor N(i) yielding an element of L is defined by:

(Def. 8) N(i) =(the mapping of N)(i).

Let L be a non empty relational structure and let N be a non empty net structure over L. We say that N is monotone if and only if:

(Def. 9) netmap(N,L) is monotone.

We say that *N* is antitone if and only if:

(Def. 10) netmap(N, L) is antitone.

Let L be a non empty 1-sorted structure, let N be a non empty net structure over L, and let X be a set. We say that N is eventually in X if and only if:

(Def. 11) There exists an element i of N such that for every element j of N such that $i \le j$ holds $N(j) \in X$.

We say that N is often in X if and only if:

(Def. 12) For every element i of N there exists an element j of N such that $i \le j$ and $N(j) \in X$.

We now state three propositions:

- (8) Let L be a non empty 1-sorted structure, N be a non empty net structure over L, and X, Y be sets such that $X \subseteq Y$. Then
- (i) if N is eventually in X, then N is eventually in Y, and
- (ii) if N is often in X, then N is often in Y.
- (9) Let *L* be a non empty 1-sorted structure, *N* be a non empty net structure over *L*, and *X* be a set. Then *N* is eventually in *X* if and only if *N* is not often in (the carrier of *L*) $\setminus X$.
- (10) Let L be a non empty 1-sorted structure, N be a non empty net structure over L, and X be a set. Then N is often in X if and only if N is not eventually in (the carrier of L) $\setminus X$.

Let L be a non empty relational structure and let N be a non empty net structure over L. We say that N is eventually-directed if and only if:

(Def. 13) For every element i of N holds N is eventually in $\{N(j); j \text{ ranges over elements of } N: N(i) \le N(j)\}$.

We say that *N* is eventually-filtered if and only if:

(Def. 14) For every element i of N holds N is eventually in $\{N(j); j \text{ ranges over elements of } N: N(i) \ge N(j)\}$.

The following two propositions are true:

(11) Let L be a non empty relational structure and N be a non empty net structure over L. Then N is eventually-directed if and only if for every element i of N there exists an element j of N such that for every element k of N such that $j \le k$ holds $N(i) \le N(k)$.

(12) Let L be a non empty relational structure and N be a non empty net structure over L. Then N is eventually-filtered if and only if for every element i of N there exists an element j of N such that for every element k of N such that $j \le k$ holds $N(i) \ge N(k)$.

Let L be a non empty relational structure. One can verify that every prenet over L which is monotone is also eventually-directed and every prenet over L which is antitone is also eventually-filtered.

Let L be a non empty reflexive relational structure. Observe that there exists a prenet over L which is monotone, antitone, and strict.

3. Lower and upper subsets

Let *L* be a relational structure and let *X* be a subset of *L*. The functor $\downarrow X$ yields a subset of *L* and is defined by:

(Def. 15) For every element x of L holds $x \in JX$ iff there exists an element y of L such that $y \ge x$ and $y \in X$.

The functor $\uparrow X$ yielding a subset of *L* is defined by:

(Def. 16) For every element x of L holds $x \in \uparrow X$ iff there exists an element y of L such that $y \le x$ and $y \in X$.

One can prove the following propositions:

- (13) Let L_1 , L_2 be relational structures. Suppose the relational structure of L_1 = the relational structure of L_2 . Let X be a subset of L_1 and Y be a subset of L_2 . If X = Y, then $\downarrow X = \downarrow Y$ and $\uparrow X = \uparrow Y$.
- (14) Let *L* be a non empty relational structure and *X* be a subset of *L*. Then $\downarrow X = \{x; x \text{ ranges over elements of } L: \bigvee_{y:\text{element of } L} (x \leq y \land y \in X)\}.$
- (15) Let L be a non empty relational structure and X be a subset of L. Then $\uparrow X = \{x; x \text{ ranges over elements of } L: \bigvee_{y:\text{element of } L} (x \ge y \land y \in X)\}.$

Let L be a non empty reflexive relational structure and let X be a non empty subset of L. Note that $\downarrow X$ is non empty and $\uparrow X$ is non empty.

One can prove the following proposition

(16) For every reflexive relational structure L and for every subset X of L holds $X \subseteq J X$ and $X \subseteq J X$.

Let *L* be a non empty relational structure and let *x* be an element of *L*. The functor $\downarrow x$ yielding a subset of *L* is defined as follows:

(Def. 17) $\downarrow x = \downarrow \{x\}.$

The functor $\uparrow x$ yields a subset of L and is defined as follows:

(Def. 18) $\uparrow x = \uparrow \{x\}.$

One can prove the following propositions:

- (17) For every non empty relational structure L and for all elements x, y of L holds $y \in \downarrow x$ iff $y \in x$
- (18) For every non empty relational structure *L* and for all elements *x*, *y* of *L* holds $y \in \uparrow x$ iff $x \le y$.
- (19) Let *L* be a non empty reflexive antisymmetric relational structure and *x*, *y* be elements of *L*. If $\downarrow x = \downarrow y$, then x = y.

- (20) Let *L* be a non empty reflexive antisymmetric relational structure and *x*, *y* be elements of *L*. If $\uparrow x = \uparrow y$, then x = y.
- (21) For every non empty transitive relational structure L and for all elements x, y of L such that $x \le y$ holds $\downarrow x \subseteq \downarrow y$.
- (22) For every non empty transitive relational structure L and for all elements x, y of L such that $x \le y$ holds $\uparrow y \subseteq \uparrow x$.

Let *L* be a non empty reflexive relational structure and let *x* be an element of *L*. One can check that $\downarrow x$ is non empty and directed and $\uparrow x$ is non empty and filtered.

Let L be a relational structure and let X be a subset of L. We say that X is lower if and only if:

(Def. 19) For all elements x, y of L such that $x \in X$ and $y \le x$ holds $y \in X$.

We say that *X* is upper if and only if:

(Def. 20) For all elements x, y of L such that $x \in X$ and $x \le y$ holds $y \in X$.

Let *L* be a relational structure. Note that there exists a subset of *L* which is lower and upper. We now state several propositions:

- (23) For every relational structure L and for every subset X of L holds X is lower iff $\bigcup X \subseteq X$.
- (24) For every relational structure L and for every subset X of L holds X is upper iff $\uparrow X \subseteq X$.
- (25) Let L_1 , L_2 be relational structures. Suppose the relational structure of L_1 = the relational structure of L_2 . Let X_1 be a subset of L_1 and X_2 be a subset of L_2 such that $X_1 = X_2$. Then
 - (i) if X_1 is lower, then X_2 is lower, and
- (ii) if X_1 is upper, then X_2 is upper.
- (26) Let L be a relational structure and A be a subset of $2^{\text{the carrier of } L}$. Suppose that for every subset X of L such that $X \in A$ holds X is lower. Then $\bigcup A$ is a lower subset of L.
- (27) Let *L* be a relational structure and *X*, *Y* be subsets of *L*. If *X* is lower and *Y* is lower, then $X \cap Y$ is lower and $X \cup Y$ is lower.
- (28) Let L be a relational structure and A be a subset of $2^{\text{the carrier of } L}$. Suppose that for every subset X of L such that $X \in A$ holds X is upper. Then $\bigcup A$ is an upper subset of L.
- (29) Let *L* be a relational structure and *X*, *Y* be subsets of *L*. If *X* is upper and *Y* is upper, then $X \cap Y$ is upper and $X \cup Y$ is upper.

Let *L* be a non empty transitive relational structure and let *X* be a subset of *L*. Note that $\downarrow X$ is lower and $\uparrow X$ is upper.

Let *L* be a non empty transitive relational structure and let *x* be an element of *L*. Note that $\downarrow x$ is lower and $\uparrow x$ is upper.

Let L be a non empty relational structure. Observe that Ω_L is lower and upper.

Let L be a non empty relational structure. Note that there exists a subset of L which is non empty, lower, and upper.

Let L be a non empty reflexive transitive relational structure. Observe that there exists a subset of L which is non empty, lower, and directed and there exists a subset of L which is non empty, upper, and filtered.

Let L be a poset with g.l.b.'s and l.u.b.'s. One can check that there exists a subset of L which is non empty, directed, filtered, lower, and upper.

Next we state the proposition

(30) Let L be a non empty transitive reflexive relational structure and X be a subset of L. Then X is directed if and only if $\downarrow X$ is directed.

Let L be a non empty transitive reflexive relational structure and let X be a directed subset of L. One can check that $\downarrow X$ is directed.

The following propositions are true:

- (31) Let *L* be a non empty transitive reflexive relational structure, *X* be a subset of *L*, and *x* be an element of *L*. Then $x \ge X$ if and only if $x \ge \bigcup X$.
- (32) Let L be a non empty transitive reflexive relational structure and X be a subset of L. Then $\sup X$ exists in L if and only if $\sup \downarrow X$ exists in L.
- (33) Let L be a non empty transitive reflexive relational structure and X be a subset of L. If sup X exists in L, then $\sup X = \sup X$.
- (34) For every non empty poset *L* and for every element *x* of *L* holds sup $\downarrow x$ exists in *L* and $\sup \downarrow x = x$.
- (35) Let L be a non empty transitive reflexive relational structure and X be a subset of L. Then X is filtered if and only if $\uparrow X$ is filtered.

Let *L* be a non empty transitive reflexive relational structure and let *X* be a filtered subset of *L*. One can verify that $\uparrow X$ is filtered.

We now state four propositions:

- (36) Let *L* be a non empty transitive reflexive relational structure, *X* be a subset of *L*, and *x* be an element of *L*. Then $x \le X$ if and only if $x \le \uparrow X$.
- (37) Let *L* be a non empty transitive reflexive relational structure and *X* be a subset of *L*. Then inf *X* exists in *L* if and only if inf $\uparrow X$ exists in *L*.
- (38) Let L be a non empty transitive reflexive relational structure and X be a subset of L. If inf X exists in L, then $\inf X = \inf \uparrow X$.
- (39) For every non empty poset L and for every element x of L holds inf $\uparrow x$ exists in L and $\inf \uparrow x = x$.

4. IDEALS AND FILTERS

Let L be a non empty reflexive transitive relational structure. An ideal of L is a directed lower non empty subset of L. A filter of L is a filtered upper non empty subset of L.

One can prove the following propositions:

- (40) Let L be an antisymmetric relational structure with l.u.b.'s and X be a lower subset of L. Then X is directed if and only if for all elements x, y of L such that $x \in X$ and $y \in X$ holds $x \sqcup y \in X$.
- (41) Let L be an antisymmetric relational structure with g.l.b.'s and X be an upper subset of L. Then X is filtered if and only if for all elements x, y of L such that $x \in X$ and $y \in X$ holds $x \sqcap y \in X$.
- (42) Let *L* be a poset with l.u.b.'s and *X* be a non empty lower subset of *L*. Then *X* is directed if and only if for every finite subset *Y* of *X* such that $Y \neq \emptyset$ holds $\bigsqcup_L Y \in X$.
- (43) Let *L* be a poset with g.l.b.'s and *X* be a non empty upper subset of *L*. Then *X* is filtered if and only if for every finite subset *Y* of *X* such that $Y \neq \emptyset$ holds $\bigcap_L Y \in X$.
- (44) Let L be a non empty antisymmetric relational structure. Suppose L has l.u.b.'s and g.l.b.'s. Let X, Y be subsets of L. Suppose X is lower and directed and Y is lower and directed. Then $X \cap Y$ is directed.

- (45) Let L be a non empty antisymmetric relational structure. Suppose L has l.u.b.'s and g.l.b.'s. Let X, Y be subsets of L. Suppose X is upper and filtered and Y is upper and filtered. Then $X \cap Y$ is filtered.
- (46) Let L be a relational structure and A be a subset of $2^{\text{the carrier of } L}$. Suppose that
 - (i) for every subset X of L such that $X \in A$ holds X is directed, and
- (ii) for all subsets X, Y of L such that $X \in A$ and $Y \in A$ there exists a subset Z of L such that $Z \in A$ and $X \cup Y \subseteq Z$.

Let X be a subset of L. If $X = \bigcup A$, then X is directed.

- (47) Let L be a relational structure and A be a subset of $2^{\text{the carrier of } L}$. Suppose that
 - (i) for every subset X of L such that $X \in A$ holds X is filtered, and
- (ii) for all subsets X, Y of L such that $X \in A$ and $Y \in A$ there exists a subset Z of L such that $Z \in A$ and $X \cup Y \subseteq Z$.

Let *X* be a subset of *L*. If $X = \bigcup A$, then *X* is filtered.

Let L be a non empty reflexive transitive relational structure and let I be an ideal of L. We say that I is principal if and only if:

(Def. 21) There exists an element x of L such that $x \in I$ and $x \ge I$.

Let L be a non empty reflexive transitive relational structure and let F be a filter of L. We say that F is principal if and only if:

(Def. 22) There exists an element x of L such that $x \in F$ and $x \le F$.

We now state two propositions:

- (48) Let *L* be a non empty reflexive transitive relational structure and *I* be an ideal of *L*. Then *I* is principal if and only if there exists an element *x* of *L* such that $I = \downarrow x$.
- (49) Let L be a non empty reflexive transitive relational structure and F be a filter of L. Then F is principal if and only if there exists an element x of L such that $F = \uparrow x$.

Let L be a non empty reflexive transitive relational structure. The functor Ids(L) yielding a set is defined as follows:

(Def. 23) $Ids(L) = \{X : X \text{ ranges over ideals of } L\}.$

The functor Filt(L) yields a set and is defined by:

(Def. 24) Filt(L) = {X : X ranges over filters of L}.

Let L be a non empty reflexive transitive relational structure. The functor $\mathrm{Ids}_0(L)$ yields a set and is defined by:

(Def. 25) $Ids_0(L) = Ids(L) \cup \{\emptyset\}.$

The functor $Filt_0(L)$ yielding a set is defined by:

(Def. 26) $\operatorname{Filt}_0(L) = \operatorname{Filt}(L) \cup \{\emptyset\}.$

Let L be a non empty relational structure and let X be a subset of L. The functor finsups(X) yields a subset of L and is defined by:

(Def. 27) finsups(X) = { $\bigcup_L Y$; Y ranges over finite subsets of X: sup Y exists in L}.

The functor fininfs(X) yielding a subset of L is defined by:

(Def. 28) fininfs(X) = { $\bigcap_L Y$; Y ranges over finite subsets of X: inf Y exists in L}.

Let L be a non empty antisymmetric lower-bounded relational structure and let X be a subset of L. Note that finsups(X) is non empty.

Let L be a non empty antisymmetric upper-bounded relational structure and let X be a subset of L. One can verify that fininfs(X) is non empty.

Let L be a non empty reflexive antisymmetric relational structure and let X be a non empty subset of L. Observe that finsups(X) is non empty and fininfs(X) is non empty.

One can prove the following propositions:

- (50) For every non empty reflexive antisymmetric relational structure L and for every subset X of L holds $X \subseteq \text{finsups}(X)$ and $X \subseteq \text{fininfs}(X)$.
- (51) Let L be a non empty transitive relational structure and X, F be subsets of L. Suppose that
 - (i) for every finite subset Y of X such that $Y \neq \emptyset$ holds sup Y exists in L,
- (ii) for every element x of L such that $x \in F$ there exists a finite subset Y of X such that sup Y exists in L and $x = \bigsqcup_{L} Y$, and
- (iii) for every finite subset *Y* of *X* such that $Y \neq \emptyset$ holds $\bigsqcup_L Y \in F$. Then *F* is directed.

Let L be a poset with l.u.b.'s and let X be a subset of L. Note that finsups(X) is directed. One can prove the following propositions:

- (52) Let L be a non empty transitive reflexive relational structure and X, F be subsets of L. Suppose that
 - (i) for every finite subset Y of X such that $Y \neq \emptyset$ holds sup Y exists in L,
- (ii) for every element x of L such that $x \in F$ there exists a finite subset Y of X such that sup Y exists in L and $x = \bigcup_{L} Y$, and
- (iii) for every finite subset Y of X such that $Y \neq \emptyset$ holds $\bigsqcup_L Y \in F$. Let x be an element of L. Then $x \geq X$ if and only if $x \geq F$.
- (53) Let L be a non empty transitive reflexive relational structure and X, F be subsets of L. Suppose that
 - (i) for every finite subset Y of X such that $Y \neq \emptyset$ holds sup Y exists in L,
- (ii) for every element x of L such that $x \in F$ there exists a finite subset Y of X such that sup Y exists in L and $x = \bigsqcup_{L} Y$, and
- (iii) for every finite subset Y of X such that $Y \neq \emptyset$ holds $\bigsqcup_L Y \in F$. Then sup X exists in L if and only if sup F exists in L.
- (54) Let L be a non empty transitive reflexive relational structure and X, F be subsets of L. Suppose that
 - (i) for every finite subset Y of X such that $Y \neq \emptyset$ holds sup Y exists in L,
- (ii) for every element x of L such that $x \in F$ there exists a finite subset Y of X such that sup Y exists in L and $x = \bigcup_{L} Y$,
- (iii) for every finite subset Y of X such that $Y \neq \emptyset$ holds $\bigsqcup_L Y \in F$, and
- (iv) $\sup X$ exists in L. Then $\sup F = \sup X$.
- (55) Let L be a poset with l.u.b.'s and X be a subset of L. If sup X exists in L or L is complete, then sup $X = \sup \text{finsups}(X)$.

- (56) Let L be a non empty transitive relational structure and X, F be subsets of L. Suppose that
 - (i) for every finite subset Y of X such that $Y \neq \emptyset$ holds inf Y exists in L,
- (ii) for every element x of L such that $x \in F$ there exists a finite subset Y of X such that inf Y exists in L and $x = \bigcap_L Y$, and
- (iii) for every finite subset *Y* of *X* such that $Y \neq \emptyset$ holds $\bigcap_L Y \in F$.

Then F is filtered.

Let L be a poset with g.l.b.'s and let X be a subset of L. Observe that fininfs(X) is filtered. One can prove the following propositions:

- (57) Let L be a non empty transitive reflexive relational structure and X, F be subsets of L. Suppose that
 - (i) for every finite subset Y of X such that $Y \neq \emptyset$ holds inf Y exists in L,
- (ii) for every element x of L such that $x \in F$ there exists a finite subset Y of X such that inf Y exists in L and $x = \prod_{L} Y$, and
- (iii) for every finite subset Y of X such that $Y \neq \emptyset$ holds $\bigcap_L Y \in F$. Let X be an element of X. Then $X \leq X$ if and only if $X \leq Y$.
- (58) Let L be a non empty transitive reflexive relational structure and X, F be subsets of L. Suppose that
 - (i) for every finite subset Y of X such that $Y \neq \emptyset$ holds inf Y exists in L,
- (ii) for every element x of L such that $x \in F$ there exists a finite subset Y of X such that inf Y exists in L and $x = \bigcap_{L} Y$, and
- (iii) for every finite subset Y of X such that $Y \neq \emptyset$ holds $\bigcap_L Y \in F$. Then inf X exists in L if and only if inf F exists in L.
- (59) Let L be a non empty transitive reflexive relational structure and X, F be subsets of L. Suppose that
 - (i) for every finite subset Y of X such that $Y \neq \emptyset$ holds inf Y exists in L,
- (ii) for every element x of L such that $x \in F$ there exists a finite subset Y of X such that inf Y exists in L and $x = \prod_{L} Y$,
- (iii) for every finite subset Y of X such that $Y \neq \emptyset$ holds $\bigcap_L Y \in F$, and
- (iv) $\inf X$ exists in L.

Then $\inf F = \inf X$.

- (60) Let L be a poset with g.l.b.'s and X be a subset of L. If inf X exists in L or L is complete, then $\inf X = \inf \liminf S(X)$.
- (61) Let *L* be a poset with l.u.b.'s and *X* be a subset of *L*. Then $X \subseteq \downarrow \text{finsups}(X)$ and for every ideal *I* of *L* such that $X \subseteq I$ holds $\downarrow \text{finsups}(X) \subseteq I$.
- (62) Let *L* be a poset with g.l.b.'s and *X* be a subset of *L*. Then $X \subseteq \uparrow \text{fininfs}(X)$ and for every filter *F* of *L* such that $X \subseteq F$ holds $\uparrow \text{fininfs}(X) \subseteq F$.

5. CHAINS

Let *L* be a non empty relational structure. We say that *L* is connected if and only if:

(Def. 29) For all elements x, y of L holds $x \le y$ or $y \le x$.

Let us observe that every non empty reflexive relational structure which is trivial is also connected.

Let us note that there exists a non empty poset which is connected.

A chain is a connected non empty poset.

Let L be a chain. Note that L^{\smile} is connected.

6. SEMILATTICES

A semilattice is a poset with g.l.b.'s. A sup-semilattice is a poset with l.u.b.'s. A lattice is a poset with g.l.b.'s and l.u.b.'s.

The following propositions are true:

- (63) Let L be a semilattice and X be an upper non empty subset of L. Then X is a filter of L if and only if sub(X) is meet-inheriting.
- (64) Let L be a sup-semilattice and X be a lower non empty subset of L. Then X is an ideal of L if and only if sub(X) is join-inheriting.

7. Maps

Let S, T be non empty relational structures, let f be a map from S into T, and let X be a subset of S. We say that f preserves inf of X if and only if:

(Def. 30) If $\inf X$ exists in S, then $\inf f^{\circ}X$ exists in T and $\inf (f^{\circ}X) = f(\inf X)$.

We say that f preserves sup of X if and only if:

(Def. 31) If sup X exists in S, then sup $f^{\circ}X$ exists in T and sup $(f^{\circ}X) = f(\sup X)$.

We now state the proposition

- (65) Let S_1 , S_2 , T_1 , T_2 be non empty relational structures. Suppose that
 - (i) the relational structure of S_1 = the relational structure of T_1 , and
- (ii) the relational structure of S_2 = the relational structure of T_2 . Let f be a map from S_1 into S_2 and g be a map from T_1 into T_2 . Suppose f = g. Let X be a subset of S_1 and Y be a subset of T_1 such that X = Y. Then
- (iii) if f preserves sup of X, then g preserves sup of Y, and
- (iv) if f preserves inf of X, then g preserves inf of Y.

Let L_1 , L_2 be non empty relational structures and let f be a map from L_1 into L_2 . We say that f is infs-preserving if and only if:

(Def. 32) For every subset X of L_1 holds f preserves inf of X.

We say that f is sups-preserving if and only if:

(Def. 33) For every subset X of L_1 holds f preserves sup of X.

We say that f is meet-preserving if and only if:

(Def. 34) For all elements x, y of L_1 holds f preserves inf of $\{x, y\}$.

We say that f is join-preserving if and only if:

(Def. 35) For all elements x, y of L_1 holds f preserves sup of $\{x,y\}$.

We say that f is filtered-infs-preserving if and only if:

(Def. 36) For every subset X of L_1 such that X is non empty and filtered holds f preserves inf of X.

We say that f is directed-sups-preserving if and only if:

(Def. 37) For every subset X of L_1 such that X is non empty and directed holds f preserves sup of X.

Let L_1 , L_2 be non empty relational structures. Observe that every map from L_1 into L_2 which is infs-preserving is also filtered-infs-preserving and meet-preserving and every map from L_1 into L_2 which is sups-preserving is also directed-sups-preserving and join-preserving.

Let S, T be relational structures and let f be a map from S into T. We say that f is isomorphic if and only if:

- (Def. 38)(i) f is one-to-one and monotone and there exists a map g from T into S such that $g = f^{-1}$ and g is monotone if S is non empty and T is non empty,
 - (ii) S is empty and T is empty, otherwise.

Next we state the proposition

- (66) Let S, T be non empty relational structures and f be a map from S into T. Then f is isomorphic if and only if the following conditions are satisfied:
 - (i) f is one-to-one,
- (ii) $\operatorname{rng} f = \operatorname{the carrier of} T$, and
- (iii) for all elements x, y of S holds $x \le y$ iff $f(x) \le f(y)$.
- Let S, T be non empty relational structures. Observe that every map from S into T which is isomorphic is also one-to-one and monotone.

The following propositions are true:

- (67) Let S, T be non empty relational structures and f be a map from S into T. Suppose f is isomorphic. Then f^{-1} is a map from T into S and $rng(f^{-1}) = the$ carrier of S.
- (68) Let S, T be non empty relational structures and f be a map from S into T. Suppose f is isomorphic. Let g be a map from T into S. If $g = f^{-1}$, then g is isomorphic.
- (69) Let S, T be non empty posets and f be a map from S into T. Suppose that for every filter X of S holds f preserves inf of X. Then f is monotone.
- (70) Let S, T be non empty posets and f be a map from S into T. Suppose that for every filter X of S holds f preserves inf of X. Then f is filtered-infs-preserving.
- (71) Let S be a semilattice, T be a non empty poset, and f be a map from S into T. Suppose that
 - (i) for every finite subset X of S holds f preserves inf of X, and
- (ii) for every non empty filtered subset *X* of *S* holds *f* preserves inf of *X*. Then *f* is infs-preserving.
- (72) Let S, T be non empty posets and f be a map from S into T. Suppose that for every ideal X of S holds f preserves sup of X. Then f is monotone.
- (73) Let S, T be non empty posets and f be a map from S into T. Suppose that for every ideal X of S holds f preserves sup of X. Then f is directed-sups-preserving.
- (74) Let S be a sup-semilattice, T be a non empty poset, and f be a map from S into T. Suppose that
 - (i) for every finite subset X of S holds f preserves sup of X, and
- (ii) for every non empty directed subset X of S holds f preserves sup of X.

Then f is sups-preserving.

8. Completeness wrt directed sets

Let L be a non empty reflexive relational structure. We say that L is up-complete if and only if the condition (Def. 39) is satisfied.

(Def. 39) Let X be a non empty directed subset of L. Then there exists an element x of L such that $x \ge X$ and for every element y of L such that $y \ge X$ holds $x \le y$.

Let us mention that every reflexive relational structure with l.u.b.'s which is up-complete is also upper-bounded.

One can prove the following proposition

(75) Let L be a non empty reflexive antisymmetric relational structure. Then L is up-complete if and only if for every non empty directed subset X of L holds sup X exists in L.

Let L be a non empty reflexive relational structure. We say that L is inf-complete if and only if the condition (Def. 40) is satisfied.

(Def. 40) Let X be a non empty subset of L. Then there exists an element x of L such that $x \le X$ and for every element y of L such that $y \le X$ holds $x \ge y$.

Next we state the proposition

(76) Let L be a non empty reflexive antisymmetric relational structure. Then L is inf-complete if and only if for every non empty subset X of L holds inf X exists in L.

One can check the following observations:

- * every non empty reflexive relational structure which is complete is also up-complete and inf-complete,
- * every non empty reflexive relational structure which is inf-complete is also lower-bounded, and
- * every non empty poset which is up-complete and lower-bounded and has l.u.b.'s is also complete.

One can verify that every non empty reflexive antisymmetric relational structure which is infcomplete has also g.l.b.'s.

One can verify that every non empty reflexive antisymmetric upper-bounded relational structure which is inf-complete has also l.u.b.'s.

Let us note that there exists a lattice which is complete and strict.

REFERENCES

- $[1] \begin{tabular}{ll} Grzegorz Bancerek. Complete lattices. {\it Journal of Formalized Mathematics}, 4, 1992. http://mizar.org/JFM/Vol4/lattice3.html. {\it Journal of Formalized Mathematics}, 4, 1992. html. {\it Journal o$
- [2] Grzegorz Bancerek. Bounds in posets and relational substructures. Journal of Formalized Mathematics, 8, 1996. http://mizar.org/ JFM/Vol8/yellow_0.html.
- [3] Józef Białas. Group and field definitions. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/realsetl. html.
- [4] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/funct_1.html.
- [5] Czesław Byliński. Functions from a set to a set. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [6] Czesław Byliński. Some basic properties of sets. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/zfmisc_1.html.
- [7] Agata Darmochwał. Finite sets. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/finset_1.html.
- [8] G. Gierz, K.H. Hofmann, K. Keimel, J.D. Lawson, M. Mislove, and D.S. Scott. A Compendium of Continuous Lattices. Springer-Verlag, Berlin, Heidelberg, New York, 1980.
- [9] Adam Grabowski. On the category of posets. Journal of Formalized Mathematics, 8, 1996. http://mizar.org/JFM/Vol8/orders_3.html.
- [10] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/pre_topc.html.
- [11] Andrzej Trybulec. Binary operations applied to functions. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/funcop_1.html.
- [12] Andrzej Trybulec. Tarski Grothendieck set theory. Journal of Formalized Mathematics, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.
- [13] Wojciech A. Trybulec. Partially ordered sets. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/orders_ 1.html.
- [14] Zinaida Trybulec. Properties of subsets. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

- [15] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Voll/relat_1.html.
- [16] Edmund Woronowicz. Relations defined on sets. Journal of Formalized Mathematics, 1, 1989. http://mizar.org/JFM/Voll/relset_1.html.
- [17] Edmund Woronowicz and Anna Zalewska. Properties of binary relations. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_2.html.

Received September 12, 1996

Published January 2, 2004