

Introduction to Meet-Continuous Topological Lattices¹

Artur Korniłowicz
University of Białystok

MML Identifier: YELLOW13.

WWW: <http://mizar.org/JFM/Vol10/yellow13.html>

The articles [21], [8], [26], [27], [6], [7], [11], [24], [19], [28], [25], [10], [15], [14], [1], [20], [4], [22], [5], [2], [3], [13], [12], [9], [29], [16], [17], [23], and [18] provide the notation and terminology for this paper.

1. PRELIMINARIES

Let S be a finite 1-sorted structure. Note that the carrier of S is finite.

Let S be a trivial 1-sorted structure. Note that the carrier of S is trivial.

Let us mention that every set which is trivial is also finite.

Let us mention that every 1-sorted structure which is trivial is also finite.

Let us note that every 1-sorted structure which is non trivial is also non empty.

One can verify the following observations:

- * there exists a 1-sorted structure which is strict, non empty, and trivial,
- * there exists a relational structure which is strict, non empty, and trivial, and
- * there exists a FR-structure which is strict, non empty, and trivial.

We now state the proposition

- (1) For every T_1 non empty topological space T holds every finite subset of T is closed.

Let T be a T_1 non empty topological space. One can check that every subset of T which is finite is also closed.

Let T be a compact topological structure. Note that Ω_T is compact.

Let us note that there exists a topological space which is strict, non empty, and trivial.

Let us observe that every non empty topological space which is finite and T_1 is also discrete.

Let us mention that every topological space which is finite is also compact.

One can prove the following propositions:

- (2) Every discrete non empty topological space is a T_4 space.
- (3) Every discrete non empty topological space is a T_3 space.
- (4) Every discrete non empty topological space is a T_2 space.

¹This work has been supported by KBN Grant 8 T11C 018 12.

- (5) Every discrete non empty topological space is a T_1 space.

One can verify that every topological space which is discrete and non empty is also T_4 , T_3 , T_2 , and T_1 .

One can check that every non empty topological space which is T_4 and T_1 is also T_3 .

Let us note that every non empty topological space which is T_3 and T_1 is also T_2 .

Let us note that every topological space which is T_2 is also T_1 .

Let us observe that every topological space which is T_1 is also T_0 .

We now state three propositions:

- (6) Let S be a reflexive relational structure, T be a reflexive transitive relational structure, f be a map from S into T , and X be a subset of S . Then $\downarrow(f^\circ X) \subseteq \downarrow(f^\circ \downarrow X)$.
- (7) Let S be a reflexive relational structure, T be a reflexive transitive relational structure, f be a map from S into T , and X be a subset of S . If f is monotone, then $\downarrow(f^\circ X) = \downarrow(f^\circ \downarrow X)$.
- (8) For every non empty poset N holds $\text{IdsMap}(N)$ is one-to-one.

Let N be a non empty poset. Note that $\text{IdsMap}(N)$ is one-to-one.

The following proposition is true

- (9) For every finite lattice N holds $\text{SupMap}(N)$ is one-to-one.

Let N be a finite lattice. Observe that $\text{SupMap}(N)$ is one-to-one.

One can prove the following three propositions:

- (10) For every finite lattice N holds N and $\langle \text{Ids}(N), \subseteq \rangle$ are isomorphic.
- (11) Let N be a complete non empty poset, x be an element of N , and X be a non empty subset of N . Then $x \sqcap \square$ preserves inf of X .
- (12) For every complete non empty poset N and for every element x of N holds $x \sqcap \square$ is meet-preserving.

Let N be a complete non empty poset and let x be an element of N . Observe that $x \sqcap \square$ is meet-preserving.

2. ON THE BASIS OF TOPOLOGICAL SPACES

We now state several propositions:

- (13) Let T be an anti-discrete non empty topological structure and p be a point of T . Then $\{\text{the carrier of } T\}$ is a basis of p .
- (14) Let T be an anti-discrete non empty topological structure, p be a point of T , and D be a basis of p . Then $D = \{\text{the carrier of } T\}$.
- (15) Let T be a non empty topological space, P be a basis of T , and p be a point of T . Then $\{A; A \text{ ranges over subsets of } T: A \in P \wedge p \in A\}$ is a basis of p .
- (16) Let T be a non empty topological structure, A be a subset of T , and p be a point of T . Then $p \in \bar{A}$ if and only if for every basis K of p and for every subset Q of T such that $Q \in K$ holds A meets Q .
- (17) Let T be a non empty topological structure, A be a subset of T , and p be a point of T . Then $p \in \bar{A}$ if and only if there exists a basis K of p such that for every subset Q of T such that $Q \in K$ holds A meets Q .

Let T be a topological structure and let p be a point of T . A family of subsets of T is said to be a generalized basis of p if:

(Def. 1) For every subset A of T such that $p \in \text{Int}A$ there exists a subset P of T such that $P \in \text{it}$ and $p \in \text{Int}P$ and $P \subseteq A$.

Let T be a non empty topological space and let p be a point of T . Let us note that the generalized basis of p can be characterized by the following (equivalent) condition:

(Def. 2) For every neighbourhood A of p there exists a neighbourhood P of p such that $P \in \text{it}$ and $P \subseteq A$.

Next we state two propositions:

(18) Let T be a topological structure and p be a point of T . Then $2^{\text{the carrier of } T}$ is a generalized basis of p .

(19) For every non empty topological space T and for every point p of T holds every generalized basis of p is non empty.

Let T be a non empty topological space and let p be a point of T . One can verify that every generalized basis of p is non empty.

Let T be a topological structure and let p be a point of T . Note that there exists a generalized basis of p which is non empty.

Let T be a topological structure, let p be a point of T , and let P be a generalized basis of p . We say that P is correct if and only if:

(Def. 3) For every subset A of T holds $A \in P$ iff $p \in \text{Int}A$.

Let T be a topological structure and let p be a point of T . One can verify that there exists a generalized basis of p which is correct.

Next we state the proposition

(20) Let T be a topological structure and p be a point of T . Then $\{A; A \text{ ranges over subsets of } T: p \in \text{Int}A\}$ is a correct generalized basis of p .

Let T be a non empty topological space and let p be a point of T . Observe that there exists a generalized basis of p which is non empty and correct.

The following propositions are true:

(21) Let T be an anti-discrete non empty topological structure and p be a point of T . Then $\{\text{the carrier of } T\}$ is a correct generalized basis of p .

(22) Let T be an anti-discrete non empty topological structure, p be a point of T , and D be a correct generalized basis of p . Then $D = \{\text{the carrier of } T\}$.

(23) For every non empty topological space T and for every point p of T holds every basis of p is a generalized basis of p .

Let T be a topological structure. A family of subsets of T is said to be a generalized basis of T if:

(Def. 4) For every point p of T holds it is a generalized basis of p .

One can prove the following two propositions:

(24) For every topological structure T holds $2^{\text{the carrier of } T}$ is a generalized basis of T .

(25) For every non empty topological space T holds every generalized basis of T is non empty.

Let T be a non empty topological space. Observe that every generalized basis of T is non empty.

Let T be a topological structure. Observe that there exists a generalized basis of T which is non empty.

Next we state two propositions:

(26) For every non empty topological space T and for every generalized basis P of T holds the topology of $T \subseteq \text{UniCl}(\text{Int}P)$.

(27) For every topological space T holds every basis of T is a generalized basis of T .

Let T be a non empty topological space-like FR-structure. We say that T satisfies conditions of topological semilattice if and only if:

(Def. 5) For every map f from $[T, (T \text{ qua topological space})]$ into T such that $f = \sqcap_T$ holds f is continuous.

One can verify that every non empty topological space-like FR-structure which is reflexive and trivial satisfies also conditions of topological semilattice.

Let us mention that there exists a FR-structure which is reflexive, trivial, non empty, and topological space-like.

The following proposition is true

(28) Let T be a non empty topological space-like FR-structure satisfying conditions of topological semilattice and x be an element of T . Then $x \sqcap \square$ is continuous.

Let T be a non empty topological space-like FR-structure satisfying conditions of topological semilattice and let x be an element of T . Observe that $x \sqcap \square$ is continuous.

REFERENCES

- [1] Grzegorz Bancerek. Complete lattices. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/lattice3.html>.
- [2] Grzegorz Bancerek. Bounds in posets and relational substructures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_0.html.
- [3] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_0.html.
- [4] Józef Białas. Group and field definitions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/realset1.html>.
- [5] Józef Białas and Yatsuka Nakamura. Dyadic numbers and T_4 topological spaces. *Journal of Formalized Mathematics*, 7, 1995. <http://mizar.org/JFM/Vol7/urysohn1.html>.
- [6] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [7] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [8] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [9] Czesław Byliński. Galois connections. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_1.html.
- [10] Agata Darmochwał. Compact spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/compts_1.html.
- [11] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.
- [12] Adam Grabowski. On the category of posets. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/orders_3.html.
- [13] Adam Grabowski and Robert Milewski. Boolean posets, posets under inclusion and products of relational structures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_1.html.
- [14] Zbigniew Karno. The lattice of domains of an extremally disconnected space. *Journal of Formalized Mathematics*, 4, 1992. http://mizar.org/JFM/Vol4/tlat_3.html.
- [15] Zbigniew Karno and Toshihiko Watanabe. Completeness of the lattices of domains of a topological space. *Journal of Formalized Mathematics*, 4, 1992. http://mizar.org/JFM/Vol4/tlat_2.html.
- [16] Artur Kornilowicz. Cartesian products of relations and relational structures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_3.html.
- [17] Artur Kornilowicz. Meet – continuous lattices. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_2.html.
- [18] Artur Kornilowicz. On the topological properties of meet-continuous lattices. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_9.html.

- [19] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/pre_topc.html.
- [20] Alexander Yu. Shibakov and Andrzej Trybulec. The Cantor set. *Journal of Formalized Mathematics*, 7, 1995. http://mizar.org/JFM/Vol7/cantor_1.html.
- [21] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [22] Andrzej Trybulec. A Borsuk theorem on homotopy types. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/borsuk_1.html.
- [23] Andrzej Trybulec. Baire spaces, Sober spaces. *Journal of Formalized Mathematics*, 9, 1997. http://mizar.org/JFM/Vol9/yellow_8.html.
- [24] Wojciech A. Trybulec. Partially ordered sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/orders_1.html.
- [25] Wojciech A. Trybulec. Groups. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/group_1.html.
- [26] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [27] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.
- [28] Mirosław Wysocki and Agata Darmochwał. Subsets of topological spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/tops_1.html.
- [29] Mariusz Żynel and Czesław Byliński. Properties of relational structures, posets, lattices and maps. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_2.html.

Received November 3, 1998

Published January 2, 2004
