

Retracts and Inheritance¹

Grzegorz Bancerek
University of Białystok

MML Identifier: YELLOW16.

WWW: <http://mizar.org/JFM/Vol11/yellow16.html>

The articles [22], [12], [26], [27], [9], [11], [10], [8], [1], [18], [24], [19], [13], [25], [2], [20], [21], [15], [29], [23], [3], [4], [5], [16], [28], [14], [7], [6], and [17] provide the notation and terminology for this paper.

1. POSET RETRACTS

One can prove the following propositions:

- (1) For all binary relations a, b holds $a \cdot b = a b$.
- (2) Let X be a set, L be a non empty relational structure, S be a non empty relational substructure of L , f, g be functions from X into the carrier of S , and f', g' be functions from X into the carrier of L . If $f' = f$ and $g' = g$ and $f \leq g$, then $f' \leq g'$.
- (3) Let X be a set, L be a non empty relational structure, S be a full non empty relational substructure of L , f, g be functions from X into the carrier of S , and f', g' be functions from X into the carrier of L . If $f' = f$ and $g' = g$ and $f' \leq g'$, then $f \leq g$.

Let S be a non empty relational structure and let T be a non empty reflexive antisymmetric relational structure. One can verify that there exists a map from S into T which is directed-sups-preserving and monotone.

Next we state the proposition

- (4) For all functions f, g such that f is idempotent and $\text{rng } g \subseteq \text{rng } f$ and $\text{rng } g \subseteq \text{dom } f$ holds $f \cdot g = g$.

Let S be a 1-sorted structure. One can check that there exists a map from S into S which is idempotent.

The following four propositions are true:

- (5) For every up-complete non empty poset L holds every directed-sups-inheriting full non empty relational substructure of L is up-complete.
- (6) Let L be an up-complete non empty poset and f be a map from L into L . Suppose f is idempotent and directed-sups-preserving. Then $\text{Im } f$ is directed-sups-inheriting.
- (8)¹ Let S, T be non empty relational structures, f be a map from T into S , and g be a map from S into T . If $f \cdot g = \text{id}_S$, then $\text{rng } f = \text{the carrier of } S$.

¹This work has been supported by KBN Grant 8 T11C 018 12.

¹ The proposition (7) has been removed.

- (9) Let T be a non empty relational structure, S be a non empty relational substructure of T , and f be a map from T into S . If $f \cdot \text{incl}(S, T) = \text{id}_S$, then f is an idempotent map from T into T .

Let S, T be non empty posets and let f be a function. We say that f is a retraction of T into S if and only if the conditions (Def. 1) are satisfied.

- (Def. 1)(i) f is a directed-sups-preserving map from T into S ,
(ii) $f \upharpoonright \text{the carrier of } S = \text{id}_S$, and
(iii) S is a directed-sups-inheriting full relational substructure of T .

We say that f is a UPS retraction of T into S if and only if the conditions (Def. 2) are satisfied.

- (Def. 2)(i) f is a directed-sups-preserving map from T into S , and
(ii) there exists a directed-sups-preserving map g from S into T such that $f \cdot g = \text{id}_S$.

Let S, T be non empty posets. We say that S is a retract of T if and only if:

- (Def. 3) There exists a map f from T into S such that f is a retraction of T into S .

We say that S is a UPS retract of T if and only if:

- (Def. 4) There exists a map f from T into S such that f is a UPS retraction of T into S .

The following propositions are true:

- (10) For all non empty posets S, T and for every function f such that f is a retraction of T into S holds $f \cdot \text{incl}(S, T) = \text{id}_S$.
- (11) Let S be a non empty poset, T be an up-complete non empty poset, and f be a function. Suppose f is a retraction of T into S . Then f is a UPS retraction of T into S .
- (12) Let S, T be non empty posets and f be a function. If f is a retraction of T into S , then $\text{rng } f = \text{the carrier of } S$.
- (13) Let S, T be non empty posets and f be a function. If f is a UPS retraction of T into S , then $\text{rng } f = \text{the carrier of } S$.
- (14) Let S, T be non empty posets and f be a function. Suppose f is a retraction of T into S . Then f is an idempotent map from T into T .
- (15) Let T, S be non empty posets and f be a map from T into T . Suppose f is a retraction of T into S . Then $\text{Im } f = \text{the relational structure of } S$.
- (16) Let T be an up-complete non empty poset, S be a non empty poset, and f be a map from T into T . Suppose f is a retraction of T into S . Then f is directed-sups-preserving and projection.
- (17) Let S, T be non empty reflexive transitive relational structures and f be a map from S into T . Then f is isomorphic if and only if the following conditions are satisfied:
(i) f is monotone, and
(ii) there exists a monotone map g from T into S such that $f \cdot g = \text{id}_T$ and $g \cdot f = \text{id}_S$.
- (18) Let S, T be non empty posets. Then S and T are isomorphic if and only if there exists a monotone map f from S into T and there exists a monotone map g from T into S such that $f \cdot g = \text{id}_T$ and $g \cdot f = \text{id}_S$.
- (19) Let S, T be up-complete non empty posets. Suppose S and T are isomorphic. Then S is a UPS retract of T and T is a UPS retract of S .

- (20) Let T, S be non empty posets, f be a monotone map from T into S , and g be a monotone map from S into T . Suppose $f \cdot g = \text{id}_S$. Then there exists a projection map h from T into T such that $h = g \cdot f$ and $h|_{\text{Im } h} = \text{id}_{\text{Im } h}$ and S and $\text{Im } h$ are isomorphic.
- (21) Let T be an up-complete non empty poset, S be a non empty poset, and f be a function. Suppose f is a UPS retraction of T into S . Then there exists a directed-sups-preserving projection map h from T into T such that h is a retraction of T into $\text{Im } h$ and S and $\text{Im } h$ are isomorphic.
- (22) For every up-complete non empty poset L and for every non empty poset S such that S is a retract of L holds S is up-complete.
- (23) For every complete non empty poset L and for every non empty poset S such that S is a retract of L holds S is complete.
- (24) Let L be a continuous complete lattice and S be a non empty poset. If S is a retract of L , then S is continuous.
- (25) Let L be an up-complete non empty poset and S be a non empty poset. If S is a UPS retract of L , then S is up-complete.
- (26) Let L be a complete non empty poset and S be a non empty poset. If S is a UPS retract of L , then S is complete.
- (27) Let L be a continuous complete lattice and S be a non empty poset. If S is a UPS retract of L , then S is continuous.
- (28) Let L be a relational structure, S be a full relational substructure of L , and R be a relational substructure of S . Then R is full if and only if R is a full relational substructure of L .
- (29) Let L be a non empty transitive relational structure and S be a directed-sups-inheriting non empty full relational substructure of L . Then every directed-sups-inheriting non empty relational substructure of S is a directed-sups-inheriting relational substructure of L .
- (30) Let L be an up-complete non empty poset and S_1, S_2 be directed-sups-inheriting full non empty relational substructures of L . Suppose S_1 is a relational substructure of S_2 . Then S_1 is a directed-sups-inheriting full relational substructure of S_2 .

2. PRODUCTS

Let R be a binary relation. We say that R is poset-yielding if and only if:

(Def. 5) For every set x such that $x \in \text{rng } R$ holds x is a poset.

Let us mention that every binary relation which is poset-yielding is also relational structure yielding and reflexive-yielding.

Let X be a non empty set, let L be a non empty relational structure, let i be an element of X , and let Y be a subset of L^X . Then $\pi_i Y$ is a subset of L .

Let X be a set and let S be a poset. One can check that $X \mapsto S$ is poset-yielding.

Let I be a set. Observe that there exists a many sorted set indexed by I which is poset-yielding and nonempty.

Let I be a non empty set and let J be a poset-yielding nonempty many sorted set indexed by I . Note that $\prod J$ is transitive and antisymmetric.

Let I be a non empty set, let J be a poset-yielding nonempty many sorted set indexed by I , and let i be an element of I . Then $J(i)$ is a non empty poset.

Next we state a number of propositions:

- (31) Let I be a non empty set, J be a poset-yielding nonempty many sorted set indexed by I , f be an element of $\prod J$, and X be a subset of $\prod J$. Then $f \geq X$ if and only if for every element i of I holds $f(i) \geq \pi_i X$.

- (32) Let I be a non empty set, J be a poset-yielding nonempty many sorted set indexed by I , f be an element of $\prod J$, and X be a subset of $\prod J$. Then $f \leq X$ if and only if for every element i of I holds $f(i) \leq \pi_i X$.
- (33) Let I be a non empty set, J be a poset-yielding nonempty many sorted set indexed by I , and X be a subset of $\prod J$. Then $\sup X$ exists in $\prod J$ if and only if for every element i of I holds $\sup \pi_i X$ exists in $J(i)$.
- (34) Let I be a non empty set, J be a poset-yielding nonempty many sorted set indexed by I , and X be a subset of $\prod J$. Then $\inf X$ exists in $\prod J$ if and only if for every element i of I holds $\inf \pi_i X$ exists in $J(i)$.
- (35) Let I be a non empty set, J be a poset-yielding nonempty many sorted set indexed by I , and X be a subset of $\prod J$. If $\sup X$ exists in $\prod J$, then for every element i of I holds $(\sup X)(i) = \sup \pi_i X$.
- (36) Let I be a non empty set, J be a poset-yielding nonempty many sorted set indexed by I , and X be a subset of $\prod J$. If $\inf X$ exists in $\prod J$, then for every element i of I holds $(\inf X)(i) = \inf \pi_i X$.
- (37) Let I be a non empty set, J be a relational structure yielding nonempty reflexive-yielding many sorted set indexed by I , X be a directed subset of $\prod J$, and i be an element of I . Then $\pi_i X$ is directed.
- (38) Let I be a non empty set and J, K be relational structure yielding nonempty many sorted sets indexed by I . Suppose that for every element i of I holds $K(i)$ is a relational substructure of $J(i)$. Then $\prod K$ is a relational substructure of $\prod J$.
- (39) Let I be a non empty set and J, K be relational structure yielding nonempty many sorted sets indexed by I . Suppose that for every element i of I holds $K(i)$ is a full relational substructure of $J(i)$. Then $\prod K$ is a full relational substructure of $\prod J$.
- (40) Let L be a non empty relational structure, S be a non empty relational substructure of L , and X be a set. Then S^X is a relational substructure of L^X .
- (41) Let L be a non empty relational structure, S be a full non empty relational substructure of L , and X be a set. Then S^X is a full relational substructure of L^X .

3. INHERITANCE

Let S, T be non empty relational structures and let X be a set. We say that S inherits sup of X from T if and only if:

(Def. 6) If $\sup X$ exists in T , then $\sqcup_T X \in$ the carrier of S .

We say that S inherits inf of X from T if and only if:

(Def. 7) If $\inf X$ exists in T , then $\sqcap_T X \in$ the carrier of S .

The following propositions are true:

- (42) Let T be a non empty transitive relational structure, S be a full non empty relational substructure of T , and X be a subset of S . Then S inherits sup of X from T if and only if if $\sup X$ exists in T , then $\sup X$ exists in S and $\sup X = \sqcup_T X$.
- (43) Let T be a non empty transitive relational structure, S be a full non empty relational substructure of T , and X be a subset of S . Then S inherits inf of X from T if and only if if $\inf X$ exists in T , then $\inf X$ exists in S and $\inf X = \sqcap_T X$.

In this article we present several logical schemes. The scheme *ProductSupsInheriting* deals with a non empty set \mathcal{A} , poset-yielding nonempty many sorted sets \mathcal{B}, \mathcal{C} indexed by \mathcal{A} , and a binary predicate \mathcal{P} , and states that:

For every subset X of $\prod C$ such that $\mathcal{P}[X, \prod C]$ holds $\prod C$ inherits sup of X from $\prod \mathcal{B}$ provided the parameters satisfy the following conditions:

- For every subset X of $\prod C$ such that $\mathcal{P}[X, \prod C]$ and for every element i of \mathcal{A} holds $\mathcal{P}[\pi_i X, C(i)]$,
- For every element i of \mathcal{A} holds $C(i)$ is a full relational substructure of $\mathcal{B}(i)$, and
- For every element i of \mathcal{A} and for every subset X of $C(i)$ such that $\mathcal{P}[X, C(i)]$ holds $C(i)$ inherits sup of X from $\mathcal{B}(i)$.

The scheme *PowerSupsInheriting* deals with a non empty set \mathcal{A} , a non empty poset \mathcal{B} , a non empty full relational substructure C of \mathcal{B} , and a binary predicate \mathcal{P} , and states that:

For every subset X of $C^{\mathcal{A}}$ such that $\mathcal{P}[X, C^{\mathcal{A}}]$ holds $C^{\mathcal{A}}$ inherits sup of X from $\mathcal{B}^{\mathcal{A}}$ provided the parameters meet the following conditions:

- For every subset X of $C^{\mathcal{A}}$ such that $\mathcal{P}[X, C^{\mathcal{A}}]$ and for every element i of \mathcal{A} holds $\mathcal{P}[\pi_i X, C]$, and
- For every subset X of C such that $\mathcal{P}[X, C]$ holds C inherits sup of X from \mathcal{B} .

The scheme *ProductInfsInheriting* deals with a non empty set \mathcal{A} , poset-yielding nonempty many sorted sets \mathcal{B}, \mathcal{C} indexed by \mathcal{A} , and a binary predicate \mathcal{P} , and states that:

For every subset X of $\prod C$ such that $\mathcal{P}[X, \prod C]$ holds $\prod C$ inherits inf of X from $\prod \mathcal{B}$ provided the following conditions are met:

- For every subset X of $\prod C$ such that $\mathcal{P}[X, \prod C]$ and for every element i of \mathcal{A} holds $\mathcal{P}[\pi_i X, C(i)]$,
- For every element i of \mathcal{A} holds $C(i)$ is a full relational substructure of $\mathcal{B}(i)$, and
- For every element i of \mathcal{A} and for every subset X of $C(i)$ such that $\mathcal{P}[X, C(i)]$ holds $C(i)$ inherits inf of X from $\mathcal{B}(i)$.

The scheme *PowerInfsInheriting* deals with a non empty set \mathcal{A} , a non empty poset \mathcal{B} , a non empty full relational substructure C of \mathcal{B} , and a binary predicate \mathcal{P} , and states that:

For every subset X of $C^{\mathcal{A}}$ such that $\mathcal{P}[X, C^{\mathcal{A}}]$ holds $C^{\mathcal{A}}$ inherits inf of X from $\mathcal{B}^{\mathcal{A}}$ provided the parameters have the following properties:

- For every subset X of $C^{\mathcal{A}}$ such that $\mathcal{P}[X, C^{\mathcal{A}}]$ and for every element i of \mathcal{A} holds $\mathcal{P}[\pi_i X, C]$, and
- For every subset X of C such that $\mathcal{P}[X, C]$ holds C inherits inf of X from \mathcal{B} .

Let I be a set, let L be a non empty relational structure, let X be a non empty subset of L^I , and let i be a set. One can verify that $\pi_i X$ is non empty.

We now state the proposition

- (44) Let L be a non empty poset, S be a directed-sup-inheriting non empty full relational substructure of L , and X be a non empty set. Then S^X is a directed-sup-inheriting relational substructure of L^X .

Let I be a non empty set, let J be a relational structure yielding nonempty many sorted set indexed by I , let X be a non empty subset of $\prod J$, and let i be a set. One can check that $\pi_i X$ is non empty.

Next we state the proposition

- (45) For every non empty set X and for every up-complete non empty poset L holds L^X is up-complete.

Let L be an up-complete non empty poset and let X be a non empty set. Observe that L^X is up-complete.

4. TOPOLOGICAL RETRACTS

Let T be a topological space. One can check that the topology of T is non empty.

One can prove the following propositions:

- (46) Let T be a non empty topological space, S be a non empty subspace of T , and f be a map from T into S . If f is a retraction, then $\text{rng } f = \text{the carrier of } S$.
- (47) Let T be a non empty topological space, S be a non empty subspace of T , and f be a continuous map from T into S . If f is a retraction, then f is idempotent.
- (48) Let T be a non empty topological space and V be an open subset of T . Then $\chi_{V, \text{the carrier of } T}$ is a continuous map from T into the Sierpiński space.
- (49) Let T be a non empty topological space and x, y be elements of T . Suppose that for every open subset W of T such that $y \in W$ holds $x \in W$. Then $[0 \mapsto y, 1 \mapsto x]$ is a continuous map from the Sierpiński space into T .
- (50) Let T be a non empty topological space, x, y be elements of T , and V be an open subset of T . Suppose $x \in V$ and $y \notin V$. Then $\chi_{V, \text{the carrier of } T} \cdot [0 \mapsto y, 1 \mapsto x] = \text{id}_{\text{the Sierpiński space}}$.
- (51) Let T be a non empty 1-sorted structure, V, W be subsets of T , S be a topological augmentation of 2_{\subseteq}^1 , and f, g be maps from T into S . Suppose $f = \chi_{V, \text{the carrier of } T}$ and $g = \chi_{W, \text{the carrier of } T}$. Then $V \subseteq W$ if and only if $f \leq g$.
- (52) Let L be a non empty relational structure, X be a non empty set, and R be a full non empty relational substructure of L^X . Suppose that for every set a holds a is an element of R iff there exists an element x of L such that $a = X \mapsto x$. Then L and R are isomorphic.
- (53) Let S, T be non empty topological spaces. Then S and T are homeomorphic if and only if there exists a continuous map f from S into T and there exists a continuous map g from T into S such that $f \cdot g = \text{id}_T$ and $g \cdot f = \text{id}_S$.
- (54) Let T, S, R be non empty topological spaces, f be a map from T into S , g be a map from S into T , and h be a map from S into R . If $f \cdot g = \text{id}_S$ and h is a homeomorphism, then $h \cdot f \cdot (g \cdot h^{-1}) = \text{id}_R$.
- (55) Let T, S, R be non empty topological spaces. Suppose S is a topological retract of T and S and R are homeomorphic. Then R is a topological retract of T .
- (56) For every non empty topological space T and for every non empty subspace S of T holds $\text{incl}(S, T)$ is continuous.
- (57) Let T be a non empty topological space, S be a non empty subspace of T , and f be a continuous map from T into S . If f is a retraction, then $f \cdot \text{incl}(S, T) = \text{id}_S$.
- (58) Let T be a non empty topological space and S be a non empty subspace of T . If S is a retract of T , then S is a topological retract of T .
- (59) Let R, T be non empty topological spaces. Then R is a topological retract of T if and only if there exists a non empty subspace S of T such that S is a retract of T and S and R are homeomorphic.

REFERENCES

- [1] Grzegorz Bancerek. König's theorem. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/card_3.html.
- [2] Grzegorz Bancerek. Complete lattices. *Journal of Formalized Mathematics*, 4, 1992. <http://mizar.org/JFM/Vol4/lattice3.html>.
- [3] Grzegorz Bancerek. Quantales. *Journal of Formalized Mathematics*, 6, 1994. <http://mizar.org/JFM/Vol6/quantall.html>.
- [4] Grzegorz Bancerek. Bounds in posets and relational substructures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_0.html.
- [5] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_0.html.
- [6] Grzegorz Bancerek. The "way-below" relation. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_3.html.

- [7] Grzegorz Bancerek. Bases and refinements of topologies. *Journal of Formalized Mathematics*, 10, 1998. http://mizar.org/JFM/Vol10/yellow_9.html.
- [8] Czesław Byliński. Basic functions and operations on functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_3.html.
- [9] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.
- [10] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.
- [11] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. <http://mizar.org/JFM/Vol1/partfun1.html>.
- [12] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.
- [13] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/funct_4.html.
- [14] Czesław Byliński. Galois connections. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_1.html.
- [15] Agata Darmochwał. Families of subsets, subspaces and mappings in topological spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/tops_2.html.
- [16] Adam Grabowski and Robert Milewski. Boolean posets, posets under inclusion and products of relational structures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_1.html.
- [17] Jarosław Gryko. Injective spaces. *Journal of Formalized Mathematics*, 10, 1998. <http://mizar.org/JFM/Vol10/waybell18.html>.
- [18] Andrzej Kondracki. Mostowski's fundamental operations — part I. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/zf_fund1.html.
- [19] Beata Madras. Product of family of universal algebras. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/pralg_1.html.
- [20] Michał Muzalewski. Categories of groups. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/grcat_1.html.
- [21] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/pre_topc.html.
- [22] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. <http://mizar.org/JFM/Axiomatics/tarski.html>.
- [23] Andrzej Trybulec. A Borsuk theorem on homotopy types. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/borsuk_1.html.
- [24] Andrzej Trybulec. Many-sorted sets. *Journal of Formalized Mathematics*, 5, 1993. <http://mizar.org/JFM/Vol5/pboole.html>.
- [25] Wojciech A. Trybulec. Partially ordered sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/orders_1.html.
- [26] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.
- [27] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.
- [28] Mariusz Żynel and Czesław Byliński. Properties of relational structures, posets, lattices and maps. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_2.html.
- [29] Mariusz Żynel and Adam Guzowski. T_0 topological spaces. *Journal of Formalized Mathematics*, 6, 1994. http://mizar.org/JFM/Vol6/t_0topsp.html.

Received September 7, 1999

Published January 2, 2004
