JOURNAL OF FORMALIZED MATHEMATICS
Volume 6, Released 1994, Published 2003
Inst. of Computer Science, Univ. of Białystok

On the Group of Inner Automorphisms

Artur Korniłowicz
Institute of Mathematics
Warsaw University
Białystok

MML Identifier: AUTGROUP.

WWW: http://mizar.org/JFM/Vol6/autgroup.html

The articles [7], [5], [15], [16], [2], [4], [3], [1], [8], [9], [6], [11], [12], [10], [13], and [14] provide the notation and terminology for this paper.

For simplicity, we use the following convention: G denotes a strict group, H denotes a subgroup of G, a, b, x denote elements of G, and h denotes a homomorphism from G to G.

We now state the proposition

(1) For all a, b such that b is an element of H holds $b^a \in H$ iff H is normal.

Let us consider G. The functor $\mathrm{Aut}(G)$ yielding a non empty set of functions from the carrier of G to the carrier of G is defined as follows:

(Def. 1) Every element of $\mathrm{Aut}(G)$ is a homomorphism from G to G and for every h holds $h \in \mathrm{Aut}(G)$ iff h is one-to-one and an epimorphism.

We now state several propositions:

(3)[1] $\mathrm{Aut}(G) \subseteq (\text{the carrier of } G)^{\text{the carrier of } G}$.

(4) $\mathrm{id}_{\text{the carrier of } G}$ is an element of $\mathrm{Aut}(G)$.

(5) For every h holds $h \in \mathrm{Aut}(G)$ iff h is an isomorphism.

(6) For every element f of $\mathrm{Aut}(G)$ holds f^{-1} is a homomorphism from G to G.

(7) For every element f of $\mathrm{Aut}(G)$ holds f^{-1} is an element of $\mathrm{Aut}(G)$.

(8) For all elements f_1, f_2 of $\mathrm{Aut}(G)$ holds $f_1 \cdot f_2$ is an element of $\mathrm{Aut}(G)$.

Let us consider G. The functor $\mathrm{AutComp}(G)$ yielding a binary operation on $\mathrm{Aut}(G)$ is defined as follows:

(Def. 2) For all elements x, y of $\mathrm{Aut}(G)$ holds $(\mathrm{AutComp}(G))(x, y) = x \cdot y$.

Let us consider G. The functor $\mathrm{AutGroup}(G)$ yields a strict group and is defined as follows:

(Def. 3) $\mathrm{AutGroup}(G) = \langle \mathrm{Aut}(G), \mathrm{AutComp}(G) \rangle$.

The following three propositions are true:

[1] The proposition (2) has been removed.

© Association of Mizar Users

(9) For all elements x, y of $\mathrm{AutGroup}(G)$ and for all elements f, g of $\mathrm{Aut}(G)$ such that $x = f$ and $y = g$ holds $x \cdot y = f \cdot g$.

(10) $\mathrm{id}_{\text{the carrier of } G} = 1_{\mathrm{AutGroup}(G)}$.

(11) For every element f of $\mathrm{Aut}(G)$ and for every element g of $\mathrm{AutGroup}(G)$ such that $f = g$ holds $f^{-1} = g^{-1}$.

Let us consider G. The functor $\mathrm{InnAut}(G)$ yielding a non empty set of functions from the carrier of G to the carrier of G is defined by the condition (Def. 4).

(Def. 4) Let f be an element of (the carrier of G)$^{\text{the carrier of } G}$. Then $f \in \mathrm{InnAut}(G)$ if and only if there exists a such that for every x holds $f(x) = x^a$.

One can prove the following propositions:

(12) $\mathrm{InnAut}(G) \subseteq$ (the carrier of G)$^{\text{the carrier of } G}$.

(13) Every element of $\mathrm{InnAut}(G)$ is an element of $\mathrm{Aut}(G)$.

(14) $\mathrm{InnAut}(G) \subseteq \mathrm{Aut}(G)$.

(15) For all elements f, g of $\mathrm{InnAut}(G)$ holds $(\mathrm{AutComp}(G))(f, g) = f \cdot g$.

(16) $\mathrm{id}_{\text{the carrier of } G}$ is an element of $\mathrm{InnAut}(G)$.

(17) For every element f of $\mathrm{InnAut}(G)$ holds f^{-1} is an element of $\mathrm{InnAut}(G)$.

(18) For all elements f, g of $\mathrm{InnAut}(G)$ holds $f \cdot g$ is an element of $\mathrm{InnAut}(G)$.

Let us consider G. The functor $\mathrm{InnAutGroup}(G)$ yields a normal strict subgroup of $\mathrm{AutGroup}(G)$ and is defined as follows:

(Def. 5) The carrier of $\mathrm{InnAutGroup}(G) = \mathrm{InnAut}(G)$.

We now state three propositions:

(20)[2] For all elements x, y of $\mathrm{InnAutGroup}(G)$ and for all elements f, g of $\mathrm{InnAut}(G)$ such that $x = f$ and $y = g$ holds $x \cdot y = f \cdot g$.

(21) $\mathrm{id}_{\text{the carrier of } G} = 1_{\mathrm{InnAutGroup}(G)}$.

(22) For every element f of $\mathrm{InnAut}(G)$ and for every element g of $\mathrm{InnAutGroup}(G)$ such that $f = g$ holds $f^{-1} = g^{-1}$.

Let us consider G, b. The functor $\mathrm{Conjugate}(b)$ yielding an element of $\mathrm{InnAut}(G)$ is defined by:

(Def. 6) For every a holds $(\mathrm{Conjugate}(b))(a) = a^b$.

Next we state a number of propositions:

(23) For all a, b holds $\mathrm{Conjugate}(a \cdot b) = \mathrm{Conjugate}(b) \cdot \mathrm{Conjugate}(a)$.

(24) $\mathrm{Conjugate}(1_G) = \mathrm{id}_{\text{the carrier of } G}$.

(25) For every a holds $(\mathrm{Conjugate}(1_G))(a) = a$.

(26) For every a holds $\mathrm{Conjugate}(a) \cdot \mathrm{Conjugate}(a^{-1}) = \mathrm{Conjugate}(1_G)$.

(27) For every a holds $\mathrm{Conjugate}(a^{-1}) \cdot \mathrm{Conjugate}(a) = \mathrm{Conjugate}(1_G)$.

(28) For every a holds $\mathrm{Conjugate}(a^{-1}) = (\mathrm{Conjugate}(a))^{-1}$.

[2] The proposition (19) has been removed.

(29) For every a holds $\mathrm{Conjugate}(a) \cdot \mathrm{Conjugate}(1_G) = \mathrm{Conjugate}(a)$ and $\mathrm{Conjugate}(1_G) \cdot \mathrm{Conjugate}(a) = \mathrm{Conjugate}(a)$.

(30) For every element f of $\mathrm{InnAut}(G)$ holds $f \cdot \mathrm{Conjugate}(1_G) = f$ and $\mathrm{Conjugate}(1_G) \cdot f = f$.

(31) For every G holds $\mathrm{InnAutGroup}(G)$ and ${}^{G}/_{\mathrm{Z}(G)}$ are isomorphic.

(32) For every G such that G is a commutative group and for every element f of $\mathrm{InnAutGroup}(G)$ holds $f = 1_{\mathrm{InnAutGroup}(G)}$.

REFERENCES

[1] Czesław Byliński. Binary operations. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/binop_1.html.

[2] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.

[3] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.

[4] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/partfun1.html.

[5] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.

[6] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/vectsp_1.html.

[7] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.

[8] Andrzej Trybulec. Function domains and Frænkel operator. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/fraenkel.html.

[9] Wojciech A. Trybulec. Vectors in real linear space. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/rlvect_1.html.

[10] Wojciech A. Trybulec. Classes of conjugation. Normal subgroups. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/group_3.html.

[11] Wojciech A. Trybulec. Groups. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/group_1.html.

[12] Wojciech A. Trybulec. Subgroup and cosets of subgroups. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/group_2.html.

[13] Wojciech A. Trybulec. Commutator and center of a group. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/group_5.html.

[14] Wojciech A. Trybulec and Michał J. Trybulec. Homomorphisms and isomorphisms of groups. Quotient group. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/group_6.html.

[15] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

[16] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received April 22, 1994

Published January 2, 2004