JOURNAL OF FORMALIZED MATHEMATICS
Volume 4, Released 1992, Published 2003
Inst. of Computer Science, Univ. of Białystok

Series in Banach and Hilbert Spaces

Elżbieta Kraszewska
Warsaw University
Białystok

Jan Popiołek
Warsaw University
Białystok

Summary. In [14] the series of real numbers were investigated. The introduction to Banach and Hilbert spaces ([10], [11],[12]), enables us to arrive at the concept of series in Hilbert space. We start with the notions: partial sums of series, sum and n-th sum of series, convergent series (summable series), absolutely convergent series. We prove some basic theorems: the necessary condition for a series to converge, Weierstrass' test, d'Alembert's test, Cauchy's test.

MML Identifier: `BHSP_4`.

WWW: `http://mizar.org/JFM/Vol4/bhsp_4.html`

The articles [17], [2], [15], [4], [1], [3], [7], [5], [6], [14], [8], [16], [9], [10], [11], [12], and [13] provide the notation and terminology for this paper.

For simplicity, we adopt the following convention: X is a real unitary space, a, b, r are real numbers, s_1, s_2, s_3 are sequences of X, R_1, R_2, R_3 are sequences of real numbers, and k, n, m are natural numbers.

The scheme *Rec Func Ex RUS* deals with a real unitary space $\mathcal{A}$, a point $\mathcal{B}$ of $\mathcal{A}$, and a binary functor $\mathcal{F}$ yielding a point of $\mathcal{A}$, and states that:

> There exists a function f from $\mathbb{N}$ into the carrier of $\mathcal{A}$ such that $f(0) = \mathcal{B}$ and for every element n of $\mathbb{N}$ and for every point x of $\mathcal{A}$ such that $x = f(n)$ holds $f(n+1) = \mathcal{F}(n,x)$

for all values of the parameters.

Let us consider X and let us consider s_1. The functor $(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}}$ yields a sequence of X and is defined by:

(Def. 1) $(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}}(0) = s_1(0)$ and for every n holds $(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}}(n+1) = (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}}(n) + s_1(n+1)$.

The following propositions are true:

(1) $(\sum_{\alpha=0}^{\kappa}(s_2)(\alpha))_{\kappa\in\mathbb{N}} + (\sum_{\alpha=0}^{\kappa}(s_3)(\alpha))_{\kappa\in\mathbb{N}} = (\sum_{\alpha=0}^{\kappa}(s_2+s_3)(\alpha))_{\kappa\in\mathbb{N}}$.

(2) $(\sum_{\alpha=0}^{\kappa}(s_2)(\alpha))_{\kappa\in\mathbb{N}} - (\sum_{\alpha=0}^{\kappa}(s_3)(\alpha))_{\kappa\in\mathbb{N}} = (\sum_{\alpha=0}^{\kappa}(s_2-s_3)(\alpha))_{\kappa\in\mathbb{N}}$.

(3) $(\sum_{\alpha=0}^{\kappa}(a\cdot s_1)(\alpha))_{\kappa\in\mathbb{N}} = a\cdot(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}}$.

(4) $(\sum_{\alpha=0}^{\kappa}(-s_1)(\alpha))_{\kappa\in\mathbb{N}} = -(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}}$.

(5) $a\cdot(\sum_{\alpha=0}^{\kappa}(s_2)(\alpha))_{\kappa\in\mathbb{N}} + b\cdot(\sum_{\alpha=0}^{\kappa}(s_3)(\alpha))_{\kappa\in\mathbb{N}} = (\sum_{\alpha=0}^{\kappa}(a\cdot s_2+b\cdot s_3)(\alpha))_{\kappa\in\mathbb{N}}$.

Let us consider X and let us consider s_1. We say that s_1 is summable if and only if:

(Def. 2) $(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}}$ is convergent.

The functor $\sum s_1$ yielding a point of X is defined as follows:

(Def. 3) $\sum s_1 = \lim((\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}})$.

Next we state several propositions:

(6) If s_2 is summable and s_3 is summable, then s_2+s_3 is summable and $\sum(s_2+s_3) = \sum s_2 + \sum s_3$.

(7) If s_2 is summable and s_3 is summable, then s_2-s_3 is summable and $\sum(s_2-s_3) = \sum s_2 - \sum s_3$.

(8) If s_1 is summable, then $a\cdot s_1$ is summable and $\sum(a\cdot s_1) = a\cdot\sum s_1$.

(9) If s_1 is summable, then s_1 is convergent and $\lim s_1 = 0_X$.

(10) Suppose X is a Hilbert space. Then s_1 is summable if and only if for every r such that $r>0$ there exists k such that for all n, m such that $n\geq k$ and $m\geq k$ holds $\|(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}}(n) - (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}}(m)\| < r$.

(11) If s_1 is summable, then $(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}}$ is bounded.

(12) For all s_1, s_2 such that for every n holds $s_2(n) = s_1(0)$ holds $(\sum_{\alpha=0}^{\kappa}(s_1\uparrow 1)(\alpha))_{\kappa\in\mathbb{N}} = (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}}\uparrow 1 - s_2$.

(13) If s_1 is summable, then for every k holds $s_1\uparrow k$ is summable.

(14) If there exists k such that $s_1\uparrow k$ is summable, then s_1 is summable.

Let us consider X, s_1, n. The functor $\sum_{\kappa=0}^{n}s_1(\kappa)$ yielding a point of X is defined by:

(Def. 4) $\sum_{\kappa=0}^{n}s_1(\kappa) = (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}}(n)$.

We now state several propositions:

(16)[1] $\sum_{\kappa=0}^{0}s_1(\kappa) = s_1(0)$.

(17) $\sum_{\kappa=0}^{1}s_1(\kappa) = \sum_{\kappa=0}^{0}s_1(\kappa) + s_1(1)$.

(18) $\sum_{\kappa=0}^{1}s_1(\kappa) = s_1(0) + s_1(1)$.

(19) $\sum_{\kappa=0}^{n+1}s_1(\kappa) = \sum_{\kappa=0}^{n}s_1(\kappa) + s_1(n+1)$.

(20) $s_1(n+1) = \sum_{\kappa=0}^{n+1}s_1(\kappa) - \sum_{\kappa=0}^{n}s_1(\kappa)$.

(21) $s_1(1) = \sum_{\kappa=0}^{1}s_1(\kappa) - \sum_{\kappa=0}^{0}s_1(\kappa)$.

Let us consider X, s_1, n, m. The functor $\sum_{\kappa=n+1}^{m}s_1(\kappa)$ yields a point of X and is defined as follows:

(Def. 5) $\sum_{\kappa=n+1}^{m}s_1(\kappa) = \sum_{\kappa=0}^{n}s_1(\kappa) - \sum_{\kappa=0}^{m}s_1(\kappa)$.

One can prove the following propositions:

(23)[2] $\sum_{\kappa=1+1}^{0}s_1(\kappa) = s_1(1)$.

(24) $\sum_{\kappa=n+1+1}^{n}s_1(\kappa) = s_1(n+1)$.

(25) Suppose X is a Hilbert space. Then s_1 is summable if and only if for every r such that $r>0$ there exists k such that for all n, m such that $n\geq k$ and $m\geq k$ holds $\|\sum_{\kappa=0}^{n}s_1(\kappa) - \sum_{\kappa=0}^{m}s_1(\kappa)\| < r$.

[1] The proposition (15) has been removed.
[2] The proposition (22) has been removed.

(26) Suppose X is a Hilbert space. Then s_1 is summable if and only if for every r such that $r > 0$ there exists k such that for all n, m such that $n \geq k$ and $m \geq k$ holds $\|\sum_{\kappa=n+1}^{m} s_1(\kappa)\| < r$.

Let us consider R_1, n. The functor $\sum_{\kappa=0}^{n} R_1(\kappa)$ yielding a real number is defined by:

(Def. 6) $\sum_{\kappa=0}^{n} R_1(\kappa) = (\sum_{\alpha=0}^{\kappa} (R_1)(\alpha))_{\kappa\in\mathbb{N}}(n)$.

Let us consider R_1, n, m. The functor $\sum_{\kappa=n+1}^{m} R_1(\kappa)$ yielding a real number is defined as follows:

(Def. 7) $\sum_{\kappa=n+1}^{m} R_1(\kappa) = \sum_{\kappa=0}^{n} R_1(\kappa) - \sum_{\kappa=0}^{m} R_1(\kappa)$.

Let us consider X, s_1. We say that s_1 is absolutely summable if and only if:

(Def. 8) $\|s_1\|$ is summable.

Next we state a number of propositions:

(27) If s_2 is absolutely summable and s_3 is absolutely summable, then $s_2 + s_3$ is absolutely summable.

(28) If s_1 is absolutely summable, then $a \cdot s_1$ is absolutely summable.

(29) If for every n holds $\|s_1\|(n) \leq R_1(n)$ and R_1 is summable, then s_1 is absolutely summable.

(30) If for every n holds $s_1(n) \neq 0_X$ and $R_1(n) = \frac{\|s_1(n+1)\|}{\|s_1(n)\|}$ and R_1 is convergent and $\lim R_1 < 1$, then s_1 is absolutely summable.

(31) If $r > 0$ and there exists m such that for every n such that $n \geq m$ holds $\|s_1(n)\| \geq r$, then s_1 is not convergent or $\lim s_1 \neq 0_X$.

(32) If for every n holds $s_1(n) \neq 0_X$ and there exists m such that for every n such that $n \geq m$ holds $\frac{\|s_1(n+1)\|}{\|s_1(n)\|} \geq 1$, then s_1 is not summable.

(33) If for every n holds $s_1(n) \neq 0_X$ and for every n holds $R_1(n) = \frac{\|s_1(n+1)\|}{\|s_1(n)\|}$ and R_1 is convergent and $\lim R_1 > 1$, then s_1 is not summable.

(34) If for every n holds $R_1(n) = \sqrt[n]{\|s_1(n)\|}$ and R_1 is convergent and $\lim R_1 < 1$, then s_1 is absolutely summable.

(35) If for every n holds $R_1(n) = \sqrt[n]{\|s_1\|(n)}$ and there exists m such that for every n such that $n \geq m$ holds $R_1(n) \geq 1$, then s_1 is not summable.

(36) If for every n holds $R_1(n) = \sqrt[n]{\|s_1\|(n)}$ and R_1 is convergent and $\lim R_1 > 1$, then s_1 is not summable.

(37) $(\sum_{\alpha=0}^{\kappa} \|s_1\|(\alpha))_{\kappa\in\mathbb{N}}$ is non-decreasing.

(38) For every n holds $(\sum_{\alpha=0}^{\kappa} \|s_1\|(\alpha))_{\kappa\in\mathbb{N}}(n) \geq 0$.

(39) For every n holds $\|(\sum_{\alpha=0}^{\kappa} (s_1)(\alpha))_{\kappa\in\mathbb{N}}(n)\| \leq (\sum_{\alpha=0}^{\kappa} \|s_1\|(\alpha))_{\kappa\in\mathbb{N}}(n)$.

(40) For every n holds $\|\sum_{\kappa=0}^{n} s_1(\kappa)\| \leq \sum_{\kappa=0}^{n} \|s_1\|(\kappa)$.

(41) For all n, m holds $\|(\sum_{\alpha=0}^{\kappa} (s_1)(\alpha))_{\kappa\in\mathbb{N}}(m) - (\sum_{\alpha=0}^{\kappa} (s_1)(\alpha))_{\kappa\in\mathbb{N}}(n)\| \leq |(\sum_{\alpha=0}^{\kappa} \|s_1\|(\alpha))_{\kappa\in\mathbb{N}}(m) - (\sum_{\alpha=0}^{\kappa} \|s_1\|(\alpha))_{\kappa\in\mathbb{N}}(n)|$.

(42) For all n, m holds $\|\sum_{\kappa=0}^{m} s_1(\kappa) - \sum_{\kappa=0}^{n} s_1(\kappa)\| \leq |\sum_{\kappa=0}^{m} \|s_1\|(\kappa) - \sum_{\kappa=0}^{n} \|s_1\|(\kappa)|$.

(43) For all n, m holds $\|\sum_{\kappa=m+1}^{n} s_1(\kappa)\| \leq |\sum_{\kappa=m+1}^{n} \|s_1\|(\kappa)|$.

(44) If X is a Hilbert space, then if s_1 is absolutely summable, then s_1 is summable.

Let us consider X, s_1, R_1. The functor $R_1 \cdot s_1$ yielding a sequence of X is defined by:

(Def. 9) For every n holds $(R_1 \cdot s_1)(n) = R_1(n) \cdot s_1(n)$.

The following propositions are true:

(45) $R_1 \cdot (s_2 + s_3) = R_1 \cdot s_2 + R_1 \cdot s_3$.

(46) $(R_2 + R_3) \cdot s_1 = R_2 \cdot s_1 + R_3 \cdot s_1$.

(47) $(R_2\, R_3) \cdot s_1 = R_2 \cdot (R_3 \cdot s_1)$.

(48) $(a\, R_1) \cdot s_1 = a \cdot (R_1 \cdot s_1)$.

(49) $R_1 \cdot -s_1 = (-R_1) \cdot s_1$.

(50) If R_1 is convergent and s_1 is convergent, then $R_1 \cdot s_1$ is convergent.

(51) If R_1 is bounded and s_1 is bounded, then $R_1 \cdot s_1$ is bounded.

(52) If R_1 is convergent and s_1 is convergent, then $R_1 \cdot s_1$ is convergent and $\lim(R_1 \cdot s_1) = \lim R_1 \cdot \lim s_1$.

Let us consider R_1. We say that R_1 is Cauchy if and only if:

(Def. 10) For every r such that $r > 0$ there exists k such that for all n, m such that $n \geq k$ and $m \geq k$ holds $|R_1(n) - R_1(m)| < r$.

We introduce R_1 is a Cauchy sequence as a synonym of R_1 is Cauchy.

Next we state four propositions:

(53) If X is a Hilbert space, then if s_1 is a Cauchy sequence and R_1 is a Cauchy sequence, then $R_1 \cdot s_1$ is a Cauchy sequence.

(54) For every n holds $(\sum_{\alpha=0}^{\kappa}((R_1 - R_1 \uparrow 1) \cdot (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}})(\alpha))_{\kappa\in\mathbb{N}}(n) = (\sum_{\alpha=0}^{\kappa}(R_1 \cdot s_1)(\alpha))_{\kappa\in\mathbb{N}}(n+1) - (R_1 \cdot (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}})(n+1)$.

(55) For every n holds $(\sum_{\alpha=0}^{\kappa}(R_1 \cdot s_1)(\alpha))_{\kappa\in\mathbb{N}}(n+1) = (R_1 \cdot (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}})(n+1) - (\sum_{\alpha=0}^{\kappa}((R_1 \uparrow 1 - R_1) \cdot (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}})(\alpha))_{\kappa\in\mathbb{N}}(n)$.

(56) For every n holds $\sum_{\kappa=0}^{n+1}(R_1 \cdot s_1)(\kappa) = (R_1 \cdot (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}})(n+1) - \sum_{\kappa=0}^{n}((R_1 \uparrow 1 - R_1) \cdot (\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}})(\kappa)$.

REFERENCES

[1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/nat_1.html`.

[2] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/ordinal1.html`.

[3] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/funct_2.html`.

[4] Krzysztof Hryniewiecki. Basic properties of real numbers. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/real_1.html`.

[5] Jarosław Kotowicz. Convergent sequences and the limit of sequences. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/seq_2.html`.

[6] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/seqm_3.html`.

[7] Jarosław Kotowicz. Real sequences and basic operations on them. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/seq_1.html`.

[8] Jan Popiołek. Some properties of functions modul and signum. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/absvalue.html`.

[9] Jan Popiołek. Real normed space. *Journal of Formalized Mathematics*, 2, 1990. `http://mizar.org/JFM/Vol2/normsp_1.html`.

[10] Jan Popiołek. Introduction to Banach and Hilbert spaces — part I. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/bhsp_1.html.

[11] Jan Popiołek. Introduction to Banach and Hilbert spaces — part II. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/bhsp_2.html.

[12] Jan Popiołek. Introduction to Banach and Hilbert spaces — part III. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/bhsp_3.html.

[13] Konrad Raczkowski and Andrzej Nędzusiak. Real exponents and logarithms. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/power.html.

[14] Konrad Raczkowski and Andrzej Nędzusiak. Series. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/series_1.html.

[15] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. http://mizar.org/JFM/Addenda/numbers.html.

[16] Wojciech A. Trybulec. Vectors in real linear space. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/rlvect_1.html.

[17] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

Received April 1, 1992

Published January 2, 2004