JOURNAL OF FORMALIZED MATHEMATICS
Volume 15, Released 2003, Published 2003
Inst. of Computer Science, Univ. of Białystok

Bessel's Inequality

Hiroshi Yamazaki
Shinshu University
Nagano

Yasunari Shidama
Shinshu University
Nagano

Yatsuka Nakamura
Shinshu University
Nagano

Summary. In this article we defined the operation of a set and proved Bessel's inequality. In the first section, we defined the sum of all results of an operation, in which the results are given by taking each element of a set. In the second section, we defined Orthogonal Family and Orthonormal Family. In the last section, we proved some properties of operation of set and Bessel's inequality.

MML Identifier: `BHSP_5`.

WWW: `http://mizar.org/JFM/Vol15/bhsp_5.html`

The articles [11], [16], [12], [7], [5], [6], [17], [14], [9], [13], [3], [8], [1], [10], [4], [2], and [15] provide the notation and terminology for this paper.

1. Sum of the Result of Operation with Each Element of a Set

For simplicity, we adopt the following convention: X denotes a real unitary space, x, y, y_1, y_2 denote points of X, i, j denote natural numbers, D_1 denotes a non empty set, and p_1, p_2 denote finite sequences of elements of D_1.

One can prove the following proposition

(1) Suppose p_1 is one-to-one and p_2 is one-to-one and $\operatorname{rng} p_1 = \operatorname{rng} p_2$. Then $\operatorname{dom} p_1 = \operatorname{dom} p_2$ and there exists a permutation P of $\operatorname{dom} p_1$ such that $p_2 = p_1 \cdot P$ and $\operatorname{dom} P = \operatorname{dom} p_1$ and $\operatorname{rng} P = \operatorname{dom} p_1$.

Let D_1 be a non empty set and let f be a binary operation on D_1. Let us assume that f is commutative and associative and has a unity. Let Y be a finite subset of D_1. The functor $f \oplus Y$ yields an element of D_1 and is defined as follows:

(Def. 1) There exists a finite sequence p of elements of D_1 such that p is one-to-one and $\operatorname{rng} p = Y$ and $f \oplus Y = f \odot p$.

Let us consider X and let Y be a finite subset of X. The functor $\operatorname{SetopSum}(Y,X)$ is defined by:

(Def. 2) $\operatorname{SetopSum}(Y,X) = \begin{cases} (\text{the addition of } X) \oplus Y, & \text{if } Y \neq \emptyset, \\ 0_X, & \text{otherwise.} \end{cases}$

Let us consider X, x, let p be a finite sequence, and let us consider i. The functor $\operatorname{PO}(i,p,x)$ is defined as follows:

(Def. 3) $\operatorname{PO}(i,p,x) = (\text{the scalar product of } X)(\langle x, p(i) \rangle)$.

Let D_2, D_1 be non empty sets, let F be a function from D_1 into D_2, and let p be a finite sequence of elements of D_1. The functor $\operatorname{FuncSeq}(F,p)$ yields a finite sequence of elements of D_2 and is defined by:

(Def. 4) $\mathrm{FuncSeq}(F, p) = F \cdot p$.

Let D_2, D_1 be non empty sets and let f be a binary operation on D_2. Let us assume that f is commutative and associative and has a unity. Let Y be a finite subset of D_1 and let F be a function from D_1 into D_2. Let us assume that $Y \subseteq \mathrm{dom}\, F$. The functor $\mathrm{setopfunc}(Y, D_1, D_2, F, f)$ yielding an element of D_2 is defined as follows:

(Def. 5) There exists a finite sequence p of elements of D_1 such that p is one-to-one and $\mathrm{rng}\, p = Y$ and $\mathrm{setopfunc}(Y, D_1, D_2, F, f) = f \odot \mathrm{FuncSeq}(F, p)$.

Let us consider X, x and let Y be a finite subset of X. The functor $\mathrm{SetopPreProd}(x, Y, X)$ yielding a real number is defined by the condition (Def. 6).

(Def. 6) There exists a finite sequence p of elements of the carrier of X such that

(i) p is one-to-one,

(ii) $\mathrm{rng}\, p = Y$, and

(iii) there exists a finite sequence q of elements of $\mathbb{R}$ such that $\mathrm{dom}\, q = \mathrm{dom}\, p$ and for every i such that $i \in \mathrm{dom}\, q$ holds $q(i) = \mathrm{PO}(i, p, x)$ and $\mathrm{SetopPreProd}(x, Y, X) = +_{\mathbb{R}} \odot q$.

Let us consider X, x and let Y be a finite subset of X. The functor $\mathrm{SetopProd}(x, Y, X)$ yields a real number and is defined as follows:

(Def. 7) $\mathrm{SetopProd}(x, Y, X) = \begin{cases} \mathrm{SetopPreProd}(x, Y, X), & \text{if } Y \neq \emptyset, \\ 0, & \text{otherwise.} \end{cases}$

2. Orthogonal Family and Orthonormal Family

Let us consider X. A subset of X is called an orthogonal family of X if:

(Def. 8) For all x, y such that $x \in$ it and $y \in$ it and $x \neq y$ holds $(x|y) = 0$.

The following proposition is true

(2) $\emptyset$ is an orthogonal family of X.

Let us consider X. One can check that there exists an orthogonal family of X which is finite.

Let us consider X. A subset of X is called an orthonormal family of X if:

(Def. 9) It is an orthogonal family of X and for every x such that $x \in$ it holds $(x|x) = 1$.

One can prove the following proposition

(3) $\emptyset$ is an orthonormal family of X.

Let us consider X. Observe that there exists an orthonormal family of X which is finite.

Next we state the proposition

(4) $x = 0_X$ iff for every y holds $(x|y) = 0$.

3. Bessel's Inequality

The following propositions are true:

(5) $\|x + y\|^2 + \|x - y\|^2 = 2 \cdot \|x\|^2 + 2 \cdot \|y\|^2$.

(6) If x, y are orthogonal, then $\|x + y\|^2 = \|x\|^2 + \|y\|^2$.

(7) Let p be a finite sequence of elements of the carrier of X. Suppose $\mathrm{len}\, p \geq 1$ and for all i, j such that $i \in \mathrm{dom}\, p$ and $j \in \mathrm{dom}\, p$ and $i \neq j$ holds (the scalar product of X)$(\langle p(i), p(j) \rangle) = 0$. Let q be a finite sequence of elements of $\mathbb{R}$. Suppose $\mathrm{dom}\, p = \mathrm{dom}\, q$ and for every i such that $i \in \mathrm{dom}\, q$ holds $q(i) =$ (the scalar product of X)$(\langle p(i), p(i) \rangle)$. Then ((the addition of $X \odot p$)|(the addition of $X \odot p$)) $= +_{\mathbb{R}} \odot q$.

(8) Let p be a finite sequence of elements of the carrier of X. Suppose $\operatorname{len} p \geq 1$. Let q be a finite sequence of elements of $\mathbb{R}$. Suppose $\operatorname{dom} p = \operatorname{dom} q$ and for every i such that $i \in \operatorname{dom} q$ holds $q(i) =$ (the scalar product of X)$(\langle x, p(i) \rangle)$. Then $(x|$(the addition of $X \odot p)) = +_{\mathbb{R}} \odot q$.

(9) Let S be a finite non empty subset of X and F be a function from the carrier of X into the carrier of X. Suppose $S \subseteq \operatorname{dom} F$ and for all y_1, y_2 such that $y_1 \in S$ and $y_2 \in S$ and $y_1 \neq y_2$ holds (the scalar product of X)$(\langle F(y_1), F(y_2) \rangle) = 0$. Let H be a function from the carrier of X into $\mathbb{R}$. Suppose $S \subseteq \operatorname{dom} H$ and for every y such that $y \in S$ holds $H(y) =$ (the scalar product of X)$(\langle F(y), F(y) \rangle)$. Let p be a finite sequence of elements of the carrier of X. Suppose p is one-to-one and $\operatorname{rng} p = S$. Then (the scalar product of X)($\langle$the addition of $X \odot \operatorname{FuncSeq}(F, p)$, the addition of $X \odot \operatorname{FuncSeq}(F, p) \rangle) = +_{\mathbb{R}} \odot \operatorname{FuncSeq}(H, p)$.

(10) Let S be a finite non empty subset of X and F be a function from the carrier of X into the carrier of X. Suppose $S \subseteq \operatorname{dom} F$. Let H be a function from the carrier of X into $\mathbb{R}$. Suppose $S \subseteq \operatorname{dom} H$ and for every y such that $y \in S$ holds $H(y) =$ (the scalar product of X)$(\langle x, F(y) \rangle)$. Let p be a finite sequence of elements of the carrier of X. Suppose p is one-to-one and $\operatorname{rng} p = S$. Then (the scalar product of X)($\langle x$, the addition of $X \odot \operatorname{FuncSeq}(F, p) \rangle) = +_{\mathbb{R}} \odot \operatorname{FuncSeq}(H, p)$.

(11) Let given X. Suppose the addition of X is commutative and associative and the addition of X has a unity. Let given x and S be a finite orthonormal family of X. Suppose S is non empty. Let H be a function from the carrier of X into $\mathbb{R}$. Suppose $S \subseteq \operatorname{dom} H$ and for every y such that $y \in S$ holds $H(y) = (x|y)^{\mathbf{2}}$. Let F be a function from the carrier of X into the carrier of X. Suppose $S \subseteq \operatorname{dom} F$ and for every y such that $y \in S$ holds $F(y) = (x|y) \cdot y$. Then $(x|\operatorname{setopfunc}(S,$ the carrier of X, the carrier of X, $F,$ the addition of $X)) = \operatorname{setopfunc}(S,$ the carrier of $X, \mathbb{R}, H, +_{\mathbb{R}})$.

(12) Let given X. Suppose the addition of X is commutative and associative and the addition of X has a unity. Let given x and S be a finite orthonormal family of X. Suppose S is non empty. Let F be a function from the carrier of X into the carrier of X. Suppose $S \subseteq \operatorname{dom} F$ and for every y such that $y \in S$ holds $F(y) = (x|y) \cdot y$. Let H be a function from the carrier of X into $\mathbb{R}$. Suppose $S \subseteq \operatorname{dom} H$ and for every y such that $y \in S$ holds $H(y) = (x|y)^{\mathbf{2}}$. Then $(\operatorname{setopfunc}(S,$ the carrier of X, the carrier of X, $F,$ the addition of $X)|\operatorname{setopfunc}(S,$ the carrier of X, the carrier of X, $F,$ the addition of $X)) = \operatorname{setopfunc}(S,$ the carrier of $X, \mathbb{R}, H, +_{\mathbb{R}})$.

(13) Let given X. Suppose the addition of X is commutative and associative and the addition of X has a unity. Let given x and S be a finite orthonormal family of X. Suppose S is non empty. Let H be a function from the carrier of X into $\mathbb{R}$. Suppose $S \subseteq \operatorname{dom} H$ and for every y such that $y \in S$ holds $H(y) = (x|y)^{\mathbf{2}}$. Then $\operatorname{setopfunc}(S,$ the carrier of $X, \mathbb{R}, H, +_{\mathbb{R}}) \leq \|x\|^{\mathbf{2}}$.

(14) Let D_2, D_1 be non empty sets and f be a binary operation on D_2. Suppose f is commutative and associative and has a unity. Let Y_1, Y_2 be finite subsets of D_1. Suppose Y_1 misses Y_2. Let F be a function from D_1 into D_2. Suppose $Y_1 \subseteq \operatorname{dom} F$ and $Y_2 \subseteq \operatorname{dom} F$. Let Z be a finite subset of D_1. If $Z = Y_1 \cup Y_2$, then $\operatorname{setopfunc}(Z, D_1, D_2, F, f) = f(\operatorname{setopfunc}(Y_1, D_1, D_2, F, f)$, $\operatorname{setopfunc}(Y_2, D_1, D_2, F, f))$.

REFERENCES

[1] Grzegorz Bancerek. Cardinal numbers. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/card_1.html`.

[2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/finseq_1.html`.

[3] Czesław Byliński. Binary operations. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/binop_1.html`.

[4] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/funct_1.html`.

[5] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/funct_2.html`.

[6] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/finset_1.html`.

[7] Krzysztof Hryniewiecki. Basic properties of real numbers. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/real_1.html`.

[8] Eugeniusz Kusak, Wojciech Leończuk, and Michał Muzalewski. Abelian groups, fields and vector spaces. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/vectsp_1.html`.

[9] Jan Popiołek. Introduction to Banach and Hilbert spaces — part I. *Journal of Formalized Mathematics*, 3, 1991. `http://mizar.org/JFM/Vol3/bhsp_1.html`.

[10] Andrzej Trybulec. Semilattice operations on finite subsets. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/setwiseo.html`.

[11] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. `http://mizar.org/JFM/Axiomatics/tarski.html`.

[12] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. `http://mizar.org/JFM/Addenda/numbers.html`.

[13] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers operations: min, max, square, and square root. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/square_1.html`.

[14] Wojciech A. Trybulec. Vectors in real linear space. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/rlvect_1.html`.

[15] Wojciech A. Trybulec. Binary operations on finite sequences. *Journal of Formalized Mathematics*, 2, 1990. `http://mizar.org/JFM/Vol2/finsop_1.html`.

[16] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/subset_1.html`.

[17] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/relat_1.html`.

Received January 30, 2003

Published January 2, 2004