JOURNAL OF FORMALIZED MATHEMATICS
Volume 11, Released 1999, Published 2003
Inst. of Computer Science, Univ. of Białystok

Propositional Calculus for Boolean Valued Functions. Part III

Shunichi Kobayashi
Shinshu University
Nagano

Summary. In this paper, we have proved some elementary propositional calculus formulae for Boolean valued functions.

MML Identifier: BVFUNC_7.

WWW: http://mizar.org/JFM/Vol11/bvfunc_7.html

The articles [3], [5], [8], [7], [6], [1], [4], and [2] provide the notation and terminology for this paper.

In this paper Y denotes a non empty set.

The following propositions are true:

(1) For all elements a, b of $Boolean^Y$ holds $(a \Rightarrow b) \wedge (\neg a \Rightarrow b) = b$.

(2) For all elements a, b of $Boolean^Y$ holds $(a \Rightarrow b) \wedge (a \Rightarrow \neg b) = \neg a$.

(3) For all elements a, b, c of $Boolean^Y$ holds $a \Rightarrow b \vee c = (a \Rightarrow b) \vee (a \Rightarrow c)$.

(4) For all elements a, b, c of $Boolean^Y$ holds $a \Rightarrow b \wedge c = (a \Rightarrow b) \wedge (a \Rightarrow c)$.

(5) For all elements a, b, c of $Boolean^Y$ holds $a \vee b \Rightarrow c = (a \Rightarrow c) \wedge (b \Rightarrow c)$.

(6) For all elements a, b, c of $Boolean^Y$ holds $a \wedge b \Rightarrow c = (a \Rightarrow c) \vee (b \Rightarrow c)$.

(7) For all elements a, b, c of $Boolean^Y$ holds $a \wedge b \Rightarrow c = a \Rightarrow b \Rightarrow c$.

(8) For all elements a, b, c of $Boolean^Y$ holds $a \wedge b \Rightarrow c = a \Rightarrow \neg b \vee c$.

(9) For all elements a, b, c of $Boolean^Y$ holds $a \Rightarrow b \vee c = a \wedge \neg b \Rightarrow c$.

(10) For all elements a, b of $Boolean^Y$ holds $a \wedge (a \Rightarrow b) = a \wedge b$.

(11) For all elements a, b of $Boolean^Y$ holds $(a \Rightarrow b) \wedge \neg b = \neg a \wedge \neg b$.

(12) For all elements a, b, c of $Boolean^Y$ holds $(a \Rightarrow b) \wedge (b \Rightarrow c) = (a \Rightarrow b) \wedge (b \Rightarrow c) \wedge (a \Rightarrow c)$.

(13) For every element a of $Boolean^Y$ holds $true(Y) \Rightarrow a = a$.

(14) For every element a of $Boolean^Y$ holds $a \Rightarrow false(Y) = \neg a$.

(15) For every element a of $Boolean^Y$ holds $false(Y) \Rightarrow a = true(Y)$.

(16) For every element a of $Boolean^Y$ holds $a \Rightarrow true(Y) = true(Y)$.

© Association of Mizar Users

(17) For every element a of $Boolean^Y$ holds $a \Rightarrow \neg a = \neg a$.

(18) For all elements a, b, c of $Boolean^Y$ holds $a \Rightarrow b \Subset c \Rightarrow a \Rightarrow c \Rightarrow b$.

(19) For all elements a, b, c of $Boolean^Y$ holds $a \Leftrightarrow b \Subset a \Leftrightarrow c \Leftrightarrow b \Leftrightarrow c$.

(20) For all elements a, b, c of $Boolean^Y$ holds $a \Leftrightarrow b \Subset a \Rightarrow c \Leftrightarrow b \Rightarrow c$.

(21) For all elements a, b, c of $Boolean^Y$ holds $a \Leftrightarrow b \Subset c \Rightarrow a \Leftrightarrow c \Rightarrow b$.

(22) For all elements a, b, c of $Boolean^Y$ holds $a \Leftrightarrow b \Subset a \wedge c \Leftrightarrow b \wedge c$.

(23) For all elements a, b, c of $Boolean^Y$ holds $a \Leftrightarrow b \Subset a \vee c \Leftrightarrow b \vee c$.

(24) For all elements a, b of $Boolean^Y$ holds $a \Subset a \Leftrightarrow b \Leftrightarrow b \Leftrightarrow a \Leftrightarrow a$.

(25) For all elements a, b of $Boolean^Y$ holds $a \Subset a \Rightarrow b \Leftrightarrow b$.

(26) For all elements a, b of $Boolean^Y$ holds $a \Subset b \Rightarrow a \Leftrightarrow a$.

(27) For all elements a, b of $Boolean^Y$ holds $a \Subset a \wedge b \Leftrightarrow b \wedge a \Leftrightarrow a$.

REFERENCES

[1] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.

[2] Shunichi Kobayashi and Kui Jia. A theory of Boolean valued functions and partitions. *Journal of Formalized Mathematics*, 10, 1998. http://mizar.org/JFM/Vol10/bvfunc_1.html.

[3] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.

[4] Andrzej Trybulec. Function domains and Frænkel operator. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/fraenkel.html.

[5] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

[6] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

[7] Edmund Woronowicz. Interpretation and satisfiability in the first order logic. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/valuat_1.html.

[8] Edmund Woronowicz. Many-argument relations. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/margrel1.html.

Received April 23, 1999

Published January 2, 2004