JOURNAL OF FORMALIZED MATHEMATICS
Volume 14, Released 2002, Published 2003
Inst. of Computer Science, Univ. of Białystok

Fibonacci Numbers

Robert M. Solovay
P. O. Box 5949
Eugene OR 97405
U. S. A.

Summary. We show that Fibonacci commutes with g.c.d.; we then derive the formula connecting the Fibonacci sequence with the roots of the polynomial $x^2 - x - 1$.

MML Identifier: FIB_NUM.

WWW: http://mizar.org/JFM/Vol14/fib_num.html

The articles [2], [9], [10], [5], [1], [3], [4], [7], [6], and [8] provide the notation and terminology for this paper.

1. Fibonacci Commutes with gcd

The following propositions are true:

(1) For all natural numbers m, n holds $\gcd(m,n) = \gcd(m,n+m)$.

(2) For all natural numbers k, m, n such that $\gcd(k,m) = 1$ holds $\gcd(k,m \cdot n) = \gcd(k,n)$.

(3) For every real number s such that $s > 0$ there exists a natural number n such that $n > 0$ and $0 < \frac{1}{n}$ and $\frac{1}{n} \leq s$.

In this article we present several logical schemes. The scheme *Fib Ind* concerns a unary predicate $\mathcal{P}$, and states that:

For every natural number k holds $\mathcal{P}[k]$

provided the following conditions are met:

- $\mathcal{P}[0]$,
- $\mathcal{P}[1]$, and
- For every natural number k such that $\mathcal{P}[k]$ and $\mathcal{P}[k+1]$ holds $\mathcal{P}[k+2]$.

The scheme *Bin Ind* concerns a binary predicate $\mathcal{P}$, and states that:

For all natural numbers m, n holds $\mathcal{P}[m,n]$

provided the following conditions are satisfied:

- For all natural numbers m, n such that $\mathcal{P}[m,n]$ holds $\mathcal{P}[n,m]$, and
- Let k be a natural number. Suppose that for all natural numbers m, n such that $m < k$ and $n < k$ holds $\mathcal{P}[m,n]$. Let m be a natural number. If $m \leq k$, then $\mathcal{P}[k,m]$.

We now state two propositions:

(4) For all natural numbers m, n holds $\mathrm{Fib}(m+(n+1)) = \mathrm{Fib}(n) \cdot \mathrm{Fib}(m) + \mathrm{Fib}(n+1) \cdot \mathrm{Fib}(m+1)$.

(5) For all natural numbers m, n holds $\gcd(\mathrm{Fib}(m), \mathrm{Fib}(n)) = \mathrm{Fib}(\gcd(m,n))$.

© Association of Mizar Users

2. Fibonacci Numbers and the Golden Mean

We now state the proposition

(6) Let x, a, b, c be real numbers. Suppose $a \neq 0$ and $\Delta(a,b,c) \geq 0$. Then $a \cdot x^2 + b \cdot x + c = 0$ if and only if $x = \frac{-b-\sqrt{\Delta(a,b,c)}}{2 \cdot a}$ or $x = \frac{-b+\sqrt{\Delta(a,b,c)}}{2 \cdot a}$.

The real number τ is defined by:

(Def. 1) $\tau = \frac{1+\sqrt{5}}{2}$.

The real number $\overline{\tau}$ is defined as follows:

(Def. 2) $\overline{\tau} = \frac{1-\sqrt{5}}{2}$.

Next we state several propositions:

(7) For every natural number n holds $\mathrm{Fib}(n) = \frac{\tau^n - \overline{\tau}^n}{\sqrt{5}}$.

(8) For every natural number n holds $|\mathrm{Fib}(n) - \frac{\tau^n}{\sqrt{5}}| < 1$.

(9) For all sequences F, G of real numbers such that for every natural number n holds $F(n) = G(n)$ holds $F = G$.

(10) For all sequences f, g, h of real numbers such that g is non-zero holds $(f/g)\,(g/h) = f/h$.

(11) For all sequences f, g of real numbers and for every natural number n holds $(f/g)(n) = \frac{f(n)}{g(n)}$ and $(f/g)(n) = f(n) \cdot g(n)^{-1}$.

(12) Let F be a sequence of real numbers. Suppose that for every natural number n holds $F(n) = \frac{\mathrm{Fib}(n+1)}{\mathrm{Fib}(n)}$. Then F is convergent and $\lim F = \tau$.

Acknowledgments

My thanks to Freek Wiedijk for helping me learn Mizar and to Piotr Rudnicki for instructive comments on an earlier version of this article. This article was finished while I was visiting Bialystok and Adam Naumowicz and Josef Urban helped me through some difficult moments.

References

[1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/nat_1.html.

[2] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/ordinal1.html.

[3] Grzegorz Bancerek and Piotr Rudnicki. Two programs for **scm**. Part I - preliminaries. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/pre_ff.html.

[4] Jarosław Kotowicz. Convergent sequences and the limit of sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/seq_2.html.

[5] Jarosław Kotowicz. Real sequences and basic operations on them. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/seq_1.html.

[6] Jan Popiołek. Some properties of functions modul and signum. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/absvalue.html.

[7] Jan Popiołek. Quadratic inequalities. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/quin_1.html.

[8] Konrad Raczkowski and Andrzej Nędzusiak. Real exponents and logarithms. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/power.html.

[9] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. http://mizar.org/JFM/Addenda/numbers.html.

[10] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers operations: min, max, square, and square root. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/square_1.html.

Received April 19, 2002

Published January 2, 2004