JOURNAL OF FORMALIZED MATHEMATICS
Volume 4, Released 1992, Published 2003
Inst. of Computer Science, Univ. of Białystok

Go-Board Theorem

Jarosław Kotowicz
Warsaw University
Białystok

Yatsuka Nakamura
Shinshu University
Nagano

Summary. We prove the Go-board theorem which is a special case of Hex Theorem. The article is based on [11].

MML Identifier: GOBOARD4.

WWW: http://mizar.org/JFM/Vol4/goboard4.html

The articles [12], [4], [15], [13], [1], [3], [2], [10], [14], [7], [5], [6], [9], and [8] provide the notation and terminology for this paper.

For simplicity, we use the following convention: p, p_1, p_2, q_1, q_2 denote points of $\mathcal{E}^2_{\rm T}$, P_1, P_2 denote subsets of $\mathcal{E}^2_{\rm T}$, f, f_1, f_2 denote finite sequences of elements of $\mathcal{E}^2_{\rm T}$, n denotes a natural number, and G denotes a Go-board.

One can prove the following propositions:

(1) Let given G, f_1, f_2. Suppose that $1 \leq \operatorname{len} f_1$ and $1 \leq \operatorname{len} f_2$ and f_1 is a sequence which elements belong to G and f_2 is a sequence which elements belong to G and $(f_1)_1 \in \operatorname{rng} \operatorname{Line}(G, 1)$ and $(f_1)_{\operatorname{len} f_1} \in \operatorname{rng} \operatorname{Line}(G, \operatorname{len} G)$ and $(f_2)_1 \in \operatorname{rng}(G_{\square,1})$ and $(f_2)_{\operatorname{len} f_2} \in \operatorname{rng}(G_{\square,\operatorname{width} G})$. Then $\operatorname{rng} f_1$ meets $\operatorname{rng} f_2$.

(2) Let given G, f_1, f_2. Suppose that $2 \leq \operatorname{len} f_1$ and $2 \leq \operatorname{len} f_2$ and f_1 is a sequence which elements belong to G and f_2 is a sequence which elements belong to G and $(f_1)_1 \in \operatorname{rng} \operatorname{Line}(G, 1)$ and $(f_1)_{\operatorname{len} f_1} \in \operatorname{rng} \operatorname{Line}(G, \operatorname{len} G)$ and $(f_2)_1 \in \operatorname{rng}(G_{\square,1})$ and $(f_2)_{\operatorname{len} f_2} \in \operatorname{rng}(G_{\square,\operatorname{width} G})$. Then $\widetilde{\mathcal{L}}(f_1)$ meets $\widetilde{\mathcal{L}}(f_2)$.

(3) Let given G, f_1, f_2. Suppose that $2 \leq \operatorname{len} f_1$ and $2 \leq \operatorname{len} f_2$ and f_1 is a sequence which elements belong to G and f_2 is a sequence which elements belong to G and $(f_1)_1 \in \operatorname{rng} \operatorname{Line}(G, 1)$ and $(f_1)_{\operatorname{len} f_1} \in \operatorname{rng} \operatorname{Line}(G, \operatorname{len} G)$ and $(f_2)_1 \in \operatorname{rng}(G_{\square,1})$ and $(f_2)_{\operatorname{len} f_2} \in \operatorname{rng}(G_{\square,\operatorname{width} G})$. Then $\widetilde{\mathcal{L}}(f_1)$ meets $\widetilde{\mathcal{L}}(f_2)$.

Let f be a finite sequence of elements of $\mathbb{R}$ and let r, s be real numbers. We say that f lies between r and s if and only if:

(Def. 1) For every n such that $n \in \operatorname{dom} f$ holds $r \leq f(n)$ and $f(n) \leq s$.

Let D be a non empty set, let f_1 be a finite sequence of elements of D, and let f_2 be a non empty finite sequence of elements of D. Note that $f_1 \frown f_2$ is non empty and $f_2 \frown f_1$ is non empty.

Next we state three propositions:

(4) Let f_1, f_2 be finite sequences of elements of $\mathcal{E}^2_{\rm T}$. Suppose that $2 \leq \operatorname{len} f_1$ and $2 \leq \operatorname{len} f_2$ and f_1 is special and f_2 is special and for every n such that $n \in \operatorname{dom} f_1$ and $n+1 \in \operatorname{dom} f_1$ holds $(f_1)_n \neq (f_1)_{n+1}$ and for every n such that $n \in \operatorname{dom} f_2$ and $n+1 \in \operatorname{dom} f_2$ holds $(f_2)_n \neq (f_2)_{n+1}$ and $\mathbf{X}$-coordinate(f_1) lies between $(\mathbf{X}\text{-coordinate}(f_1))(1)$ and $(\mathbf{X}\text{-coordinate}(f_1))(\operatorname{len} f_1)$

and **X**-coordinate(f_2) lies between (**X**-coordinate(f_1))(1) and (**X**-coordinate(f_1))(len f_1) and **Y**-coordinate(f_1) lies between (**Y**-coordinate(f_2))(1) and (**Y**-coordinate(f_2))(len f_2) and **Y**-coordinate(f_2) lies between (**Y**-coordinate(f_2))(1) and (**Y**-coordinate(f_2))(len f_2). Then $\widetilde{\mathcal{L}}(f_1)$ meets $\widetilde{\mathcal{L}}(f_2)$.

(5) Let f_1, f_2 be finite sequences of elements of $\mathcal{E}^2_{\mathrm{T}}$. Suppose that f_1 is one-to-one and special and $2 \leq \mathrm{len}\, f_1$ and f_2 is one-to-one and special and $2 \leq \mathrm{len}\, f_2$ and **X**-coordinate(f_1) lies between (**X**-coordinate(f_1))(1) and (**X**-coordinate(f_1))(len f_1) and **X**-coordinate(f_2) lies between (**X**-coordinate(f_1))(1) and (**X**-coordinate(f_1))(len f_1) and **Y**-coordinate(f_1) lies between (**Y**-coordinate(f_2))(1) and (**Y**-coordinate(f_2))(len f_2) and **Y**-coordinate(f_2) lies between (**Y**-coordinate(f_2))(1) and (**Y**-coordinate(f_2))(len f_2). Then $\widetilde{\mathcal{L}}(f_1)$ meets $\widetilde{\mathcal{L}}(f_2)$.

(8)[1] Let given P_1, P_2, p_1, p_2, q_1, q_2. Suppose that

(i) P_1 is a special polygonal arc joining p_1 and q_1,

(ii) P_2 is a special polygonal arc joining p_2 and q_2,

(iii) for every p such that $p \in P_1 \cup P_2$ holds $(p_1)_{\mathbf{1}} \leq p_{\mathbf{1}}$ and $p_{\mathbf{1}} \leq (q_1)_{\mathbf{1}}$, and

(iv) for every p such that $p \in P_1 \cup P_2$ holds $(p_2)_{\mathbf{2}} \leq p_{\mathbf{2}}$ and $p_{\mathbf{2}} \leq (q_2)_{\mathbf{2}}$.

Then P_1 meets P_2.

REFERENCES

[1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/nat_1.html.

[2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.

[3] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.

[4] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.

[5] Agata Darmochwał. The Euclidean space. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/euclid.html.

[6] Agata Darmochwał and Yatsuka Nakamura. The topological space $\mathcal{E}^2_{\mathrm{T}}$. Arcs, line segments and special polygonal arcs. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/topreal1.html.

[7] Katarzyna Jankowska. Matrices. Abelian group of matrices. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/matrix_1.html.

[8] Jarosław Kotowicz and Yatsuka Nakamura. Introduction to Go-Board — part I. *Journal of Formalized Mathematics*, 4, 1992. http://mizar.org/JFM/Vol4/goboard1.html.

[9] Yatsuka Nakamura and Jarosław Kotowicz. Connectedness conditions using polygonal arcs. *Journal of Formalized Mathematics*, 4, 1992. http://mizar.org/JFM/Vol4/topreal4.html.

[10] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/pre_topc.html.

[11] Yukio Takeuchi and Yatsuka Nakamura. On the Jordan curve theorem. Technical Report 19804, Dept. of Information Eng., Shinshu University, 500 Wakasato, Nagano city, Japan, April 1980.

[12] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.

[13] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. http://mizar.org/JFM/Addenda/numbers.html.

[14] Wojciech A. Trybulec. Pigeon hole principle. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_4.html.

[1] The propositions (6) and (7) have been removed.

[15] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

Received August 24, 1992

Published January 2, 2004