JOURNAL OF FORMALIZED MATHEMATICS
Volume 14, Released 2002, Published 2003
Inst. of Computer Science, Univ. of Białystok

Preparing the Internal Approximations of Simple Closed Curves[1]

Andrzej Trybulec
University of Białystok

Summary. We mean by an internal approximation of a simple closed curve a special polygon disjoint with it but sufficiently close to it, i.e. such that it is clock-wise oriented and its right cells meet the curve. We prove lemmas used in the next article to construct a sequence of internal approximations.

MML Identifier: JORDAN11.

WWW: http://mizar.org/JFM/Vol14/jordan11.html

The articles [18], [6], [21], [2], [20], [12], [1], [16], [3], [22], [5], [4], [13], [17], [8], [9], [10], [11], [14], [15], [7], and [19] provide the notation and terminology for this paper.

In this paper j, k, n denote natural numbers and C denotes a subset of $\mathcal{E}^2_{\mathrm{T}}$ satisfying conditions of simple closed curve.

Let us consider C. The functor ApproxIndex$C$ yields a natural number and is defined as follows:

(Def. 1) ApproxIndex$C$ is sufficiently large for C and for every j such that j is sufficiently large for C holds $j \geq$ ApproxIndex$C$.

The following proposition is true

(1) ApproxIndex$C \geq 1$.

Let us consider C. The functor Y-InitStart$C$ yields a natural number and is defined by:

(Def. 2) Y-InitStart$C <$ width Gauge$(C,$ ApproxIndex$C)$ and cell$($Gauge$(C,$ ApproxIndex$C),$ X-SpanStart$(C,$ ApproxIndex$C) -' 1,$ Y-InitStart$C) \subseteq$ BDDC and for every j such that $j <$ width Gauge$(C,$ ApproxIndex$C)$ and cell$($Gauge$(C,$ ApproxIndex$C),$ X-SpanStart$(C,$ ApproxIndex$C) -' 1, j) \subseteq$ BDDC holds $j \geq$ Y-InitStart$C$.

One can prove the following propositions:

(2) Y-InitStart$C > 1$.

(3) Y-InitStart$C + 1 <$ width Gauge$(C,$ ApproxIndex$C)$.

Let us consider C, n. Let us assume that n is sufficiently large for C. The functor Y-SpanStart(C,n) yields a natural number and is defined by the conditions (Def. 3).

(Def. 3)(i) Y-SpanStart$(C,n) \leq$ width Gauge(C,n),

(ii) for every k such that Y-SpanStart$(C,n) \leq k$ and $k \leq 2^{n-'\mathrm{ApproxIndex}\,C} \cdot ($Y-InitStart$C -' 2) +$ 2 holds cell$($Gauge$(C,n),$ X-SpanStart$(C,n) -' 1, k) \subseteq$ BDDC, and

[1] This work has been partially supported by CALCULEMUS grant HPRN-CT-2000-00102.

© Association of Mizar Users

(iii) for every j such that $j \leq \operatorname{width}\operatorname{Gauge}(C,n)$ and for every k such that $j \leq k$ and $k \leq 2^{n-'\operatorname{ApproxIndex}C} \cdot (\text{Y-InitStart}\,C -' 2) + 2$ holds $\operatorname{cell}(\operatorname{Gauge}(C,n), \text{X-SpanStart}(C,n) -' 1, k) \subseteq \operatorname{BDD}C$ holds $j \geq \text{Y-SpanStart}(C,n)$.

The following propositions are true:

(4) If n is sufficiently large for C, then $\text{X-SpanStart}(C,n) = 2^{n-'\operatorname{ApproxIndex}C} \cdot (\text{X-SpanStart}(C, \operatorname{ApproxIndex}C) - 2) + 2$.

(5) If n is sufficiently large for C, then $\text{Y-SpanStart}(C,n) \leq 2^{n-'\operatorname{ApproxIndex}C} \cdot (\text{Y-InitStart}\,C -' 2) + 2$.

(6) If n is sufficiently large for C, then $\operatorname{cell}(\operatorname{Gauge}(C,n), \text{X-SpanStart}(C,n) -' 1, \text{Y-SpanStart}(C,n)) \subseteq \operatorname{BDD}C$.

(7) If n is sufficiently large for C, then $1 < \text{Y-SpanStart}(C,n)$ and $\text{Y-SpanStart}(C,n) \leq \operatorname{width}\operatorname{Gauge}(C,n)$.

(8) If n is sufficiently large for C, then $\langle \text{X-SpanStart}(C,n), \text{Y-SpanStart}(C,n) \rangle \in$ the indices of $\operatorname{Gauge}(C,n)$.

(9) If n is sufficiently large for C, then $\langle \text{X-SpanStart}(C,n) -' 1, \text{Y-SpanStart}(C,n) \rangle \in$ the indices of $\operatorname{Gauge}(C,n)$.

(10) If n is sufficiently large for C, then $\operatorname{cell}(\operatorname{Gauge}(C,n), \text{X-SpanStart}(C,n) -' 1, \text{Y-SpanStart}(C,n) -' 1)$ meets C.

(11) If n is sufficiently large for C, then $\operatorname{cell}(\operatorname{Gauge}(C,n), \text{X-SpanStart}(C,n) -' 1, \text{Y-SpanStart}(C,n))$ misses C.

REFERENCES

[1] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/nat_1.html.

[2] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/ordinal1.html.

[3] Grzegorz Bancerek. Countable sets and Hessenberg's theorem. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/card_4.html.

[4] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.

[5] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.

[6] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.

[7] Czesław Byliński. Gauges. *Journal of Formalized Mathematics*, 11, 1999. http://mizar.org/JFM/Vol11/jordan8.html.

[8] Agata Darmochwał. Compact spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/compts_1.html.

[9] Agata Darmochwał. The Euclidean space. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/euclid.html.

[10] Agata Darmochwał and Yatsuka Nakamura. The topological space $\mathcal{E}^2_{\mathrm{T}}$. Arcs, line segments and special polygonal arcs. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/topreal1.html.

[11] Agata Darmochwał and Yatsuka Nakamura. The topological space $\mathcal{E}^2_{\mathrm{T}}$. Simple closed curves. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/topreal2.html.

[12] Krzysztof Hryniewiecki. Basic properties of real numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/real_1.html.

[13] Katarzyna Jankowska. Matrices. Abelian group of matrices. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/matrix_1.html.

[14] Yatsuka Nakamura and Andrzej Trybulec. Decomposing a Go-Board into cells. *Journal of Formalized Mathematics*, 7, 1995. http://mizar.org/JFM/Vol7/goboard5.html.

[15] Yatsuka Nakamura, Andrzej Trybulec, and Czesław Byliński. Bounded domains and unbounded domains. *Journal of Formalized Mathematics*, 11, 1999. http://mizar.org/JFM/Vol11/jordan2c.html.

[16] Takaya Nishiyama and Yasuho Mizuhara. Binary arithmetics. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/binarith.html.

[17] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/pre_topc.html.

[18] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.

[19] Andrzej Trybulec. More on external approximation of a continuum. *Journal of Formalized Mathematics*, 13, 2001. http://mizar.org/JFM/Vol13/jordan1h.html.

[20] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. http://mizar.org/JFM/Addenda/numbers.html.

[21] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

[22] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received May 21, 2002

Published January 2, 2004