JOURNAL OF FORMALIZED MATHEMATICS
Volume 15, Released 2003, Published 2003
Inst. of Computer Science, Univ. of Białystok

The Class of Series-Parallel Graphs. Part II[1]

Krzysztof Retel
University of Białystok

Summary. In this paper we introduce two new operations on graphs: sum and union corresponding to parallel and series operation respectively. We determine N-free graph as the graph that does not embed Necklace 4. We define "fin_RelStr" as the set of all graphs with finite carriers. We also define the smallest class of graphs which contains the one-element graph and which is closed under parallel and series operations. The goal of the article is to prove the theorem that the class of finite series-parallel graphs is the class of finite N-free graphs. This paper formalizes the next part of [12].

MML Identifier: NECKLA_2.

WWW: http://mizar.org/JFM/Vol15/neckla_2.html

The articles [14], [13], [18], [7], [20], [8], [1], [2], [3], [15], [17], [4], [16], [19], [11], [5], [6], [9], and [10] provide the notation and terminology for this paper.

In this paper U is a universal class.

Next we state two propositions:

(1) For all sets X, Y such that $X \in U$ and $Y \in U$ and for every relation R between X and Y holds $R \in U$.

(2) The internal relation of Necklace 4 $= \{\langle 0,1\rangle, \langle 1,0\rangle, \langle 1,2\rangle, \langle 2,1\rangle, \langle 2,3\rangle, \langle 3,2\rangle\}$.

Let n be a natural number. Note that every element of $\mathbf{R}_n$ is finite.

One can prove the following proposition

(3) For every set x such that $x \in \mathbf{U}_0$ holds x is finite.

One can verify that every element of $\mathbf{U}_0$ is finite.

Let us mention that every number which is finite and ordinal is also natural.

Let G be a non empty relational structure. We say that G is N-free if and only if:

(Def. 1) G does not embed Necklace 4.

Let us mention that there exists a non empty relational structure which is strict, finite, and N-free.

Let R, S be relational structures. The functor UnionOf(R,S) yielding a strict relational structure is defined by the conditions (Def. 2).

(Def. 2)(i) The carrier of UnionOf(R,S) = (the carrier of R) $\cup$ (the carrier of S), and

(ii) the internal relation of UnionOf(R,S) = (the internal relation of R) $\cup$ (the internal relation of S).

[1] This work has been partially supported by CALCULEMUS grant HPRN-CT-2000-00102.

© Association of Mizar Users

Let R, S be relational structures. The functor SumOf(R,S) yields a strict relational structure and is defined by the conditions (Def. 3).

(Def. 3)(i) The carrier of SumOf(R,S) = (the carrier of R) $\cup$ (the carrier of S), and

(ii) the internal relation of SumOf(R,S) = (the internal relation of R) $\cup$ (the internal relation of S) $\cup$ [: the carrier of R, the carrier of S :] $\cup$ [: the carrier of S, the carrier of R :].

The functor FinRelStr is defined by the condition (Def. 4).

(Def. 4) Let X be a set. Then $X \in$ FinRelStr if and only if there exists a strict relational structure R such that $X = R$ and the carrier of $R \in \mathbf{U}_0$.

One can check that FinRelStr is non empty.
The subset FinRelStrSp of FinRelStr is defined by the conditions (Def. 5).

(Def. 5)(i) For every strict relational structure R such that the carrier of R is non empty and trivial and the carrier of $R \in \mathbf{U}_0$ holds $R \in$ FinRelStrSp,

(ii) for all strict relational structures H_1, H_2 such that the carrier of H_1 misses the carrier of H_2 and $H_1 \in$ FinRelStrSp and $H_2 \in$ FinRelStrSp holds UnionOf$(H_1,H_2) \in$ FinRelStrSp and SumOf$(H_1,H_2) \in$ FinRelStrSp, and

(iii) for every subset M of FinRelStr such that for every strict relational structure R such that the carrier of R is non empty and trivial and the carrier of $R \in \mathbf{U}_0$ holds $R \in M$ and for all strict relational structures H_1, H_2 such that the carrier of H_1 misses the carrier of H_2 and $H_1 \in M$ and $H_2 \in M$ holds UnionOf$(H_1,H_2) \in M$ and SumOf$(H_1,H_2) \in M$ holds FinRelStrSp $\subseteq M$.

Let us note that FinRelStrSp is non empty.
The following four propositions are true:

(4) For every set X such that $X \in$ FinRelStrSp holds X is a finite strict non empty relational structure.

(5) For every relational structure R such that $R \in$ FinRelStrSp holds the carrier of $R \in \mathbf{U}_0$.

(6) Let X be a set. Suppose $X \in$ FinRelStrSp. Then

(i) X is a strict non empty trivial relational structure, or

(ii) there exist strict relational structures H_1, H_2 such that the carrier of H_1 misses the carrier of H_2 and $H_1 \in$ FinRelStrSp and $H_2 \in$ FinRelStrSp and $X =$ UnionOf(H_1,H_2) or $X =$ SumOf(H_1,H_2).

(7) For every strict non empty relational structure R such that $R \in$ FinRelStrSp holds R is N-free.

REFERENCES

[1] Grzegorz Bancerek. Cardinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/card_1.html.

[2] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/ordinal1.html.

[3] Grzegorz Bancerek. Sequences of ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/ordinal2.html.

[4] Józef Białas. Group and field definitions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/realset1.html.

[5] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.

[6] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.

[7] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.

[8] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.

[9] Bogdan Nowak and Grzegorz Bancerek. Universal classes. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/classes2.html.

[10] Beata Padlewska. Families of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/setfam_1.html.

[11] Krzysztof Retel. The class of series — parallel graphs. *Journal of Formalized Mathematics*, 14, 2002. http://mizar.org/JFM/Vol14/necklace.html.

[12] Stephan Thomasse. On better-quasi-ordering countable series-parallel orders. *Transactions of the American Mathematical Society*, 352(6):2491–2505, 2000.

[13] Andrzej Trybulec. Enumerated sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/enumset1.html.

[14] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.

[15] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. http://mizar.org/JFM/Addenda/numbers.html.

[16] Wojciech A. Trybulec. Partially ordered sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/orders_1.html.

[17] Wojciech A. Trybulec. Groups. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/group_1.html.

[18] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

[19] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

[20] Edmund Woronowicz. Relations defined on sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relset_1.html.

Received May 29, 2003

Published January 2, 2004