JOURNAL OF FORMALIZED MATHEMATICS
Volume 2, Released 1990, Published 2003
Inst. of Computer Science, Univ. of Białystok

Elementary Variants of Affine Configurational Theorems[1]

Krzysztof Prażmowski
Warsaw University
Białystok

Krzysztof Radziszewski
Gdańsk University

Summary. We present elementary versions of Pappus, Major Desargues and Minor Desargues Axioms (i.e. statements formulated entirely in the language of points and parallelity of segments). Evidently they are consequences of appropriate configurational axioms introduced in the article [2]. In particular it follows that there exists an affine plane satisfying all of them.

MML Identifier: PARDEPAP.

WWW: http://mizar.org/JFM/Vol2/pardepap.html

The articles [4], [1], [3], and [2] provide the notation and terminology for this paper.

In this paper S_1 denotes an affine plane.

We now state several propositions:

(1) Suppose S_1 satisfies **PAP**. Let a_1, a_2, a_3, b_1, b_2, b_3 be elements of S_1. If $a_1,a_2 \parallel a_1,a_3$ and $b_1,b_2 \parallel b_1,b_3$ and $a_1,b_2 \parallel a_2,b_1$ and $a_2,b_3 \parallel a_3,b_2$, then $a_3,b_1 \parallel a_1,b_3$.

(2) Suppose S_1 satisfies **DES**. Let o, a, a', b, b', c, c' be elements of S_1. Suppose $o,a \nparallel o,b$ and $o,a \nparallel o,c$ and $o,a \parallel o,a'$ and $o,b \parallel o,b'$ and $o,c \parallel o,c'$ and $a,b \parallel a',b'$ and $a,c \parallel a',c'$. Then $b,c \parallel b',c'$.

(3) Suppose S_1 satisfies **des**. Let a, a', b, b', c, c' be elements of S_1. Suppose $a,a' \nparallel a,b$ and $a,a' \nparallel a,c$ and $a,a' \parallel b,b'$ and $a,a' \parallel c,c'$ and $a,b \parallel a',b'$ and $a,c \parallel a',c'$. Then $b,c \parallel b',c'$.

(5)[1] There exists S_1 such that

(i) for all elements o, a, a', b, b', c, c' of S_1 such that $o,a \nparallel o,b$ and $o,a \nparallel o,c$ and $o,a \parallel o,a'$ and $o,b \parallel o,b'$ and $o,c \parallel o,c'$ and $a,b \parallel a',b'$ and $a,c \parallel a',c'$ holds $b,c \parallel b',c'$,

(ii) for all elements a, a', b, b', c, c' of S_1 such that $a,a' \nparallel a,b$ and $a,a' \nparallel a,c$ and $a,a' \parallel b,b'$ and $a,a' \parallel c,c'$ and $a,b \parallel a',b'$ and $a,c \parallel a',c'$ holds $b,c \parallel b',c'$,

(iii) for all elements a_1, a_2, a_3, b_1, b_2, b_3 of S_1 such that $a_1,a_2 \parallel a_1,a_3$ and $b_1,b_2 \parallel b_1,b_3$ and $a_1,b_2 \parallel a_2,b_1$ and $a_2,b_3 \parallel a_3,b_2$ holds $a_3,b_1 \parallel a_1,b_3$, and

(iv) for all elements a, b, c, d of S_1 such that $a,b \nparallel a,c$ and $a,b \parallel c,d$ and $a,c \parallel b,d$ holds $a,d \nparallel b,c$.

(6) Let o, a be elements of S_1. Then there exists an element p of S_1 such that for all elements b, c of S_1 holds

[1] Supported by RPBP.III-24.C2.

[1] The proposition (4) has been removed.

© Association of Mizar Users

$o,a \nparallel o,p$ and there exists an element d of S_1 such that if $o,p \nparallel o,b$, then $o,c \nparallel o,d$ and $p,c \nparallel b,d$.

REFERENCES

[1] Henryk Oryszczyszyn and Krzysztof Prażmowski. Analytical ordered affine spaces. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/analoaf.html.

[2] Henryk Oryszczyszyn and Krzysztof Prażmowski. Classical configurations in affine planes. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/aff_2.html.

[3] Henryk Oryszczyszyn and Krzysztof Prażmowski. Ordered affine spaces defined in terms of directed parallelity — part I. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/diraf.html.

[4] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

Received November 30, 1990

Published January 2, 2004