JOURNAL OF FORMALIZED MATHEMATICS
Volume 12, Released 2000, Published 2003
Inst. of Computer Science, Univ. of Białystok

On Segre's Product of Partial Line Spaces

Adam Naumowicz
University of Białystok

Summary. In this paper the concept of partial line spaces is presented. We also construct the Segre's product for a family of partial line spaces indexed by an arbitrary nonempty set.

MML Identifier: PENCIL_1.

WWW: http://mizar.org/JFM/Vol12/pencil_1.html

The articles [16], [9], [19], [13], [2], [10], [20], [8], [6], [4], [1], [5], [3], [14], [17], [18], [7], [11], [12], and [15] provide the notation and terminology for this paper.

1. Preliminaries

The following propositions are true:

(1) For all functions f, g such that $\prod f = \prod g$ holds if f is non-empty, then g is non-empty.

(2) For every set X holds $2 \subseteq \overline{\overline{X}}$ iff there exist sets x, y such that $x \in X$ and $y \in X$ and $x \neq y$.

(3) For every set X such that $2 \subseteq \overline{\overline{X}}$ and for every set x there exists a set y such that $y \in X$ and $x \neq y$.

(4) For every set X holds $2 \subseteq \overline{\overline{X}}$ iff X is non trivial.

(5) For every set X holds $3 \subseteq \overline{\overline{X}}$ iff there exist sets x, y, z such that $x \in X$ and $y \in X$ and $z \in X$ and $x \neq y$ and $x \neq z$ and $y \neq z$.

(6) For every set X such that $3 \subseteq \overline{\overline{X}}$ and for all sets x, y there exists a set z such that $z \in X$ and $x \neq z$ and $y \neq z$.

2. Partial Line Spaces

Let S be a topological structure. A block of S is an element of the topology of S.

Let S be a topological structure and let x, y be points of S. We say that x, y are collinear if and only if:

(Def. 1) $x = y$ or there exists a block l of S such that $\{x, y\} \subseteq l$.

Let S be a topological structure and let T be a subset of S. We say that T is closed under lines if and only if:

(Def. 2) For every block l of S such that $2 \subseteq \overline{\overline{l \cap T}}$ holds $l \subseteq T$.

We say that T is strong if and only if:

(Def. 3) For all points x, y of S such that $x \in T$ and $y \in T$ holds x, y are collinear.

Let S be a topological structure. We say that S is void if and only if:

(Def. 4) The topology of S is empty.

We say that S is degenerated if and only if:

(Def. 5) The carrier of S is a block of S.

We say that S has non trivial blocks if and only if:

(Def. 6) For every block k of S holds $2 \subseteq \overline{\overline{k}}$.

We say that S is identifying close blocks if and only if:

(Def. 7) For all blocks k, l of S such that $2 \subseteq \overline{\overline{k \cap l}}$ holds $k = l$.

We say that S is truly-partial if and only if:

(Def. 8) There exist points x, y of S such that x, y are not collinear.

We say that S has no isolated points if and only if:

(Def. 9) For every point x of S there exists a block l of S such that $x \in l$.

We say that S is connected if and only if the condition (Def. 10) is satisfied.

(Def. 10) Let x, y be points of S. Then there exists a finite sequence f of elements of the carrier of S such that

(i) $x = f(1)$,

(ii) $y = f(\operatorname{len} f)$, and

(iii) for every natural number i such that $1 \leq i$ and $i < \operatorname{len} f$ and for all points a, b of S such that $a = f(i)$ and $b = f(i+1)$ holds a, b are collinear.

We say that S is strongly connected if and only if the condition (Def. 11) is satisfied.

(Def. 11) Let x be a point of S and X be a subset of S. Suppose X is closed under lines and strong. Then there exists a finite sequence f of elements of $2^{\text{the carrier of } S}$ such that

(i) $X = f(1)$,

(ii) $x \in f(\operatorname{len} f)$,

(iii) for every subset W of S such that $W \in \operatorname{rng} f$ holds W is closed under lines and strong, and

(iv) for every natural number i such that $1 \leq i$ and $i < \operatorname{len} f$ holds $2 \subseteq \overline{\overline{f(i) \cap f(i+1)}}$.

We now state two propositions:

(7) Let X be a non empty set. Suppose $3 \subseteq \overline{\overline{X}}$. Let S be a topological structure. Suppose the carrier of $S = X$ and the topology of $S = \{L; L$ ranges over subsets of X: $2 = \overline{\overline{L}}\}$. Then S is non empty, non void, non degenerated, non truly-partial, and identifying close blocks and has non trivial blocks and no isolated points.

(8) Let X be a non empty set. Suppose $3 \subseteq \overline{\overline{X}}$. Let K be a subset of X. Suppose $\overline{\overline{K}} = 2$. Let S be a topological structure. Suppose the carrier of $S = X$ and the topology of $S = \{L; L$ ranges over subsets of X: $2 = \overline{\overline{L}}\} \setminus \{K\}$. Then S is non empty, non void, non degenerated, truly-partial, and identifying close blocks and has non trivial blocks and no isolated points.

One can verify that there exists a topological structure which is strict, non empty, non void, non degenerated, non truly-partial, and identifying close blocks and has non trivial blocks and no isolated points and there exists a topological structure which is strict, non empty, non void, non degenerated, truly-partial, and identifying close blocks and has non trivial blocks and no isolated points.

Let S be a non void topological structure. Note that the topology of S is non empty.

Let S be a topological structure with no isolated points and let x, y be points of S. Let us observe that x, y are collinear if and only if:

(Def. 12) There exists a block l of S such that $\{x,y\} \subseteq l$.

A PLS is a non empty non void non degenerated identifying close blocks topological structure with non trivial blocks.

Let F be a binary relation. We say that F is TopStruct-yielding if and only if:

(Def. 13) For every set x such that $x \in \operatorname{rng} F$ holds x is a topological structure.

Let us note that every function which is TopStruct-yielding is also 1-sorted yielding.

Let I be a set. One can check that there exists a many sorted set indexed by I which is TopStruct-yielding.

Let us note that there exists a function which is TopStruct-yielding.

Let F be a binary relation. We say that F is non-void-yielding if and only if:

(Def. 14) For every topological structure S such that $S \in \operatorname{rng} F$ holds S is non void.

Let F be a TopStruct-yielding function. Let us observe that F is non-void-yielding if and only if:

(Def. 15) For every set i such that $i \in \operatorname{rng} F$ holds i is a non void topological structure.

Let F be a binary relation. We say that F is trivial-yielding if and only if:

(Def. 16) For every set S such that $S \in \operatorname{rng} F$ holds S is trivial.

Let F be a binary relation. We say that F is non-Trivial-yielding if and only if:

(Def. 17) For every 1-sorted structure S such that $S \in \operatorname{rng} F$ holds S is non trivial.

Let us observe that every binary relation which is non-Trivial-yielding is also nonempty.

Let F be a 1-sorted yielding function. Let us observe that F is non-Trivial-yielding if and only if:

(Def. 18) For every set i such that $i \in \operatorname{rng} F$ holds i is a non trivial 1-sorted structure.

Let I be a non empty set, let A be a TopStruct-yielding many sorted set indexed by I, and let j be an element of I. Then $A(j)$ is a topological structure.

Let F be a binary relation. We say that F is PLS-yielding if and only if:

(Def. 19) For every set x such that $x \in \operatorname{rng} F$ holds x is a PLS.

One can verify the following observations:

* every function which is PLS-yielding is also nonempty and TopStruct-yielding,
* every TopStruct-yielding function which is PLS-yielding is also non-void-yielding, and
* every TopStruct-yielding function which is PLS-yielding is also non-Trivial-yielding.

Let I be a set. One can verify that there exists a many sorted set indexed by I which is PLS-yielding.

Let I be a non empty set, let A be a PLS-yielding many sorted set indexed by I, and let j be an element of I. Then $A(j)$ is a PLS.

Let I be a set and let A be a many sorted set indexed by I. We say that A is Segre-like if and only if:

(Def. 20) There exists an element i of I such that for every element j of I such that $i \neq j$ holds $A(j)$ is non empty and trivial.

Let I be a set and let A be a many sorted set indexed by I. Note that $\{A\}$ is trivial-yielding.
The following proposition is true

(9) Let I be a non empty set, A be a many sorted set indexed by I, i be an element of I, and S be a non trivial set. Then $A +\cdot (i,S)$ is non trivial-yielding.

Let I be a non empty set and let A be a many sorted set indexed by I. Note that $\{A\}$ is Segre-like.
We now state two propositions:

(10) For every non empty set I and for every many sorted set A indexed by I and for all sets i, S holds $\{A\} +\cdot (i,S)$ is Segre-like.

(11) Let I be a non empty set, A be a nonempty 1-sorted yielding many sorted set indexed by I, and B be an element of the support of A. Then $\{B\}$ is a many sorted subset indexed by the support of A.

Let I be a non empty set and let A be a nonempty 1-sorted yielding many sorted set indexed by I. One can check that there exists a many sorted subset indexed by the support of A which is Segre-like, trivial-yielding, and non-empty.

Let I be a non empty set and let A be a non-Trivial-yielding 1-sorted yielding many sorted set indexed by I. Observe that there exists a many sorted subset indexed by the support of A which is Segre-like, non trivial-yielding, and non-empty.

Let I be a non empty set. One can verify that there exists a many sorted set indexed by I which is Segre-like and non trivial-yielding.

Let I be a non empty set and let B be a Segre-like non trivial-yielding many sorted set indexed by I. The functor index(B) yielding an element of I is defined as follows:

(Def. 21) $B(\text{index}(B))$ is non trivial.

The following proposition is true

(12) Let I be a non empty set, A be a Segre-like non trivial-yielding many sorted set indexed by I, and i be an element of I. If $i \neq \text{index}(A)$, then $A(i)$ is non empty and trivial.

Let I be a non empty set. One can check that every many sorted set indexed by I which is Segre-like and non trivial-yielding is also non-empty.

One can prove the following proposition

(13) Let I be a non empty set and A be a many sorted set indexed by I. Then $2 \subseteq \overline{\overline{\prod A}}$ if and only if A is non-empty and non trivial-yielding.

Let I be a non empty set and let B be a Segre-like non trivial-yielding many sorted set indexed by I. Note that $\prod B$ is non trivial.

3. SEGRE'S PRODUCT

Let I be a non empty set and let A be a nonempty TopStruct-yielding many sorted set indexed by I. The functor Segre_Blocks$A$ yields a family of subsets of $\prod$(the support of A) and is defined by the condition (Def. 22).

(Def. 22) Let x be a set. Then $x \in$ Segre_Blocks$A$ if and only if there exists a Segre-like many sorted subset B indexed by the support of A such that $x = \prod B$ and there exists an element i of I such that $B(i)$ is a block of $A(i)$.

Let I be a non empty set and let A be a nonempty TopStruct-yielding many sorted set indexed by I. The functor Segre_Product$A$ yields a non empty topological structure and is defined as follows:

(Def. 23) Segre_Product$A = \langle\prod$(the support of A),Segre_Blocks$A\rangle$.

The following propositions are true:

(14) Let I be a non empty set and A be a nonempty TopStruct-yielding many sorted set indexed by I. Then every point of Segre_Product$A$ is a many sorted set indexed by I.

(15) Let I be a non empty set and A be a nonempty TopStruct-yielding many sorted set indexed by I. If there exists an element i of I such that $A(i)$ is non void, then Segre_Product$A$ is non void.

(16) Let I be a non empty set and A be a nonempty TopStruct-yielding many sorted set indexed by I. Suppose that for every element i of I holds $A(i)$ is non degenerated and there exists an element i of I such that $A(i)$ is non void. Then Segre_Product$A$ is non degenerated.

(17) Let I be a non empty set and A be a nonempty TopStruct-yielding many sorted set indexed by I. Suppose that for every element i of I holds $A(i)$ has non trivial blocks and there exists an element i of I such that $A(i)$ is non void. Then Segre_Product$A$ has non trivial blocks.

(18) Let I be a non empty set and A be a nonempty TopStruct-yielding many sorted set indexed by I. Suppose that for every element i of I holds $A(i)$ is identifying close blocks and has non trivial blocks and there exists an element i of I such that $A(i)$ is non void. Then Segre_Product$A$ is identifying close blocks.

Let I be a non empty set and let A be a PLS-yielding many sorted set indexed by I. Then Segre_Product$A$ is a PLS.

Next we state a number of propositions:

(19) Let T be a topological structure and S be a subset of T. If S is trivial, then S is strong and closed under lines.

(20) Let S be an identifying close blocks topological structure, l be a block of S, and L be a subset of S. If $L = l$, then L is closed under lines.

(21) Let S be a topological structure, l be a block of S, and L be a subset of S. If $L = l$, then L is strong.

(22) For every non void topological structure S holds Ω_S is closed under lines.

(23) Let I be a non empty set, A be a Segre-like non trivial-yielding many sorted set indexed by I, and x, y be many sorted sets indexed by I. If $x \in \prod A$ and $y \in \prod A$, then for every set i such that $i \neq \text{index}(A)$ holds $x(i) = y(i)$.

(24) Let I be a non empty set, A be a PLS-yielding many sorted set indexed by I, and x be a set. Then x is a block of Segre_Product$A$ if and only if there exists a Segre-like non trivial-yielding many sorted subset L indexed by the support of A such that $x = \prod L$ and $L(\text{index}(L))$ is a block of $A(\text{index}(L))$.

(25) Let I be a non empty set, A be a PLS-yielding many sorted set indexed by I, and P be a many sorted set indexed by I. Suppose P is a point of Segre_Product$A$. Let i be an element of I and p be a point of $A(i)$. Then $P +\cdot (i, p)$ is a point of Segre_Product$A$.

(26) Let I be a non empty set and A, B be Segre-like non trivial-yielding many sorted sets indexed by I. Suppose $2 \subseteq \overline{\overline{\prod A \cap \prod B}}$. Then $\text{index}(A) = \text{index}(B)$ and for every set i such that $i \neq \text{index}(A)$ holds $A(i) = B(i)$.

(27) Let I be a non empty set, A be a Segre-like non trivial-yielding many sorted set indexed by I, and N be a non trivial set. Then $A +\cdot (\text{index}(A), N)$ is Segre-like and non trivial-yielding.

(28) Let S be a non empty non void identifying close blocks topological structure with no isolated points. If S is strongly connected, then S is connected.

(29) Let I be a non empty set, A be a PLS-yielding many sorted set indexed by I, and S be a subset of Segre_Product$A$. Then S is non trivial, strong, and closed under lines if and only if there exists a Segre-like non trivial-yielding many sorted subset B indexed by the support of A such that $S = \prod B$ and for every subset C of $A(\mathrm{index}(B))$ such that $C = B(\mathrm{index}(B))$ holds C is strong and closed under lines.

REFERENCES

[1] Grzegorz Bancerek. Cardinal numbers. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/card_1.html`.

[2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/nat_1.html`.

[3] Grzegorz Bancerek. König's theorem. *Journal of Formalized Mathematics*, 2, 1990. `http://mizar.org/JFM/Vol2/card_3.html`.

[4] Grzegorz Bancerek. The "way-below" relation. *Journal of Formalized Mathematics*, 8, 1996. `http://mizar.org/JFM/Vol8/waybel_3.html`.

[5] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/finseq_1.html`.

[6] Józef Białas. Group and field definitions. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/realset1.html`.

[7] Ewa Burakowska. Subalgebras of many sorted algebra. Lattice of subalgebras. *Journal of Formalized Mathematics*, 6, 1994. `http://mizar.org/JFM/Vol6/msualg_2.html`.

[8] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/funct_1.html`.

[9] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/zfmisc_1.html`.

[10] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/finset_1.html`.

[11] Artur Korniłowicz. Some basic properties of many sorted sets. *Journal of Formalized Mathematics*, 7, 1995. `http://mizar.org/JFM/Vol7/pzfmisc1.html`.

[12] Beata Madras. Product of family of universal algebras. *Journal of Formalized Mathematics*, 5, 1993. `http://mizar.org/JFM/Vol5/pralg_1.html`.

[13] Beata Padlewska. Families of sets. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/setfam_1.html`.

[14] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/pre_topc.html`.

[15] Piotr Rudnicki and Andrzej Trybulec. Multivariate polynomials with arbitrary number of variables. *Journal of Formalized Mathematics*, 11, 1999. `http://mizar.org/JFM/Vol11/polynom1.html`.

[16] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. `http://mizar.org/JFM/Axiomatics/tarski.html`.

[17] Andrzej Trybulec. Many-sorted sets. *Journal of Formalized Mathematics*, 5, 1993. `http://mizar.org/JFM/Vol5/pboole.html`.

[18] Andrzej Trybulec. Many sorted algebras. *Journal of Formalized Mathematics*, 6, 1994. `http://mizar.org/JFM/Vol6/msualg_1.html`.

[19] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/subset_1.html`.

[20] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/relat_1.html`.

Received May 29, 2000

Published January 2, 2004