JOURNAL OF FORMALIZED MATHEMATICS
Volume 5, Released 1993, Published 2003
Inst. of Computer Science, Univ. of Białystok

Product of Family of Universal Algebras

Beata Madras
Warsaw University
Białystok

Summary. The product of two algebras, trivial algebra determined by an empty set and product of a family of algebras are defined. Some basic properties are shown.

MML Identifier: `PRALG_1`.

WWW: `http://mizar.org/JFM/Vol5/pralg_1.html`

The articles [17], [11], [21], [1], [2], [20], [22], [8], [5], [9], [15], [18], [16], [10], [12], [13], [3], [4], [6], [14], [7], and [19] provide the notation and terminology for this paper.

1. PRODUCT OF TWO ALGEBRAS

One can prove the following proposition

(1) For all non empty sets D_1, D_2 and for all natural numbers n, m such that ${D_1}^n = {D_2}^m$ holds $n = m$.

For simplicity, we follow the rules: U_1, U_2 denote universal algebras, n, m denote natural numbers, x, y denote sets, A, B denote non empty sets, and h_1 denotes a finite sequence of elements of $[:A, B:]$.

Let us consider A, B and let us consider h_1. Then $\mathrm{pr1}(h_1)$ is a finite sequence of elements of A and it can be characterized by the condition:

(Def. 1) $\mathrm{len}\,\mathrm{pr1}(h_1) = \mathrm{len}\,h_1$ and for every n such that $n \in \mathrm{dom}\,\mathrm{pr1}(h_1)$ holds $\mathrm{pr1}(h_1)(n) = h_1(n)_{\mathbf{1}}$.

Then $\mathrm{pr2}(h_1)$ is a finite sequence of elements of B and it can be characterized by the condition:

(Def. 2) $\mathrm{len}\,\mathrm{pr2}(h_1) = \mathrm{len}\,h_1$ and for every n such that $n \in \mathrm{dom}\,\mathrm{pr2}(h_1)$ holds $\mathrm{pr2}(h_1)(n) = h_1(n)_{\mathbf{2}}$.

Let us consider A, B, let f_1 be a homogeneous quasi total non empty partial function from A^* to A, and let f_2 be a homogeneous quasi total non empty partial function from B^* to B. Let us assume that $\mathrm{arity}\,f_1 = \mathrm{arity}\,f_2$. The functor $]\!]f_1, f_2[\![$ yields a homogeneous quasi total non empty partial function from $[:A, B:]^*$ to $[:A, B:]$ and is defined by the conditions (Def. 3).

(Def. 3)(i) $\mathrm{dom}\,]\!]f_1, f_2[\![= [:A, B:]^{\mathrm{arity}\,f_1}$, and

(ii) for every finite sequence h of elements of $[:A, B:]$ such that $h \in \mathrm{dom}\,]\!]f_1, f_2[\![$ holds $]\!]f_1, f_2[\![(h) = \langle f_1(\mathrm{pr1}(h)), f_2(\mathrm{pr2}(h))\rangle$.

In the sequel h_1 denotes a homogeneous quasi total non empty partial function from (the carrier of U_1)* to the carrier of U_1 and h_2 denotes a homogeneous quasi total non empty partial function from (the carrier of U_2)* to the carrier of U_2.

Let us consider U_1, U_2. Let us assume that U_1 and U_2 are similar. The functor $\mathrm{Opers}(U_1, U_2)$ yields a finite sequence of operational functions of [:the carrier of U_1, the carrier of U_2:] and is defined by the conditions (Def. 4).

(Def. 4)(i) $\operatorname{len}\operatorname{Opers}(U_1,U_2) = \operatorname{len}(\text{the characteristic of } U_1)$, and

(ii) for every n such that $n \in \operatorname{dom}\operatorname{Opers}(U_1,U_2)$ and for all h_1, h_2 such that $h_1 =$ (the characteristic of U_1)(n) and $h_2 =$ (the characteristic of U_2)(n) holds $(\operatorname{Opers}(U_1,U_2))(n) =]\!]h_1,h_2[\![$.

One can prove the following proposition

(2) If U_1 and U_2 are similar, then $\langle [\!:\text{the carrier of } U_1, \text{ the carrier of } U_2:\!], \operatorname{Opers}(U_1,U_2)\rangle$ is a strict universal algebra.

Let us consider U_1, U_2. Let us assume that U_1 and U_2 are similar. The functor $[\!:U_1, U_2:\!]$ yielding a strict universal algebra is defined as follows:

(Def. 5) $[\!:U_1, U_2:\!] = \langle [\!:\text{the carrier of } U_1, \text{ the carrier of } U_2:\!], \operatorname{Opers}(U_1,U_2)\rangle$.

Let A, B be non empty sets. The functor $\operatorname{Inv}(A,B)$ yielding a function from $[\!:A, B:\!]$ into $[\!:B, A:\!]$ is defined as follows:

(Def. 6) For every element a of $[\!:A, B:\!]$ holds $(\operatorname{Inv}(A,B))(a) = \langle a_{\mathbf{2}}, a_{\mathbf{1}}\rangle$.

Next we state several propositions:

(3) For all non empty sets A, B holds $\operatorname{rng}\operatorname{Inv}(A,B) = [\!:B, A:\!]$.

(4) For all non empty sets A, B holds $\operatorname{Inv}(A,B)$ is one-to-one.

(5) Suppose U_1 and U_2 are similar. Then Inv(the carrier of U_1, the carrier of U_2) is a function from the carrier of $[\!:U_1, U_2:\!]$ into the carrier of $[\!:U_2, U_1:\!]$.

(6) Suppose U_1 and U_2 are similar. Let o_1 be an operation of U_1, o_2 be an operation of U_2, o be an operation of $[\!:U_1, U_2:\!]$, and n be a natural number. Suppose that

(i) $o_1 =$ (the characteristic of U_1)(n),

(ii) $o_2 =$ (the characteristic of U_2)(n),

(iii) $o =$ (the characteristic of $[\!:U_1, U_2:\!]$)(n), and

(iv) $n \in \operatorname{dom}(\text{the characteristic of } U_1)$.

Then $\operatorname{arity} o = \operatorname{arity} o_1$ and $\operatorname{arity} o = \operatorname{arity} o_2$ and $o =]\!]o_1, o_2[\![$.

(7) If U_1 and U_2 are similar, then $[\!:U_1, U_2:\!]$ and U_1 are similar.

(8) Let U_1, U_2, U_3, U_4 be universal algebras. Suppose U_1 is a subalgebra of U_2 and U_3 is a subalgebra of U_4 and U_2 and U_4 are similar. Then $[\!:U_1, U_3:\!]$ is a subalgebra of $[\!:U_2, U_4:\!]$.

2. Trivial Algebra

Let k be a natural number. The functor $\operatorname{TrivOp}(k)$ yielding a partial function from $\{\emptyset\}^*$ to $\{\emptyset\}$ is defined by:

(Def. 7) $\operatorname{dom}\operatorname{TrivOp}(k) = \{k \mapsto \emptyset\}$ and $\operatorname{rng}\operatorname{TrivOp}(k) = \{\emptyset\}$.

Let k be a natural number. Note that $\operatorname{TrivOp}(k)$ is homogeneous, quasi total, and non empty.
The following proposition is true

(9) For every natural number k holds $\operatorname{arity}\operatorname{TrivOp}(k) = k$.

Let f be a finite sequence of elements of $\mathbb{N}$. The functor $\operatorname{TrivOps}(f)$ yields a finite sequence of operational functions of $\{\emptyset\}$ and is defined as follows:

(Def. 8) $\operatorname{len}\operatorname{TrivOps}(f) = \operatorname{len} f$ and for every n such that $n \in \operatorname{dom}\operatorname{TrivOps}(f)$ and for every m such that $m = f(n)$ holds $(\operatorname{TrivOps}(f))(n) = \operatorname{TrivOp}(m)$.

One can prove the following two propositions:

(10) For every finite sequence f of elements of $\mathbb{N}$ holds $\mathrm{TrivOps}(f)$ is homogeneous, quasi total, and non-empty.

(11) For every finite sequence f of elements of $\mathbb{N}$ such that $f \neq \emptyset$ holds $\langle \{\emptyset\}, \mathrm{TrivOps}(f) \rangle$ is a strict universal algebra.

Let D be a non empty set. Observe that there exists a finite sequence of elements of D which is non empty and there exists an element of D^* which is non empty.

Let f be a non empty finite sequence of elements of $\mathbb{N}$. The trivial algebra of f yielding a strict universal algebra is defined as follows:

(Def. 9) The trivial algebra of $f = \langle \{\emptyset\}, \mathrm{TrivOps}(f) \rangle$.

3. Product of Universal Algebras

Let I_1 be a function. We say that I_1 is universal algebra yielding if and only if:

(Def. 10) For every x such that $x \in \mathrm{dom}\, I_1$ holds $I_1(x)$ is a universal algebra.

Let I_1 be a function. We say that I_1 is 1-sorted yielding if and only if:

(Def. 11) For every x such that $x \in \mathrm{dom}\, I_1$ holds $I_1(x)$ is a 1-sorted structure.

Let us mention that there exists a function which is universal algebra yielding.

Let us mention that every function which is universal algebra yielding is also 1-sorted yielding.

Let I be a set. Observe that there exists a many sorted set indexed by I which is 1-sorted yielding.

Let I_1 be a function. We say that I_1 is equal signature if and only if:

(Def. 12) For all x, y such that $x \in \mathrm{dom}\, I_1$ and $y \in \mathrm{dom}\, I_1$ and for all U_1, U_2 such that $U_1 = I_1(x)$ and $U_2 = I_1(y)$ holds $\mathrm{signature}\, U_1 = \mathrm{signature}\, U_2$.

Let J be a non empty set. One can verify that there exists a many sorted set indexed by J which is equal signature and universal algebra yielding.

Let J be a non empty set, let A be a 1-sorted yielding many sorted set indexed by J, and let j be an element of J. Then $A(j)$ is a 1-sorted structure.

Let J be a non empty set, let A be a universal algebra yielding many sorted set indexed by J, and let j be an element of J. Then $A(j)$ is a universal algebra.

Let J be a set and let A be a 1-sorted yielding many sorted set indexed by J. The support of A yielding a many sorted set indexed by J is defined by the condition (Def. 13).

(Def. 13) Let j be a set. Suppose $j \in J$. Then there exists a 1-sorted structure R such that $R = A(j)$ and (the support of $A)(j) =$ the carrier of R.

Let J be a non empty set and let A be a universal algebra yielding many sorted set indexed by J. One can check that the support of A is non-empty.

Let J be a non empty set and let A be an equal signature universal algebra yielding many sorted set indexed by J. The functor $\mathrm{ComSign}(A)$ yields a finite sequence of elements of $\mathbb{N}$ and is defined as follows:

(Def. 14) For every element j of J holds $\mathrm{ComSign}(A) = \mathrm{signature}\, A(j)$.

Let I_1 be a function. We say that I_1 is function yielding if and only if:

(Def. 15) For every x such that $x \in \mathrm{dom}\, I_1$ holds $I_1(x)$ is a function.

Let us observe that there exists a function which is function yielding.

Let I be a set. One can check that there exists a many sorted set indexed by I which is function yielding.

Let I be a set. A many sorted function indexed by I is a function yielding many sorted set indexed by I.

Let B be a function yielding function and let j be a set. One can verify that $B(j)$ is function-like and relation-like.

Let J be a non empty set, let B be a non-empty many sorted set indexed by J, and let j be an element of J. Observe that $B(j)$ is non empty.

Let F be a function yielding function and let f be a function. Note that $F \cdot f$ is function yielding.

Let J be a non empty set and let B be a non-empty many sorted set indexed by J. Note that $\prod B$ is non empty.

Let J be a non empty set and let B be a non-empty many sorted set indexed by J. A many sorted function indexed by J is said to be a many sorted operation of B if:

(Def. 16) For every element j of J holds $\mathrm{it}(j)$ is a homogeneous quasi total non empty partial function from $B(j)^*$ to $B(j)$.

Let J be a non empty set, let B be a non-empty many sorted set indexed by J, let O be a many sorted operation of B, and let j be an element of J. Then $O(j)$ is a homogeneous quasi total non empty partial function from $B(j)^*$ to $B(j)$.

Let I_1 be a function. We say that I_1 is equal arity if and only if the condition (Def. 17) is satisfied.

(Def. 17) Let x, y be sets. Suppose $x \in \mathrm{dom} I_1$ and $y \in \mathrm{dom} I_1$. Let f, g be functions. Suppose $I_1(x) = f$ and $I_1(y) = g$. Let n, m be natural numbers and X, Y be non empty sets. Suppose $\mathrm{dom} f = X^n$ and $\mathrm{dom} g = Y^m$. Let o_1 be a homogeneous quasi total non empty partial function from X^* to X and o_2 be a homogeneous quasi total non empty partial function from Y^* to Y. If $f = o_1$ and $g = o_2$, then $\mathrm{arity}\, o_1 = \mathrm{arity}\, o_2$.

Let J be a non empty set and let B be a non-empty many sorted set indexed by J. Note that there exists a many sorted operation of B which is equal arity.

The following proposition is true

(12) Let J be a non empty set, B be a non-empty many sorted set indexed by J, and O be a many sorted operation of B. Then O is equal arity if and only if for all elements i, j of J holds $\mathrm{arity}\, O(i) = \mathrm{arity}\, O(j)$.

Let F be a function yielding function and let f be a function. The functor $F \looparrowleft f$ yields a function and is defined as follows:

(Def. 18) $\mathrm{dom}(F \looparrowleft f) = \mathrm{dom} F$ and for every set x such that $x \in \mathrm{dom} F$ holds $(F \looparrowleft f)(x) = F(x)(f(x))$.

Let I be a set, let f be a many sorted function indexed by I, and let x be a many sorted set indexed by I. Then $f \looparrowleft x$ is a many sorted set indexed by I and it can be characterized by the condition:

(Def. 19) For every set i such that $i \in I$ and for every function g such that $g = f(i)$ holds $(f \looparrowleft x)(i) = g(x(i))$.

Let J be a non empty set, let B be a non-empty many sorted set indexed by J, and let p be a finite sequence of elements of $\prod B$. Then $\mathrm{uncurry}\, p$ is a many sorted set indexed by $[: \mathrm{dom}\, p, J :]$.

Let I, J be sets and let X be a many sorted set indexed by $[: I, J :]$. Then $\curvearrowleft X$ is a many sorted set indexed by $[: J, I :]$.

Let X be a set, let Y be a non empty set, and let f be a many sorted set indexed by $[: X, Y :]$. Then $\mathrm{curry}\, f$ is a many sorted set indexed by X.

Let J be a non empty set, let B be a non-empty many sorted set indexed by J, and let O be an equal arity many sorted operation of B. The functor $\mathrm{ComAr}(O)$ yields a natural number and is defined as follows:

(Def. 20) For every element j of J holds $\mathrm{ComAr}(O) = \mathrm{arity}\, O(j)$.

Let I be a set and let A be a many sorted set indexed by I. The functor ε_A yielding a many sorted set indexed by I is defined as follows:

(Def. 21) For every set i such that $i \in I$ holds $\varepsilon_A(i) = \emptyset_{A(i)}$.

Let J be a non empty set, let B be a non-empty many sorted set indexed by J, and let O be an equal arity many sorted operation of B. The functor $]]O[[$ yields a homogeneous quasi total non empty partial function from $(\prod B)^*$ to $\prod B$ and is defined by the conditions (Def. 22).

(Def. 22)(i) $\mathrm{dom}]]O[[= (\prod B)^{\mathrm{ComAr}(O)}$, and

(ii) for every element p of $(\prod B)^*$ such that $p \in \mathrm{dom}]]O[[$ holds if $\mathrm{dom}\, p = \emptyset$, then $]]O[[(p) = O \hookleftarrow (\varepsilon_B)$ and if $\mathrm{dom}\, p \neq \emptyset$, then for every non empty set Z and for every many sorted set w indexed by $[:J, Z:]$ such that $Z = \mathrm{dom}\, p$ and $w = \curvearrowleft \mathrm{uncurry}\, p$ holds $]]O[[(p) = O \hookleftarrow \mathrm{curry}\, w$.

Let J be a non empty set, let A be an equal signature universal algebra yielding many sorted set indexed by J, and let n be a natural number. Let us assume that $n \in \mathrm{dom}\,\mathrm{ComSign}(A)$. The functor $\mathrm{ProdOp}(A,n)$ yielding an equal arity many sorted operation of the support of A is defined by the condition (Def. 23).

(Def. 23) Let j be an element of J and o be an operation of $A(j)$. If (the characteristic of $A(j))(n) = o$, then $(\mathrm{ProdOp}(A,n))(j) = o$.

Let J be a non empty set and let A be an equal signature universal algebra yielding many sorted set indexed by J. The functor $\mathrm{ProdOpSeq}(A)$ yields a finite sequence of operational functions of $\prod$(the support of A) and is defined as follows:

(Def. 24) $\mathrm{len}\,\mathrm{ProdOpSeq}(A) = \mathrm{len}\,\mathrm{ComSign}(A)$ and for every n such that $n \in \mathrm{dom}\,\mathrm{ProdOpSeq}(A)$ holds $(\mathrm{ProdOpSeq}(A))(n) =]]\mathrm{ProdOp}(A,n)[[$.

Let J be a non empty set and let A be an equal signature universal algebra yielding many sorted set indexed by J. The functor $\mathrm{ProdUnivAlg}(A)$ yields a strict universal algebra and is defined by:

(Def. 25) $\mathrm{ProdUnivAlg}(A) = \langle \prod(\text{the support of } A), \mathrm{ProdOpSeq}(A)\rangle$.

REFERENCES

[1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/ordinal1.html.

[2] Grzegorz Bancerek. Sequences of ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/ordinal2.html.

[3] Grzegorz Bancerek. Curried and uncurried functions. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/funct_5.html.

[4] Grzegorz Bancerek. König's theorem. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/card_3.html.

[5] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.

[6] Grzegorz Bancerek and Piotr Rudnicki. On defining functions on trees. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/dtconstr.html.

[7] Ewa Burakowska. Subalgebras of the universal algebra. Lattices of subalgebras. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/unialg_2.html.

[8] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.

[9] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.

[10] Czesław Byliński. Partial functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/partfun1.html.

[11] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.

[12] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/finseq_2.html.

[13] Czesław Byliński. The modification of a function by a function and the iteration of the composition of a function. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/funct_4.html.

[14] Jarosław Kotowicz, Beata Madras, and Małgorzata Korolkiewicz. Basic notation of universal algebra. *Journal of Formalized Mathematics*, 4, 1992. http://mizar.org/JFM/Vol4/unialg_1.html.

[15] Andrzej Trybulec. Binary operations applied to functions. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/funcop_1.html`.

[16] Andrzej Trybulec. Domains and their Cartesian products. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/domain_1.html`.

[17] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. `http://mizar.org/JFM/Axiomatics/tarski.html`.

[18] Andrzej Trybulec. Tuples, projections and Cartesian products. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/mcart_1.html`.

[19] Andrzej Trybulec. Many-sorted sets. *Journal of Formalized Mathematics*, 5, 1993. `http://mizar.org/JFM/Vol5/pboole.html`.

[20] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. `http://mizar.org/JFM/Addenda/numbers.html`.

[21] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/subset_1.html`.

[22] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/relat_1.html`.

Received October 12, 1993

Published January 2, 2004