JOURNAL OF FORMALIZED MATHEMATICS
Volume 10, Released 1998, Published 2003
Inst. of Computer Science, Univ. of Białystok

Trigonometric Functions and Existence of Circle Ratio

Yuguang Yang
Shinshu University
Nagano

Yasunari Shidama
Shinshu University
Nagano

Summary. In this article, we defined *sinus* and *cosine* as the real part and the imaginary part of the exponential function on complex, and also give their series expression. Then we proved the differentiablity of *sinus*, *cosine* and the exponential function of real. Finally, we showed the existence of the circle ratio, and some formulas of *sinus*, *cosine*.

MML Identifier: `SIN_COS`.

WWW: `http://mizar.org/JFM/Vol10/sin_cos.html`

The articles [23], [26], [3], [24], [7], [8], [4], [19], [5], [11], [9], [20], [27], [18], [16], [2], [13], [6], [25], [17], [10], [21], [15], [12], [1], [14], and [22] provide the notation and terminology for this paper.

1. Some Definitions and Properties of Complex Sequence

For simplicity, we use the following convention: p, q, t_1, t_2, t_3 denote real numbers, w, z, z_1, z_2 denote elements of $\mathbb{C}$, k, l, m, n denote natural numbers, s_1 denotes a complex sequence, and r_1 denotes a sequence of real numbers.

Let m, k be natural numbers. The functor $\mathrm{CHK}(m,k)$ yielding an element of $\mathbb{C}$ is defined by:

(Def. 2)[1] $$\mathrm{CHK}(m,k) = \begin{cases} 1_{\mathbb{C}}, \text{ if } m \leq k, \\ 0_{\mathbb{C}}, \text{ otherwise.} \end{cases}$$

The functor $\mathrm{RHK}(m,k)$ is defined as follows:

(Def. 3) $$\mathrm{RHK}(m,k) = \begin{cases} 1, \text{ if } m \leq k, \\ 0, \text{ otherwise.} \end{cases}$$

Let m, k be natural numbers. One can check that $\mathrm{RHK}(m,k)$ is real.

Let m, k be natural numbers. Then $\mathrm{RHK}(m,k)$ is a real number.

In this article we present several logical schemes. The scheme *ExComplex CASE* deals with a binary functor $\mathcal{F}$ yielding an element of $\mathbb{C}$, and states that:

For every k there exists s_1 such that for every n holds if $n \leq k$, then $s_1(n) = \mathcal{F}(k,n)$ and if $n > k$, then $s_1(n) = 0_{\mathbb{C}}$

for all values of the parameter.

The scheme *ExReal CASE* deals with a binary functor $\mathcal{F}$ yielding a real number, and states that:

For every k there exists r_1 such that for every n holds if $n \leq k$, then $r_1(n) = \mathcal{F}(k,n)$ and if $n > k$, then $r_1(n) = 0$

[1] The definition (Def. 1) has been removed.

for all values of the parameter.

The complex sequence Prod_complex_n is defined by:

(Def. 4) $(\text{Prod\_complex\_n})(0) = 1_{\mathbb{C}}$ and for every n holds $(\text{Prod\_complex\_n})(n+1) = (\text{Prod\_complex\_n})(n) \cdot ((n+1)+0i)$.

The sequence Prod_real_n of real numbers is defined as follows:

(Def. 5) $(\text{Prod\_real\_n})(0) = 1$ and for every n holds $(\text{Prod\_real\_n})(n+1) = (\text{Prod\_real\_n})(n) \cdot (n+1)$.

Let n be a natural number. The functor $n!_{\mathbb{C}}$ yields an element of $\mathbb{C}$ and is defined as follows:

(Def. 6) $n!_{\mathbb{C}} = (\text{Prod\_complex\_n})(n)$.

Let n be a natural number. Then $n!$ is a real number and it can be characterized by the condition:

(Def. 7) $n! = (\text{Prod\_real\_n})(n)$.

Let z be an element of $\mathbb{C}$. The functor $z\,\text{ExpSeq}$ yielding a complex sequence is defined as follows:

(Def. 8) For every n holds $z\,\text{ExpSeq}(n) = \frac{z^n_{\mathbb{N}}}{n!_{\mathbb{C}}}$.

Let a be a real number. The functor $a\,\text{ExpSeq}$ yields a sequence of real numbers and is defined by:

(Def. 9) For every n holds $a\,\text{ExpSeq}(n) = \frac{a^n}{n!}$.

One can prove the following three propositions:

(1) If $0 < n$, then $n+0i \neq 0_{\mathbb{C}}$ and $0!_{\mathbb{C}} = 1_{\mathbb{C}}$ and $n!_{\mathbb{C}} \neq 0_{\mathbb{C}}$ and $(n+1)!_{\mathbb{C}} = n!_{\mathbb{C}} \cdot ((n+1)+0i)$.

(2) $n! \neq 0$ and $(n+1)! = n! \cdot (n+1)$.

(3) For every k such that $0 < k$ holds $(k -' 1)!_{\mathbb{C}} \cdot (k+0i) = k!_{\mathbb{C}}$ and for all m, k such that $k \leq m$ holds $(m -' k)!_{\mathbb{C}} \cdot (((m+1)-k)+0i) = ((m+1) -' k)!_{\mathbb{C}}$.

Let n be a natural number. The functor $\text{Coef}\,n$ yields a complex sequence and is defined as follows:

(Def. 10) For every natural number k holds if $k \leq n$, then $(\text{Coef}\,n)(k) = \frac{n!_{\mathbb{C}}}{k!_{\mathbb{C}} \cdot (n -' k)!_{\mathbb{C}}}$ and if $k > n$, then $(\text{Coef}\,n)(k) = 0_{\mathbb{C}}$.

Let n be a natural number. The functor $\text{Coef\_e}\,n$ yielding a complex sequence is defined as follows:

(Def. 11) For every natural number k holds if $k \leq n$, then $(\text{Coef\_e}\,n)(k) = \frac{1_{\mathbb{C}}}{k!_{\mathbb{C}} \cdot (n -' k)!_{\mathbb{C}}}$ and if $k > n$, then $(\text{Coef\_e}\,n)(k) = 0_{\mathbb{C}}$.

Let us consider s_1. The functor $\text{Sift}\,s_1$ yielding a complex sequence is defined by:

(Def. 12) $(\text{Sift}\,s_1)(0) = 0_{\mathbb{C}}$ and for every natural number k holds $(\text{Sift}\,s_1)(k+1) = s_1(k)$.

Let us consider n and let z, w be elements of $\mathbb{C}$. The functor $\text{Expan}(n,z,w)$ yields a complex sequence and is defined by:

(Def. 13) For every natural number k holds if $k \leq n$, then $(\text{Expan}(n,z,w))(k) = (\text{Coef}\,n)(k) \cdot z^k_{\mathbb{N}} \cdot w^{n -' k}_{\mathbb{N}}$ and if $n < k$, then $(\text{Expan}(n,z,w))(k) = 0_{\mathbb{C}}$.

Let us consider n and let z, w be elements of $\mathbb{C}$. The functor $\text{Expan\_e}(n,z,w)$ yields a complex sequence and is defined as follows:

(Def. 14) For every natural number k holds if $k \leq n$, then $(\text{Expan\_e}(n,z,w))(k) = (\text{Coef\_e}\,n)(k) \cdot z^k_{\mathbb{N}} \cdot w^{n -' k}_{\mathbb{N}}$ and if $n < k$, then $(\text{Expan\_e}(n,z,w))(k) = 0_{\mathbb{C}}$.

Let us consider n and let z, w be elements of $\mathbb{C}$. The functor $\mathrm{Alfa}(n,z,w)$ yields a complex sequence and is defined as follows:

(Def. 15) For every natural number k holds if $k \leq n$, then $(\mathrm{Alfa}(n,z,w))(k) = z\,\mathrm{ExpSeq}(k) \cdot (\sum_{\alpha=0}^{\kappa} w\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n -' k)$ and if $n < k$, then $(\mathrm{Alfa}(n,z,w))(k) = 0_{\mathbb{C}}$.

Let a, b be real numbers and let n be a natural number. The functor $\mathrm{Conj}(n,a,b)$ yielding a sequence of real numbers is defined as follows:

(Def. 16) For every natural number k holds if $k \leq n$, then $(\mathrm{Conj}(n,a,b))(k) = a\,\mathrm{ExpSeq}(k) \cdot ((\sum_{\alpha=0}^{\kappa} b\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n) - (\sum_{\alpha=0}^{\kappa} b\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n -' k))$ and if $n < k$, then $(\mathrm{Conj}(n,a,b))(k) = 0$.

Let z, w be elements of $\mathbb{C}$ and let n be a natural number. The functor $\mathrm{Conj}(n,z,w)$ yields a complex sequence and is defined as follows:

(Def. 17) For every natural number k holds if $k \leq n$, then $(\mathrm{Conj}(n,z,w))(k) = z\,\mathrm{ExpSeq}(k) \cdot ((\sum_{\alpha=0}^{\kappa} w\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n) - (\sum_{\alpha=0}^{\kappa} w\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n -' k))$ and if $n < k$, then $(\mathrm{Conj}(n,z,w))(k) = 0_{\mathbb{C}}$.

The following propositions are true:

(4) $z\,\mathrm{ExpSeq}(n+1) = \frac{z\,\mathrm{ExpSeq}(n)\cdot z}{(n+1)+0i}$ and $z\,\mathrm{ExpSeq}(0) = 1_{\mathbb{C}}$ and $|z\,\mathrm{ExpSeq}(n)| = |z|\,\mathrm{ExpSeq}(n)$.

(5) If $0 < k$, then $(\mathrm{Sift}\, s_1)(k) = s_1(k -' 1)$.

(6) $(\sum_{\alpha=0}^{\kappa}(s_1)(\alpha))_{\kappa\in\mathbb{N}}(k) = (\sum_{\alpha=0}^{\kappa}(\mathrm{Sift}\, s_1)(\alpha))_{\kappa\in\mathbb{N}}(k) + s_1(k)$.

(7) $(z+w)_{\mathbb{N}}^{n} = (\sum_{\alpha=0}^{\kappa}(\mathrm{Expan}(n,z,w))(\alpha))_{\kappa\in\mathbb{N}}(n)$.

(8) $\mathrm{Expan\_e}(n,z,w) = \frac{1_{\mathbb{C}}}{n!_{\mathbb{C}}}\,\mathrm{Expan}(n,z,w)$.

(9) $\frac{(z+w)_{\mathbb{N}}^{n}}{n!_{\mathbb{C}}} = (\sum_{\alpha=0}^{\kappa}(\mathrm{Expan\_e}(n,z,w))(\alpha))_{\kappa\in\mathbb{N}}(n)$.

(10) $0_{\mathbb{C}}\,\mathrm{ExpSeq}$ is absolutely summable and $\sum(0_{\mathbb{C}}\,\mathrm{ExpSeq}) = 1_{\mathbb{C}}$.

Let us consider z. Note that $z\,\mathrm{ExpSeq}$ is absolutely summable.
One can prove the following propositions:

(11) $z\,\mathrm{ExpSeq}(0) = 1_{\mathbb{C}}$ and $(\mathrm{Expan}(0,z,w))(0) = 1_{\mathbb{C}}$.

(12) If $l \leq k$, then $(\mathrm{Alfa}(k+1,z,w))(l) = (\mathrm{Alfa}(k,z,w))(l) + (\mathrm{Expan\_e}(k+1,z,w))(l)$.

(13) $(\sum_{\alpha=0}^{\kappa}(\mathrm{Alfa}(k+1,z,w))(\alpha))_{\kappa\in\mathbb{N}}(k) = (\sum_{\alpha=0}^{\kappa}(\mathrm{Alfa}(k,z,w))(\alpha))_{\kappa\in\mathbb{N}}(k) + (\sum_{\alpha=0}^{\kappa}(\mathrm{Expan\_e}(k+1,z,w))(\alpha))_{\kappa\in\mathbb{N}}(k)$.

(14) $z\,\mathrm{ExpSeq}(k) = (\mathrm{Expan\_e}(k,z,w))(k)$.

(15) $(\sum_{\alpha=0}^{\kappa} z+w\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n) = (\sum_{\alpha=0}^{\kappa}(\mathrm{Alfa}(n,z,w))(\alpha))_{\kappa\in\mathbb{N}}(n)$.

(16) $(\sum_{\alpha=0}^{\kappa} z\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(k) \cdot (\sum_{\alpha=0}^{\kappa} w\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(k) - (\sum_{\alpha=0}^{\kappa} z+w\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(k) = (\sum_{\alpha=0}^{\kappa}(\mathrm{Conj}(k,z,w))(\alpha))_{\kappa\in\mathbb{N}}(k)$.

(17) $|(\sum_{\alpha=0}^{\kappa} z\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(k)| \leq (\sum_{\alpha=0}^{\kappa}|z|\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(k)$ and $(\sum_{\alpha=0}^{\kappa}|z|\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(k) \leq \sum(|z|\,\mathrm{ExpSeq})$ and $|(\sum_{\alpha=0}^{\kappa} z\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(k)| \leq \sum(|z|\,\mathrm{ExpSeq})$.

(18) $1 \leq \sum(|z|\,\mathrm{ExpSeq})$.

(19) $0 \leq |z|\,\mathrm{ExpSeq}(n)$.

(20) $|(\sum_{\alpha=0}^{\kappa}|z|\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n)| = (\sum_{\alpha=0}^{\kappa}|z|\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n)$ and if $n \leq m$, then $|(\sum_{\alpha=0}^{\kappa}|z|\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(m) - (\sum_{\alpha=0}^{\kappa}|z|\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n)| = (\sum_{\alpha=0}^{\kappa}|z|\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(m) - (\sum_{\alpha=0}^{\kappa}|z|\,\mathrm{ExpSeq}(\alpha))_{\kappa\in\mathbb{N}}(n)$.

(21) $|(\sum_{\alpha=0}^{\kappa}|\mathrm{Conj}(k,z,w)|(\alpha))_{\kappa\in\mathbb{N}}(n)| = (\sum_{\alpha=0}^{\kappa}|\mathrm{Conj}(k,z,w)|(\alpha))_{\kappa\in\mathbb{N}}(n)$.

(22) For every real number p such that $p > 0$ there exists n such that for every k such that $n \leq k$ holds $|(\sum_{\alpha=0}^{\kappa}|\mathrm{Conj}(k,z,w)|(\alpha))_{\kappa\in\mathbb{N}}(k)| < p$.

(23) For every s_1 such that for every k holds $s_1(k) = (\sum_{\alpha=0}^{\kappa}(\mathrm{Conj}(k,z,w))(\alpha))_{\kappa\in\mathbb{N}}(k)$ holds s_1 is convergent and $\lim s_1 = 0_{\mathbb{C}}$.

2. DEFINITION OF EXPONENTIAL FUNCTION ON COMPLEX

The partial function exp from $\mathbb{C}$ to $\mathbb{C}$ is defined as follows:

(Def. 18) $\mathrm{dom}\exp = \mathbb{C}$ and for every element z of $\mathbb{C}$ holds $\exp(z) = \sum(z\,\mathrm{ExpSeq})$.

Let us consider z. The functor $\exp z$ yields an element of $\mathbb{C}$ and is defined as follows:

(Def. 19) $\exp z = \exp(z)$.

The following proposition is true

(24) For all z_1, z_2 holds $\exp(z_1+z_2) = \exp z_1 \cdot \exp z_2$.

3. DEFINITION OF SINUS, COSINE, AND EXPONENTIAL FUNCTION ON $\mathbb{R}$

The partial function sin from $\mathbb{R}$ to $\mathbb{R}$ is defined as follows:

(Def. 20) $\mathrm{dom}\sin = \mathbb{R}$ and for every element d of $\mathbb{R}$ holds $\sin(d) = \Im(\sum(0+di\,\mathrm{ExpSeq}))$.

Let t_1 be a real number. The functor $\sin t_1$ is defined as follows:

(Def. 21) $\sin t_1 = \sin(t_1)$.

Let t_1 be a real number. One can check that $\sin t_1$ is real.
Let t_1 be a real number. Then $\sin t_1$ is a real number.
Next we state the proposition

(25) sin is a function from $\mathbb{R}$ into $\mathbb{R}$.

In the sequel d denotes a real number.
The partial function cos from $\mathbb{R}$ to $\mathbb{R}$ is defined as follows:

(Def. 22) $\mathrm{dom}\cos = \mathbb{R}$ and for every d holds $\cos(d) = \Re(\sum(0+di\,\mathrm{ExpSeq}))$.

Let t_1 be a real number. The functor $\cos t_1$ is defined by:

(Def. 23) $\cos t_1 = \cos(t_1)$.

Let t_1 be a real number. Note that $\cos t_1$ is real.
Let t_1 be a real number. Then $\cos t_1$ is a real number.
Next we state several propositions:

(26) cos is a function from $\mathbb{R}$ into $\mathbb{R}$.

(27) $\mathrm{dom}\sin = \mathbb{R}$ and $\mathrm{dom}\cos = \mathbb{R}$.

(28) $\exp(0+t_1 i) = \cos t_1 + \sin t_1 i$.

(29) $\overline{\exp(0+t_1 i)} = \exp(-(0+t_1 i))$.

(30) $|\exp(0+t_1 i)| = 1$ and $|\sin t_1| \leq 1$ and $|\cos t_1| \leq 1$.

(31) $\cos(t_1)^{\mathbf{2}} + \sin(t_1)^{\mathbf{2}} = 1$ and $\cos(t_1)\cdot\cos(t_1) + \sin(t_1)\cdot\sin(t_1) = 1$.

(32) $(\cos t_1)^{\mathbf{2}} + (\sin t_1)^{\mathbf{2}} = 1$ and $\cos t_1 \cdot \cos t_1 + \sin t_1 \cdot \sin t_1 = 1$.

(33) $\cos(0) = 1$ and $\sin(0) = 0$ and $\cos(-t_1) = \cos(t_1)$ and $\sin(-t_1) = -\sin(t_1)$.

(34) $\cos 0 = 1$ and $\sin 0 = 0$ and $\cos(-t_1) = \cos t_1$ and $\sin(-t_1) = -\sin t_1$.

Let t_1 be a real number. The functor t_1 P_sin yielding a sequence of real numbers is defined by:

(Def. 24) For every n holds $t_1\,\mathrm{P\_sin}(n) = \frac{(-1)^n \cdot {t_1}^{2\cdot n+1}}{(2\cdot n+1)!}$.

The functor t_1 P_cos yields a sequence of real numbers and is defined by:

(Def. 25) For every n holds $t_1\,\mathrm{P\_cos}(n) = \frac{(-1)^n \cdot {t_1}^{2\cdot n}}{(2\cdot n)!}$.

Next we state a number of propositions:

(35) For all z, k holds $z^{2\cdot k}_{\mathbb{N}} = (z^{k}_{\mathbb{N}})^{2}_{\mathbb{N}}$ and $z^{2\cdot k}_{\mathbb{N}} = (z^{2}_{\mathbb{N}})^{k}_{\mathbb{N}}$.

(36) For all k, t_1 holds $(0+t_1 i)^{2\cdot k}_{\mathbb{N}} = (-1)^k \cdot {t_1}^{2\cdot k} + 0i$ and $(0+t_1 i)^{2\cdot k+1}_{\mathbb{N}} = 0 + ((-1)^k \cdot {t_1}^{2\cdot k+1})i$.

(37) For every n holds $n!_{\mathbb{C}} = n! + 0i$.

(38) For all t_1, n holds $(\sum_{\alpha=0}^{\kappa} t_1\,\mathrm{P\_sin}(\alpha))_{\kappa\in\mathbb{N}}(n) = (\sum_{\alpha=0}^{\kappa} \Im(0+t_1 i\,\mathrm{ExpSeq})(\alpha))_{\kappa\in\mathbb{N}}(2\cdot n+1)$ and $(\sum_{\alpha=0}^{\kappa} t_1\,\mathrm{P\_cos}(\alpha))_{\kappa\in\mathbb{N}}(n) = (\sum_{\alpha=0}^{\kappa} \Re(0+t_1 i\,\mathrm{ExpSeq})(\alpha))_{\kappa\in\mathbb{N}}(2\cdot n)$.

(39) For every t_1 holds $(\sum_{\alpha=0}^{\kappa} t_1\,\mathrm{P\_sin}(\alpha))_{\kappa\in\mathbb{N}}$ is convergent and $\sum(t_1\,\mathrm{P\_sin}) = \Im(\sum(0+t_1 i\,\mathrm{ExpSeq}))$ and $(\sum_{\alpha=0}^{\kappa} t_1\,\mathrm{P\_cos}(\alpha))_{\kappa\in\mathbb{N}}$ is convergent and $\sum(t_1\,\mathrm{P\_cos}) = \Re(\sum(0+t_1 i\,\mathrm{ExpSeq}))$.

(40) For every t_1 holds $\cos(t_1) = \sum(t_1\,\mathrm{P\_cos})$ and $\sin(t_1) = \sum(t_1\,\mathrm{P\_sin})$.

(41) For all p, t_1, r_1 such that r_1 is convergent and $\lim r_1 = t_1$ and for every n holds $r_1(n) \geq p$ holds $t_1 \geq p$.

(42) For all n, k, m such that $n < k$ holds $m! > 0$ and $n! \leq k!$.

(43) For all t_1, n, k such that $0 \leq t_1$ and $t_1 \leq 1$ and $n \leq k$ holds ${t_1}^k \leq {t_1}^n$.

(44) For all t_1, n holds $(t_1+0i)^{n}_{\mathbb{N}} = {t_1}^n + 0i$.

(45) For all t_1, n holds $\frac{(t_1+0i)^{n}_{\mathbb{N}}}{n!_{\mathbb{C}}} = \frac{{t_1}^n}{n!} + 0i$.

(46) $\Im(\sum(p+0i\,\mathrm{ExpSeq})) = 0$.

(47) $\cos(1) > 0$ and $\sin(1) > 0$ and $\cos(1) < \sin(1)$.

(48) For every t_1 holds $t_1\,\mathrm{ExpSeq} = \Re(t_1+0i\,\mathrm{ExpSeq})$.

(49) For every t_1 holds $t_1\,\mathrm{ExpSeq}$ is summable and $\sum(t_1\,\mathrm{ExpSeq}) = \Re(\sum(t_1+0i\,\mathrm{ExpSeq}))$.

(50) For all p, q holds $\sum(p+q\,\mathrm{ExpSeq}) = \sum(p\,\mathrm{ExpSeq}) \cdot \sum(q\,\mathrm{ExpSeq})$.

The partial function exp from $\mathbb{R}$ to $\mathbb{R}$ is defined as follows:

(Def. 26) $\mathrm{dom}\exp = \mathbb{R}$ and for every real number d holds $\exp(d) = \sum(d\,\mathrm{ExpSeq})$.

Let t_1 be a real number. The functor $\exp t_1$ is defined as follows:

(Def. 27) $\exp t_1 = \exp(t_1)$.

Let t_1 be a real number. One can check that $\exp t_1$ is real.
Let t_1 be a real number. Then $\exp t_1$ is a real number.
The following propositions are true:

(51) $\mathrm{dom}\exp = \mathbb{R}$.

(53)[2] For every t_1 holds $\exp(t_1) = \Re(\sum(t_1 + 0i\,\mathrm{ExpSeq}))$.

(54) $\exp(t_1 + 0i) = \exp t_1 + 0i$.

(55) $\exp(p+q) = \exp p \cdot \exp q$.

(56) $\exp 0 = 1$.

(57) For every t_1 such that $t_1 > 0$ holds $\exp(t_1) \geq 1$.

(58) For every t_1 such that $t_1 < 0$ holds $0 < \exp(t_1)$ and $\exp(t_1) \leq 1$.

(59) For every t_1 holds $\exp(t_1) > 0$.

(60) For every t_1 holds $\exp t_1 > 0$.

4. Differential of Sinus, Cosine, and Exponential Function

Let z be an element of $\mathbb{C}$. The functor zP_dt yielding a complex sequence is defined as follows:

(Def. 28) For every n holds $z\mathrm{P\_dt}(n) = \frac{z_{\mathbb{N}}^{n+1}}{(n+2)!_{\mathbb{C}}}$.

The functor zP_t yielding a complex sequence is defined as follows:

(Def. 29) For every n holds $z\mathrm{P\_t}(n) = \frac{z_{\mathbb{N}}^{n}}{(n+2)!_{\mathbb{C}}}$.

Next we state a number of propositions:

(61) For every z holds zP_dt is absolutely summable.

(62) For every z holds $z \cdot \sum(z\mathrm{P\_dt}) = \sum(z\,\mathrm{ExpSeq}) - 1_{\mathbb{C}} - z$.

(63) For every p such that $p > 0$ there exists q such that $q > 0$ and for every z such that $|z| < q$ holds $|\sum(z\mathrm{P\_dt})| < p$.

(64) For all z, z_1 holds $\sum(z_1 + z\,\mathrm{ExpSeq}) - \sum(z_1\,\mathrm{ExpSeq}) = \sum(z_1\,\mathrm{ExpSeq}) \cdot z + z \cdot \sum(z\mathrm{P\_dt}) \cdot \sum(z_1\,\mathrm{ExpSeq})$.

(65) For all p, q holds $\cos(p+q) - \cos(p) = -q \cdot \sin(p) - q \cdot \Im(\sum(0 + qi\mathrm{P\_dt}) \cdot (\cos(p) + \sin(p)i))$.

(66) For all p, q holds $\sin(p+q) - \sin(p) = q \cdot \cos(p) + q \cdot \Re(\sum(0 + qi\mathrm{P\_dt}) \cdot (\cos(p) + \sin(p)i))$.

(67) For all p, q holds $\exp(p+q) - \exp(p) = q \cdot \exp(p) + q \cdot \exp(p) \cdot \Re(\sum(q + 0i\mathrm{P\_dt}))$.

(68) For every p holds cos is differentiable in p and $\cos'(p) = -\sin(p)$.

(69) For every p holds sin is differentiable in p and $\sin'(p) = \cos(p)$.

(70) For every p holds exp is differentiable in p and $\exp'(p) = \exp(p)$.

(71) exp is differentiable on $\mathbb{R}$ and for every t_1 such that $t_1 \in \mathbb{R}$ holds $\exp'(t_1) = \exp(t_1)$.

(72) cos is differentiable on $\mathbb{R}$ and for every t_1 such that $t_1 \in \mathbb{R}$ holds $\cos'(t_1) = -\sin(t_1)$.

(73) sin is differentiable on $\mathbb{R}$ and for every t_1 holds $\sin'(t_1) = \cos(t_1)$.

(74) For every t_1 such that $t_1 \in [0,1]$ holds $0 < \cos(t_1)$ and $\cos(t_1) \geq \frac{1}{2}$.

(75) $[0,1] \subseteq \mathrm{dom}(\frac{\sin}{\cos})$ and $]0,1[\subseteq \mathrm{dom}(\frac{\sin}{\cos})$.

(76) $\frac{\sin}{\cos}$ is continuous on $[0,1]$.

(77) For all t_2, t_3 such that $t_2 \in]0,1[$ and $t_3 \in]0,1[$ and $(\frac{\sin}{\cos})(t_2) = (\frac{\sin}{\cos})(t_3)$ holds $t_2 = t_3$.

[2] The proposition (52) has been removed.

5. EXISTENCE OF CIRCLE RATIO

The real number π is defined as follows:

(Def. 30) $(\frac{\sin}{\cos})(\frac{\pi}{4}) = 1$ and $\pi \in]0,4[$.

π is a real number.
The following proposition is true

(78) $\sin(\frac{\pi}{4}) = \cos(\frac{\pi}{4})$.

6. FORMULAS OF SINUS, COSINE

One can prove the following propositions:

(79) $\sin(t_2+t_3) = \sin(t_2)\cdot\cos(t_3)+\cos(t_2)\cdot\sin(t_3)$ and $\cos(t_2+t_3) = \cos(t_2)\cdot\cos(t_3)-\sin(t_2)\cdot\sin(t_3)$.

(80) $\sin(t_2+t_3) = \sin t_2\cdot\cos t_3+\cos t_2\cdot\sin t_3$ and $\cos(t_2+t_3) = \cos t_2\cdot\cos t_3-\sin t_2\cdot\sin t_3$.

(81) $\cos(\frac{\pi}{2}) = 0$ and $\sin(\frac{\pi}{2}) = 1$ and $\cos(\pi) = -1$ and $\sin(\pi) = 0$ and $\cos(\pi+\frac{\pi}{2}) = 0$ and $\sin(\pi+\frac{\pi}{2}) = -1$ and $\cos(2\cdot\pi) = 1$ and $\sin(2\cdot\pi) = 0$.

(82) $\cos(\frac{\pi}{2}) = 0$ and $\sin(\frac{\pi}{2}) = 1$ and $\cos\pi = -1$ and $\sin\pi = 0$ and $\cos(\pi+\frac{\pi}{2}) = 0$ and $\sin(\pi+\frac{\pi}{2}) = -1$ and $\cos(2\cdot\pi) = 1$ and $\sin(2\cdot\pi) = 0$.

(83) $\sin(t_1+2\cdot\pi) = \sin(t_1)$ and $\cos(t_1+2\cdot\pi) = \cos(t_1)$ and $\sin(\frac{\pi}{2}-t_1) = \cos(t_1)$ and $\cos(\frac{\pi}{2}-t_1) = \sin(t_1)$ and $\sin(\frac{\pi}{2}+t_1) = \cos(t_1)$ and $\cos(\frac{\pi}{2}+t_1) = -\sin(t_1)$ and $\sin(\pi+t_1) = -\sin(t_1)$ and $\cos(\pi+t_1) = -\cos(t_1)$.

(84) $\sin(t_1+2\cdot\pi) = \sin t_1$ and $\cos(t_1+2\cdot\pi) = \cos t_1$ and $\sin(\frac{\pi}{2}-t_1) = \cos t_1$ and $\cos(\frac{\pi}{2}-t_1) = \sin t_1$ and $\sin(\frac{\pi}{2}+t_1) = \cos t_1$ and $\cos(\frac{\pi}{2}+t_1) = -\sin t_1$ and $\sin(\pi+t_1) = -\sin t_1$ and $\cos(\pi+t_1) = -\cos t_1$.

(85) For every t_1 such that $t_1 \in]0,\frac{\pi}{2}[$ holds $\cos(t_1) > 0$.

(86) For every t_1 such that $t_1 \in]0,\frac{\pi}{2}[$ holds $\cos t_1 > 0$.

REFERENCES

[1] Agnieszka Banachowicz and Anna Winnicka. Complex sequences. *Journal of Formalized Mathematics*, 5, 1993. `http://mizar.org/JFM/Vol5/comseq_1.html`.

[2] Grzegorz Bancerek. The fundamental properties of natural numbers. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/nat_1.html`.

[3] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/ordinal1.html`.

[4] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/funct_1.html`.

[5] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/funct_2.html`.

[6] Czesław Byliński. The complex numbers. *Journal of Formalized Mathematics*, 2, 1990. `http://mizar.org/JFM/Vol2/complex1.html`.

[7] Library Committee. Introduction to arithmetic. *Journal of Formalized Mathematics*, Addenda, 2003. `http://mizar.org/JFM/Addenda/arytm_0.html`.

[8] Krzysztof Hryniewiecki. Basic properties of real numbers. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/real_1.html`.

[9] Jarosław Kotowicz. Convergent sequences and the limit of sequences. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/seq_2.html`.

[10] Jarosław Kotowicz. Monotone real sequences. Subsequences. *Journal of Formalized Mathematics*, 1, 1989. `http://mizar.org/JFM/Vol1/seqm_3.html`.

[11] Jarosław Kotowicz. Real sequences and basic operations on them. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/seq_1.html.

[12] Jarosław Kotowicz. Partial functions from a domain to the set of real numbers. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/rfunct_1.html.

[13] Rafał Kwiatek. Factorial and Newton coefficients. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/newton.html.

[14] Adam Naumowicz. Conjugate sequences, bounded complex sequences and convergent complex sequences. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/comseq_2.html.

[15] Takaya Nishiyama and Yasuho Mizuhara. Binary arithmetics. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/binarith.html.

[16] Jan Popiołek. Some properties of functions modul and signum. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/absvalue.html.

[17] Konrad Raczkowski. Integer and rational exponents. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/prepower.html.

[18] Konrad Raczkowski and Andrzej Nędzusiak. Series. *Journal of Formalized Mathematics*, 3, 1991. http://mizar.org/JFM/Vol3/series_1.html.

[19] Konrad Raczkowski and Paweł Sadowski. Real function continuity. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/fcont_1.html.

[20] Konrad Raczkowski and Paweł Sadowski. Real function differentiability. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/fdiff_1.html.

[21] Konrad Raczkowski and Paweł Sadowski. Topological properties of subsets in real numbers. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/rcomp_1.html.

[22] Yasunari Shidama and Artur Korniłowicz. Convergence and the limit of complex sequences. Series. *Journal of Formalized Mathematics*, 9, 1997. http://mizar.org/JFM/Vol9/comseq_3.html.

[23] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.

[24] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. http://mizar.org/JFM/Addenda/numbers.html.

[25] Andrzej Trybulec and Czesław Byliński. Some properties of real numbers operations: min, max, square, and square root. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/square_1.html.

[26] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

[27] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

Received October 22, 1998

Published January 2, 2004